

The spectroscopic foundation of radiative forcing of climate by carbon dioxide

Marty Mlynczak
NASA Langley Research Center
&
Co-Authors

Co-Authors

- NASA Langley
 - Taumi Daniels
 - David Kratz
 - Jeffrey Mast
 - Linda Hunt
- AER
 - Eli Mlawer
 - Matthew Alvarado
- Lawrence Berkeley Lab
 - William Collins
 - Daniel Feldman
- NOAA ESRL
 - David Fahey
- University of Wisconsin
 - Wilmer Anderson
 - James Lawler

Outline

- **Motivation**
- **Definition and Examples of Radiative Forcing**
- **Uncertainty in RF due to Spectroscopic Parameters**
 - Line shape and line mixing
 - Line Strengths
 - Air-broadened halfwidths
- **Results**
- **Utility of the Voigt lineshape**
- **Summary and Conclusions**

Motivation

- Radiative forcing (RF) by CO₂ is the leading contribution to anthropogenic climate change
- Uncertainties in CO₂ RF impact scientific and policy assessments
- Goals of this work:
 - Assess uncertainty in RF associated with infrared spectroscopy of CO₂
 - Of particular interest is line mixing and line shape function
 - Refute recent assertions that RF is greatly overestimated due to inappropriate use of Voigt lineshape (Happer, 2014)
- **Result: RF spectroscopic uncertainty is < 1% and the foundation of climate change modeling is sound**

International Journal of Modern Physics A
Vol. 29, No. 7 (2014) 1460003 (34 pages)
© World Scientific Publishing Company
DOI: 10.1142/S0217751X14600033



Why has global warming paused?*

William Happer

Department of Physics, Princeton University, Princeton, NJ 08544, USA
happer@princeton.edu

Received 29 November 2013


Revised 10 February 2014

Accepted 13 February 2014

Published 11 March 2014


Definition of Radiative Forcing (RF) - 1

- RF is the change in the net radiative flux at the tropopause
- Net flux is defined as $F(\text{down})$ minus $F(\text{up})$



Definition of Radiative Forcing (RF) - 2


- Change in net flux is difference for two different CO₂ burdens, typically doubled (D) from pre-industrial (P) minus P



Use LBLRTM v12.2 to model radiances and fluxes in computation of RF

The Spectrum of Radiative Forcing for 2 x CO₂


Instantaneous Spectral Radiative Forcing by Carbon Dioxide



Baseline Case, Mid-Latitude Summer

The Spectrum of Radiative Forcing for 2 x CO₂

Instantaneous Spectral Radiative Forcing by Carbon Dioxide



Baseline Case, Sub-Arctic Winter

Spectroscopic Uncertainty - Methodology

- Use LBLRTM v12.2 to evaluate up and down radiances/fluxes
- Compute instantaneous radiative forcing
- Only uncertainty in the transmittance function is considered:

$$T_\nu(z, z') = \exp\left(-\sum_i S_i(\Theta) g_i(\nu - \nu_0) \frac{u(z, z')}{\mu}\right)$$

- Line shape function $g(\nu - \nu_0)$
 - Line strength $S_i(\Theta)$
 - Air-broadened halfwidth (in lineshape function)
- Perturb these parameters to assess uncertainty in spectroscopy; compute difference with “baseline” case

Uncertainties in Spectroscopy – Line Shape/Mixing

- Classically, the Voigt lineshape is used to represent the effects of collision/Lorentz and Doppler broadening
- Wings of many IR-active species are sub- or super-Lorentzian
- The 15- μm bands of CO_2 , lines are densely packed and cannot be considered isolated during collisions
- Line mixing occurs, resulting in additional absorption near line centers and sub-Lorentzian behavior in the line wings
 - There is a 50-year history of modeling sub-Lorentzian wings in CO_2
- LBLRTM adopts line mixing from Niro et al. (2005)
 - Replaces classic line shape function in transmittance calculation

Approach to Computing Spectroscopic Uncertainty


- **Uncertainty determined by perturbation analysis**
- **Compute RF for “Baseline” case**
- **Compute RF for “Perturbed” case**
 - Increase line mixing by 20%
 - Perturb line strengths by assigned uncertainty on AER v3.2 database
 - Perturb line halfwidths by assigned uncertainty on AER v3.2 database
- **Compute Uncertainty = Perturbed minus Baseline**
 - Spectra, and spectrally integrated differences
 - RSS all uncertainties to get total uncertainty

Uncertainty in Radiative Forcing due to Line Mixing

RF (increased line mixing) – RF (baseline)

Mid-Latitude Summer Atmosphere

Line Mixing Uncertainty in Radiative Forcing



Significant but compensating spectral structure


RF difference is 0.016 W/m² or 0.3% of RF baseline

Uncertainty in Radiative Forcing due to Line Mixing

RF (increased line mixing) – RF (baseline)

Sub-Arctic Winter Atmosphere


Line Mixing Uncertainty in Radiative Forcing



Significant but compensating spectral structure

RF difference is 0.022 W/m² or 0.6% of RF baseline

Radiative Forcing Uncertainty due to Line Strength



Virtually no change in RF with up to 20% increase/decrease in S

Summary of Spectroscopic Uncertainty in RF

<u>Atmosphere</u>	Line Shape Uncertainty W/m ²	Line Strength Uncertainty W/m ²	Halfwidth Uncertainty W/m ²	RSS W/m ²	Error as % of Baseline RF
Mid-Latitude Summer	0.016	0.015	0.005	0.022	0.41
Mid-Latitude Winter	0.016	0.010	0.004	0.019	0.44
Sub-Arctic Summer	0.023	0.009	0.007	0.028	0.55
Sub-Arctic Winter	0.022	0.015	0.006	0.025	0.68
Tropical	0.010	0.014	0.002	0.017	0.51


Spectroscopic Uncertainty in RF is < 0.7% of Forcing in a Variety of Atmospheres




Summary

- **We have examined uncertainty in radiative forcing by CO₂ associated with spectroscopic parameters**
- **Combined uncertainty of line shape, line strength, and halfwidth is < 0.7% for a variety of standard atmospheres**
- **Line mixing is rigorously included in state of the art line-by-line models and in rapid codes used in climate models**
- **Voigt shape provides good agreement at 50 cm⁻¹ wing truncation, but does overestimate at larger truncations**
- **Foundation of climate modeling is robust with regards to uncertainties in spectroscopy**

Backup Slides

Radiative Forcing vs. Voigt Truncation Offset







Summary of Spectroscopic Uncertainty in RF


Atmosphere	RF LBLRTM Baseline W/m ²	RF LBLRTM 20% Pert. W/m ²	Uncertainty Line Shape/Line Mixing W/m ²	Uncertainty Line Strength W/m ²	Uncertainty Half Width W/m ²	RSS W/m ²	RSS as Percent of baseline LBLRTM Forcing
MLS	5.428	5.444	0.016	0.015	0.005	0.022	0.41
MLW	4.372	4.388	0.016	0.010	0.004	0.019	0.44
SAS	5.118	5.141	0.023	0.015	0.007	0.028	0.55
SAW	3.606	3.628	0.022	0.009	0.006	0.025	0.68
TRO	5.509	5.519	0.010	0.014	0.002	0.017	0.31

Instantaneous Spectral Radiative Forcing by Carbon Dioxide




LBLRTM Radiative Forcing






The Spectrum of Radiative Forcing

Instantaneous Spectral Radiative Forcing by Carbon Dioxide




The Spectrum of Radiative Forcing

Instantaneous Spectral Radiative Forcing by Carbon Dioxide




The Spectrum of Radiative Forcing


Instantaneous Spectral Radiative Forcing by Carbon Dioxide




Line Mixing Uncertainty in Radiative Forcing




Line Mixing Uncertainty in Radiative Forcing



Line Mixing Uncertainty in Radiative Forcing




Radiative Forcing vs. Voigt Truncation Offset



Definition of Radiative Forcing (RF)

- RF is the change in the net radiative flux at the tropopause
- Net flux is defined as $F(\text{down})$ minus $F(\text{up})$
- Change in net flux is difference for two different CO_2 burdens, typically doubled from pre-industrial (PI) minus PI



Methodology

- Use LBLRTM v12.2 to evaluate up and down radiances/fluxes

$$I_{\nu}(\mu, z) = B_{\nu}(\Theta_s)T_{\nu}(z, 0) + \int_0^z B_{\nu}(\Theta(z')) \frac{\partial T_{\nu}(z, z')}{\partial z'} dz'$$

$$I_{\nu}(-\mu, z) = - \int_z^{\infty} B_{\nu}(\Theta(z')) \frac{\partial T_{\nu}(z, z')}{\partial z'} dz'$$

- The key to our study is the transmittance function:

$$T_{\nu}(z, z') = \exp\left(- \sum_i S_i(\Theta) g_i(\nu - \nu_0) \frac{u(z, z')}{\mu}\right)$$

- Compute instantaneous forcing, no change in temperature
- Vary $g(\nu - \nu_0)$, S_i , and α_L to assess spectroscopic uncertainty in RF

Effect of Line Mixing

