

## ORGANIC AGGREGATES WITH $\delta D$ AND $\delta^{15}N$ ANOMALIES IN THE ZAG CLAST REVEALED BY STXM AND NANOSIMS.

Y. Kebukawa<sup>1</sup>, M. Ito<sup>2</sup>, M. E. Zolensky<sup>3</sup>, Z. Rahman<sup>4</sup>, A. L. D. Kilcoyne<sup>5</sup>, A. Nakato<sup>6</sup>, Y. Takeichi<sup>7</sup>, H. Suga<sup>8</sup>, C. Miyamoto<sup>9</sup>, K. Mase<sup>7</sup>, Y. Takahashi<sup>9</sup>, Q. Chan<sup>3</sup> and M. Fries<sup>3</sup>. <sup>1</sup>Faculty of Engineering, Yokohama National University (kebukawa@ynu.ac.jp), <sup>2</sup>Kochi Institute for Core Sample Research, JAMSTEC, <sup>3</sup>ARES, NASA Johnson Space Center, <sup>4</sup>Jacobs, NASA Johnson Space Center, <sup>5</sup>Advanced Light Source, Lawrence Berkeley National Laboratory, <sup>6</sup>Graduate School of Science, Kyoto University, <sup>7</sup>Institute of Materials Structure Science, High-Energy Accelerator Research Organization (KEK), <sup>8</sup>Department of Earth and Planetary Systems Science, Hiroshima University, <sup>9</sup>Department of Earth and Planetary Science, The University of Tokyo.

**Introduction:** Xenolithic clasts are often found in a wide variety of meteorite groups [e.g., 1]. Some ordinary chondrite clasts are interesting since these clasts might have originated from Ceres which shares crossing orbits with a possible ordinary chondrite parent body, Hebe [2]. The Zag meteorite contains a dark clast dominated by saponite, serpentine, carbonates, sulfides, magnetite, minor olivine and pyroxene, which is consistent with formation on a large, carbonaceous, aqueously active body, e.g., Ceres [3]. Abundant large C-rich grains up to 20  $\mu m$  were found in the Zag clast as well [3]. Such large C-rich grains are unique among any other meteorites in our knowledge, and will provide important clues to decipher the origin of the clast and accretion history.

**Methods:** C-rich grains were selected in the Zag dark clast using SEM and approximately 100 nm-thick sections were prepared using a focused ion beam (FIB) at NASA-JSC. The sections were analyzed using the scanning transmission X-ray microscope (STXM) on beamline 5.3.2.2 at Advanced Light Source, LBNL, and BL-13A at the Photon Factory, KEK. Subsequently, the FIB section was analyzed for H, C and N isotopic compositions using a CAMECA NanoSIMS 50L ion microprobe at Kochi Institute for Core Sample Research, JAMSTEC.

**Results and Discussion:** The FIB section showed a large C-rich area over 10  $\mu m$  width that corresponded to the C-rich grain (Fig. 1b). Carbon X-ray absorption near edge structure (XANES) of the C-rich grain showed a peak at 284.8 eV that is assigned to  $sp^2$  carbon (C=C) (Fig. 1c,d in Red). The surrounded matrix area showed a peak at 290.3 eV that is assigned to carbonates with smaller 284.8 eV peak. The C-XANES of the C-rich grain had no other functional groups (e.g., C=O) that are characteristic of primitive chondritic insoluble organic matter (IOM) (e.g., Murchison), nor  $1s-\sigma^*$  exciton at 291.7 eV of graphene structures that is characteristic of thermally metamorphosed meteorites (e.g., Allende IOM) [4]. No specific nitrogen features were observed in N-XANES. XANES results indicate that the C-rich grain may consist of a hydrogenated amorphous carbon-like structure.

The C-rich grain had large  $\delta D$  and  $\delta^{15}N$  anomaly,  $\delta D = 2,370 \pm 74 \text{ ‰}$  and  $\delta^{15}N = 696 \pm 100 \text{ ‰}$  in average.  $\delta^{13}C$  was  $-43 \pm 20 \text{ ‰}$  that was broadly consistent with the values of CRs and Bells within the analytical error [5]. Two hot spots were observed; one is D- and  $^{15}N$ -rich ( $\delta D = 4,200 \pm 550 \text{ ‰}$  and  $\delta^{15}N = 3,413 \pm 1,070 \text{ ‰}$ ), and the other is D-rich ( $\delta D = 4,500 \pm 900 \text{ ‰}$ ) with normal to moderate enrichment in  $^{15}N$  ( $724 \pm 780 \text{ ‰}$ ) (Fig. 1e,f). The origin of the C-rich grain and isotope anomaly is puzzling since no molecular heterogeneity was observed between the hot spots and the average area. Some “aromatic” globules in chondrites reported by [6] show C-XANES spectra similar to the C-rich grains in the Zag clast. In their study however, no clear correlation between molecular structure and  $\delta^{15}N$  was observed; aromatic globules have similar or higher  $\delta^{15}N$  values than IOM-like globules [6]. If the C-rich grain is an assemblage of organic matter with different origins (with different  $\delta D$  and  $\delta^{15}N$  values), isotope heterogeneity might have survived the structural homogenization that possibly occurred during subsequent aqueous processing.

**References:** [1] Rubin A. E. et al. 2002. *Meteoritics & Planetary Science* 37:125–141. [2] Fries M. et al. 2013. Abstract #5266. 76th Annual Meeting of The Meteoritical Society. [3] Zolensky M. E. et al. 2015. Abstract #5270. 78th Annual Meeting of The Meteoritical Society. [4] Cody G. D. et al. 2008. *Meteoritics & Planetary Science* 43:353–365. [5] Alexander C. M. O’D. et al. 2007. *Geochimica et Cosmochimica Acta* 71:4380–4403. [6] De Gregorio B. T. et al. 2013. *Meteoritics & Planetary Science* 48:904–928.




Fig. 1: A FIB section from Zag clast C-rich grain (a) BSE image, (b) C-map obtained with STXM, (c) color coded map with two C-XANES spectra (d), C-rich grain is in red and clast matrix is in green. (e, f)  $\delta D$  and  $\delta^{15}N$  images. Hot spots are indicated by arrows.