



Global Modeling and Assimilation Office

GMAO Office Note No. 11 (Version 1.9)

Soil Moisture Active Passive (SMAP) Mission Level 4 Carbon (L4_C) Product Specification Document

Release Date: 10/31/2015

**Global Modeling and Assimilation Office
Earth Sciences Division
NASA Goddard Space Flight Center
Greenbelt, Maryland 20771**

This page intentionally left blank.

Soil Moisture Active Passive (SMAP) Mission Level 4 Carbon (L4_C) Product Specification Document

Joe Glassy¹, John S. Kimball¹, Lucas Jones¹, Rolf H. Reichle², Joseph V. Ardizzone^{2,3}, Gi-Kong Kim², Robert A. Lucchesi^{2,3}, Edmond B. Smith^{2,3}, and Barry H. Weiss⁴

¹NTSG, University of Montana, Missoula, MT, USA

²NASA Goddard Space Flight Center, Greenbelt, MD, USA

³Science Systems and Applications, Inc., Lanham, MD, USA

⁴Jet Propulsion Laboratory, Pasadena, CA, USA

Document maintained by Rolf Reichle (GMAO)

This document should be cited as:

Glassy, J., J.S. Kimball, L. Jones, R. H., Reichle, R. A. Lucchesi, J. V. Ardizzone, G.-K. Kim, E. B. Smith, and B. H. Weiss, 2015: Soil Moisture Active Passive (SMAP) Mission Level 4 Carbon (L4_C) Product Specification Document. GMAO Office Note No. 11 (Version 1.9), 65 pp, NASA Goddard Space Flight Center, Greenbelt, MD, USA. Available from http://gmao.gsfc.nasa.gov/pubs/office_notes.

Approved by:

Steven Pawson

Date

Head, Global Modeling and Assimilation Office
Code 610.1, NASA GSFC

REVISION HISTORY

Revision	Date	Sections Changed	Reason for Change (ECR #)
1.0	26 February, 2013	All	J. Glassy
1.1	4 March, 2013	Product volume estimates added	J. Glassy
1.2	14 May, 28 May 2013	Added QA outputs partitioned by PFT, and updated narratives for QA and bit-flag scheme.	J. Glassy
1.3	17 July 2013	Refined definition of QA structure, added pft_freq{pft} and pft_mod_cnt{pft} definitions; GEO variables "latitude", "longitude" changed to "cell_lat", "cell_lon for Panoply interoperability	J.Glassy
1.4	23 August, 2013	Corrected several typos in QA field descriptions, and added gpp_method bit field to the QA bitflag variable.	J. Glassy
1.5	12 Sept 2013	Modified gpp_method bit to take on values {0,1,2}, indicating when FPAR climatology used vs FPAR 8-day or NDVI, and, added FT_method bitflag	J. Glassy
1.6	15 April, 2014	Deprecated output of 3km surface_flag variable, due to spatio-temporal incompatibility with 9km variables	J. Glassy
		carbon_model_bitflag variable, FT_method bit flag (at bit 15) definition clarified as 0=used radar F/T, 1=used Tsurf (soil surf temp) driven F/T	J. Glassy
		TBD: pending L2/L3 changes to surface_flag will introduce new Tsurf F/T vs. radar F/T and need for a F/T provenance bit indicating which was used.	J.Glassy

1.6.1	17 June 2014	Changed file name format to adopt L4 wide Science Version ID, and clarified valid_min, valid_max dataset properties	J. Glassy
1.7	23 July 2014	QA bitfield x-x GPP_Method changed to a single bit {0 1} and NDVI_Used {0 1} single bit metric was added , and adjusted metadata table to harmonize with L4_SM	J. Glassy
1.8	24 Oct, 2014	All QA spatial variables now merged into single L4C MDL output granule; added additional ISO metadata "source" sections needed for integration, and added prior missing DomainConsistency, CompletenessOmission	J. Glassy
1.8.1	22 June 2015	Updated EC min, max value documentation	J. Glassy
1.9	31 October 2015	Prefatory material: converted to GMAO Office Note for public beta-release of L4_C data. Revised definition of "Launch Indicator" in file name. Minor text updates throughout.	R. Reichle, J. Glassy, J. Kimball

Table of Contents

1 Contents

1 INTRODUCTION	2-9
1.1. Identification.....	2-9
1.2 Scope.....	2-10
1.3 The SMAP Experiment.....	2-10
1.4 SMAP Data products.....	2-11
Table 1: SMAP Data Products.....	2-12
1.5 Content Overview.....	2-13
Table 2: Plant Function Type (PFT) Classifier Summary.....	2-14
1.6 Related SMAP Project Documents.....	2-14
1.7 Applicable Documents.....	2-15
2 DATA PRODUCT ORGANIZATION	2-16
2.1 File Format.....	2-16
2.2 HDF File Notation.....	2-16
2.3 SMAP File Organization.....	2-17
2.3.1 Structure.....	2-17
2.3.2 Data.....	2-17
2.3.3 Element Types.....	2-17
Table 3: Element type definitions.....	2-18
2.3.4 File Level Metadata.....	2-19
2.3.5 Local Metadata.....	2-19
2.4 Data Definition Standards.....	2-20
Table 4: Definitions of Data element characteristics.....	2-20
2.4.1 Double Precision Time Variables.....	2-21
2.4.2 Array representation.....	2-21
3 INTERFACE CHARACTERISTICS	3-22
3.1 Coordinate Systems.....	3-22
3.2 The Global cylindrical EASE-Grid 2.0.....	3-23
3.3 Fill and Gap Values.....	3-25

Table 5: SMAP data product fill values.	3-26
4 DATA DEFINITION	4-27
4.1 Product Overview.....	4-27
4.1.1 SMAP L4_C Data Product	4-27
Table 6: L4_C Data Product Overview.....	4-28
4.1.2 HDF5 Group GEO Overview.....	4-29
4.1.3 HDF5 Group QA Overview	4-29
4.1.4 L4_C Metadata Overview	4-29
4.2 Data Groups and Volume Estimates.....	4-30
Table 7: L4_C Data Volume Estimates	4-30
4.3 File Naming Conventions.....	4-30
4.4 Metadata Detail.....	4-33
4.4.1 L4_C Mandatory Metadata	4-33
4.4.2 L4_C Product Specific metadata	4-33
Table 8: Granule level metadata in the L4_C data product.....	4-34
4.5 Data Structure.....	4-44
4.6 Element definitions	4-44
4.6.1 NEE Dataset Definition	4-45
Table 9: NEE Group Contents	4-45
4.6.2 GPP Dataset Definition	4-46
Table 10: GPP Group Contents	4-46
4.6.3 Heterotrophic respiration (R _h) Dataset definition.....	4-47
Table 11: Rh Group Contents	4-47
4.6.4 Soil Organic Carbon (SOC) Dataset definition.....	4-48
Table 12: SOC Group Contents	4-48
Table 13: EC -- Environmental Constraints Group Contents.....	4-49
4.6.5 GEO: the Geolocation Group.....	4-49
Table 14: Overview of Geolocation Group: GEO	4-50
4.6.6 QA : the Quality Assurance Group.....	4-50
Table 15: Overview of Quality Group: QA	4-51
4.6.7 The carbon_model_bit-flag	4-52
Table 16: L4_C Quality Control Variable (carbon_model_bitflag) Definition	4-52
4.6.8 The carbon_model_qual – partitioned by PFT	4-55

Table 17: L4_C Quality Control Variable.....	4-55
4.7 L4_C Data Product Volume Estimates	4-55
Table 18: L4_C Data Product Volume Estimates	4-56
5 APPENDICES	5-57
Appendix A: Acronyms and abbreviations	5-57
Appendix B: L4_C Dimensions	5-64
Table 18: Dimensions in the SMAP L4_SM data product.	5-64
Appendix C: L4_C Shapes	5-64
Table 19: Shapes in the SMAP L4_SM data product.....	5-64
Appendix D: L4_C Units	5-64
Table 20: Units in the SMAP L4_C data product.....	5-65

1 Introduction

The SMAP Level 4 Carbon (L4_C) data product is computed using the daily Level 4 Carbon (L4_C) algorithm which produces a number of global ecosystem productivity variables along with several product quality terms. The ecosystem productivity variables calculated within the L4_C daily algorithm are Gross Primary Productivity (GPP), Net Ecosystem Exchange (NEE), Heterotrophic respiration (Rh), and surface Soil Organic Carbon (SOC). In addition, several environment constraint (EC) metrics are computed, along with a set of Plant Function Type (PFT) stratified QA/QC variables and bit-flags. All L4_C variables are modeled on a daily basis and internally registered to an EASE-Grid version 2 global 1km extent (14616 lines by 34704 samples, for 507,233,664 pixels per layer), and are then posted to a EASE-Grid v2 global 9km grid (3856 samples by 1624 lines, for 6,262,144 pixels per layer). The model derivation of each of these variables is described in detail in the SMAP L4_C Algorithm Theoretical Basis Document (ATBD).

The SMAP L4_C baseline product has a 9 km spatial resolution consistent with the L4_SM product, but retains sub-grid scale heterogeneity information determined from finer scale (1-km resolution) land cover and FPAR inputs, and carbon model processing. The resulting carbon product is similar to the scale of tower CO₂ eddy covariance flux measurements (Baldocchi et al. 2008, Chen et al. 2012). The baseline L4_C domain encompasses all global vegetated land areas. The L4_C product has a targeted mean RMSE (bias-adjusted) accuracy threshold for NEE within 30 g C m⁻² yr⁻¹ or 1.6 g C m⁻² d⁻¹ over northern (≥45°N) boreal and arctic biomes, which is within the estimated ±30-100 g C m⁻² yr⁻¹ accuracy of *in situ* tower measurements (Baldocchi 2008; Richardson 2005; Richardson 2008). The L4_C baseline product has a mean daily temporal sampling to characterize the dynamic NEE response to daily variations in surface meteorology and associated moisture and thermal controls to GPP and respiration, and for greater precision in the computation of cumulative (weekly, monthly and annual) carbon fluxes. Operational implementation of the baseline L4_C algorithms using L4_SM and GMAO surface meteorology inputs enables spatially and temporally continuous daily mapping of NEE and component carbon fluxes for all vegetated land areas independent of data gaps, vegetation biomass and other constraints on SMAP parameter retrievals. The product is intended to provide NEE and component carbon fluxes over all global vegetated land areas consistent with the scale of carbon processes of the major regional plant functional types.

1.1. Identification

This is the NASA SMAP Mission Product Specification Document (PSD) for the Level 4 Carbon (L4_C) dataset produced within the NASA GMAO, affiliated with the Soil Moisture Active Passive (SMAP) project. The L4_C data product provides estimates of global ecosystem productivity. A unique Digital Object Identifier (DOI) has been assigned to L4_C as:

10.5067/22TFAUSNLO9R. This document applies to any standard L4_C data product generated by the SMAP project.

1.2 Scope

This Product Specification Document (PSD) describes the output file format of the L4_C data product. The intent of this document is to document and clarify the L4_C data product structure as needed for external software interfaces. The SMAP Science Data Management and Archive Plan provides a more comprehensive explanation of these data within the complete context of the SMAP instrument, algorithms, and software.

1.3 The SMAP Experiment

The Soil Moisture Active Passive (SMAP) mission will enhance the accuracy and the resolution of space-based measurements of terrestrial soil moisture and freeze-thaw state. SMAP data products will have a noteworthy impact on multiple relevant and current Earth Science endeavors. These include:

- Understanding of the processes that link the terrestrial water, the energy and the carbon cycles,
- Estimations of global water and energy fluxes over the land surfaces,
- Quantification of the net carbon flux in boreal landscapes
- Forecast skill of both weather and climate,
- Predictions and monitoring of natural disasters including floods, landslides and droughts, and
- Predictions of agricultural productivity.

To provide these data, the SMAP mission deploys a satellite observatory in a near polar, sun synchronous orbit. The observatory includes an L-band radiometer that operates at 1.40 GHz and an L-band radar sensor that operates at 1.26 GHz. The instruments share a rotating reflector antenna with a 6 meter aperture that scans over a 1000 km swath.

As the spacecraft flies from north to south on *descending* orbits, the SMAP instruments view Earth locations at approximately 06:00 local time. As the spacecraft flies from south to north, on *ascending* orbits, the SMAP instruments will view Earth locations at approximately 18:00 local time. The spacecraft operates in a cycle of 117 repeatable orbits.

Each time that the spacecraft repeats the orbit cycle, the nadir path on the Earth's surface may not vary by more than 20 km. The flight plan enables scientists to collect data over any region of the Earth over seasonal and annual cycles and avoid diurnal variations. The combined flight pattern and viewing design enables the observatory to view almost all of the Earth's land mass once every three days.

The SMAP radiometer records microwave emissions from the top 5 cm in the soil with a spatial resolution of about 40 km. Scientific applications based on radiometer measurements in the same frequency range have established this approach as an accurate means to detect the presence or water in near surface soil. The SMAP radar is intended to provide backscatter measurements at 3 km resolution. The combined instrumentation enables SMAP to generate accurate global soil moisture retrievals at 9 km resolution.




Figure 1: Artist's concept of the SMAP observatory.

Geophysical retrievals based on SMAP radar indicate the presence of standing water, the freeze/thaw condition of the surface as well as measures of surface roughness and vegetation. The additional information enables SMAP processors to select appropriate locations for soil moisture retrievals and modeling. In addition, the freeze/thaw data contributes to model enhanced products providing global estimates of the Net Ecosystem Exchange (NEE) of carbon (CO₂) between the Earth's surface and the atmosphere.

1.4 SMAP Data products

During the life-of-mission (LOF) the SMAP Mission produces a series of data products, generally associated with different processing levels 1-4. Lower level products embody data organization schemes closer to the observational platform (e.g swath format, calibrated, geometrically corrected, but un-gridded), while higher level products embody properties more conducive to a broader range of science users, who typically expect and work with data that possess a formal map projection, are gridded, and more importantly, come with more user-oriented tools. The following table shows the relationship of the Level 4 Carbon product among the other SMAP mission data products.

Table 1: SMAP Data Products

SMAP Data Products		
Mission Product Short Name	ECS Short Names	Description
L1A_Radar	SPL1AA	Parsed radar instrument telemetry
L1A_Radiometer	SPL1AP	Parsed radiometer instrument telemetry
L1B_S0_LoRes	SPL1BS0	Low resolution radar (x) in time order
L1C_S0_HiRes	SPL1CS0	High resolution radar (x) on swath grid
L1B_TB	SPL1BTB	Radiometer T (t) in time order
L1C_TB	SPL1CTB	Radiometer T(t)
L2_SM_A	SPL2SMA	Radar soil moisture, includes freeze thaw state
L2_SM_P	SPL2SMP	Radiometer soil moisture
L2_SM_AP	SPL2SMAP	Active Passive soil moisture
L3_FT_A	SPL3FTA	Daily global composite freeze thaw state
L3_SM_A	SPL3SMA	Daily global composite radar soil moisture
L3_SM_P	SPL3SMP	Daily global composite radiometer soil moisture
L3_SM_AP	SPL3SMAP	Daily global composite active passive soil moisture
L4_SM	SPL4TSM	Surface and root zone soil moisture
L4_C	SPL4C	Carbon net ecosystem exchange

1.5 Content Overview

The SMAP L4_C data product contains daily estimates of global ecosystem carbon stocks and fluxes (NEE, Gross Primary Productivity (GPP), heterotrophic respiration (R_h), and surface soil organic carbon (SOC) along with underlying environmental constraint (EC) metrics, and data quality assurance and quality control (QA/QC) attributes. The net ecosystem exchange (NEE) of CO_2 with the atmosphere is a fundamental measure of the balance between carbon uptake by vegetation GPP and carbon losses through autotrophic (R_a) and heterotrophic (R_h) respiration. The sum of R_a and R_h defines the total ecosystem respiration rate (R_{tot}), which encompasses most of the annual terrestrial CO_2 efflux to the atmosphere. NEE, GPP, R_h , and SOC are expressed in units of $\text{g C m}^{-2} \text{yr}^{-1}$. The L4_C data product appears on an Earth fixed, global, cylindrical Equal Area Scalable Earth grid, version 2.0 (EASE-Grid 2.0) at 9km resolution.

As a spatio-temporal data product, the daily global L4_C outputs are organized on the basis of (8) functional land-cover classes called Plant Function Types (PFTs), as defined within the Terra/Aqua MODIS land instrument team MOD/MYD12Q1 (Type-5) global land cover classification. The PFT 2D variable is stored as the Type-5, Plant Function Type scientific data set (SDS) in both the MOD12Q1 (1km resolution), and MYD12Q1 file (500meter resolution) forms. Global 1km PFT classification maps in the original Sinusoidal 1km map projection are thus distributed as part of the Terra/Aqua MODIS MOD12Q1 and related global land cover products. PFT classes directly used as driving parameters in the L4_C algorithm each possess an entry in the L4_C Biome Properties Lookup Table (BPLUT). The table below summarizes the Plant Function Type classifiers, and highlight the (8) the PFT classes used by the L4_C algorithms to partition L4_C output variables.

Table 2: Plant Function Type (PFT) Classifier Summary

Plant Function Type (PFT) Classifier Summary			
PFT Class label	PFT code	PFT Description	PFT Class used in L4_C
Water	0	For all ocean and perennial inland water bodies	No
Evergreen needleleaf	1	Evergreen needle-leaf forest (mostly conifers)	Yes
Evergreen broadleaf	2	Evergreen broadleaf forest	Yes
Deciduous needleleaf	3	Deciduous needle-leaf forest	Yes
Deciduous broadleaf	4	Deciduous broad-leaf forest	Yes
Shrubland	5	Shrub (woody perennial)	Yes
Grassland	6	Grasses (native Graminoids)	Yes
Cereal crops	7	Cereal C3 cropland (domesticated agricultural crops such as wheat, oats, barley, rye)	Yes
Broadleaf crops	8	Broadleaf C4 crop (domesticated agricultural)	Yes
Urban and Built-up	9	Urban and built-up (cities, towns, highways, etc)	No
Snow and ice	10	Snow and ice (may or may not be perennial)	No
Barren (rock) or sparsely vegetated	11	Barren, rock, or very sparsely vegetated land	No
Unclassified	254	Areas otherwise not classified as per above	No

1.6 Related SMAP Project Documents

SMAP L4_C Algorithm Theoretical Basis Document (ATBD, v.1), J. Kimball et al, JPL D-66484, DDR No. SMAP-11-012

SMAP L4_SM Algorithm Theoretical Basis Document (Initial Release v.1), R. Reichle et al., JPL D-66483, October 1, 2012.

SMAP Science Data Management and Archive Plan, JPL D-45973, August 29, 2011.

SMAP Pointing, Positioning, Phasing and Coordinate Systems, Volume 0: Definitions and Principle Coordinate Systems, JPL D-46018, Initial Release, May 18, 2010

SMAP Science Data Calibration and Validation Plan. SMAP Project, JPL D-52544, Jet Propulsion Laboratory, Pasadena, CA

SMAP Applications Plan. SMAP Project, JPL D-53082, Jet Propulsion Laboratory, Pasadena, CA

1.7 Applicable Documents

ISO 19115:2003(E) International Standard – Geographic Information – Metadata, May 1, 2003.

ISO 19115-2:2009 International Standard – Geographic Information – Part 2:Extensions for imagery and gridded data, December 12, 2009.

ISO 19139:2007 International Standard – Geographic Information – Metadata – XML schema implementation, May 14 2009.

Introduction to HDF5, The HDF Group, <http://www.hdfgroup.org/HDF5/doc/H5.intro.html>.

HDF5: API Specification Reference Manual, The HDF Group
http://www.hdfgroup.org/HDF5/doc/RM/RM_H5Front.html

HDF5 User's Guide Release 1.8.9, The HDF Group, <http://hdfgroup.com/HDF5/doc/UG>, May 2012.

EASE-Grid 2.0: Incremental but Significant Improvements for Earth-Gridded Data Sets, Brodzik, M.J., et. al., National Snow and Ice Data Center, Cooperative Institute of Environmental Sciences, University of Colorado, ISPRS International Journal of Geo-Information, ISSN 2220-9964, DOI: 10.3390/igji1010032.

NetCDF Climate and Forecast (CF) Metadata Conventions, Version 1.6, December 5, 2011.

2 Data Product Organization

2.1 File Format

All SMAP standard products are in the Hierarchical Data Format version 5 (HDF5). HDF5 is a scientific multi-platform binary file format, with access by software applications provided by an applications programming interface (API) library. A key distinguishing characteristic of the HDF5 file format is the extensive multi-dimensional, self-referential data model it is built around; these properties allow for very flexible representation in both data and metadata, while providing good I/O performance, storage efficiency, and consistent access across many different compute platforms. The National Center for Supercomputing Applications (NCSA) at the University of Illinois developed HDF to help scientists share data regardless of the source. Modern high performance computing platforms often differ in native word-size and byte ordering. These intrinsic properties govern how the machines internally represent integer and floating point numbers. Use of the HDF library gives users a consistent data model and tool to read HDF files on multiple platforms regardless of the differences in native architecture (endian byte-ordering) and word-size of the platform. Further, HDF files are equally accessible by routines written either in Fortran, C or C++.

The HDF Group is responsible for the development and maintenance of the HDF family of file formats. Users should reference The HDF Group website at <http://www.hdfgroup.org> to download HDF software and documentation.

2.2 HDF File Notation

HDF5 represents a significant departure from the conventions of previous versions of HDF, specifically, HDF v4.x. The changes that appear in HDF5 provide flexibility to overcome many of the limitations of previous releases, particularly regarding total file size and the ability to more naturally represent hierarchical groups of data and attributes. The key concepts of the HDF5 Abstract Data Model are Files, Groups, Datasets, Datatypes, Attributes and Property Lists. The most relevant of these for L4_C data product users are the group, datatype, dataset, and attributes entities. Briefly, HDF **groups** are named organizational levels, arranged in a hierarchy starting at the root or file level, within which data and attributes are encapsulated. Actual data in an HDF5 file are stored in **Datasets**, a term that refers to the primary, multi-dimensional science data stored in the file. Datasets are defined as storing elements belonging to a supported HDF model **datatype**. HDF datatypes (“uint16”, “float32”, etc) are expressively named to include a numeric variable’s ‘sign’ property as signed or unsigned, its type as integer vs. floating point, along with a word-size identifier that helps convey the variables natural numeric representational limits. Attributes refer to a special class of named HDF variable that are most often used to represent metadata, or “data-about-data”. The reader is referred to the HDF user documentation cited in Section 1.7 for formal descriptions of these foundation data model concepts.

2.3 SMAP File Organization

2.3.1 Structure

SMAP data products follow a common convention for all HDF5 Files with regard to both structure and naming. Use of this convention provides uniformity of data access and interpretation. The SMAP Project uses HDF5 Groups, which provide an additional level of data organization. All metadata that pertain to the complete data granule are members of the “/Metadata” Group and are typically stored using HDF Attributes as summarized above. All other data are organized within Groups that are designed specifically to handle the structure and content of each particular data product as specified later in this document.

2.3.2 Data

All data in HDF5 files are stored in individual Datasets. All of the Datasets in an SMAP product are assigned to an HDF5 Group, where the file’s root level (denoted by the group called “/”) also represents an implicit “base” group. A standard field name is associated with each Dataset. The field name is a unique string identifier. The field name corresponds to the name of the data element the Dataset stores. This document lists these names with the description of each data element that they identify.

Each Dataset is associated with an HDF5 Dataspace and an HDF5 Datatype. They provide a minimally sufficient set of parameters for reading the data using standard HDF5 tools.

2.3.3 Element Types

SMAP HDF5 employs the Data Attribute “Type” to classify every data field as a specific data type. The “Type” is an embellishment upon the standard HDF5 Datatypes that is designed specifically to configure SMAP data products.

Table 3 lists all of the “Type” strings that appear in the SMAP data products. The table maps each SMAP “Type” to a specific HDF5 Datatype in both the HDF5 file and in the data buffer. The table also specifies the common conceptual data type that corresponds to the “Type” in SMAP executable code.

Table 3: Element type definitions

Type	HDF5 Datatype (File)	HDF5 Datatype (Buffer)	Conceptual Type
Unsigned8	H5T_STD_U8LE	H5T_NATIVE_UCHAR	unsigned integer
Unsigned16	H5T_STD_U16LE	H5T_NATIVE_USHORT	unsigned integer
Unsigned24	H5T_STD_U16LE, with precision set to 24 bits, and size set to 3 bytes.	H5T_NATIVE_INT	unsigned integer
Unsigned32	H5T_STD_U32LE	H5T_NATIVE_UINT	unsigned integer
Unsigned64	H5T_STD_U64LE	H5T_NATIVE_ULLONG	unsigned integer
Signed8	H5T_STD_I8LE	H5T_NATIVE_SCHAR	signed integer
Signed16	H5T_STD_I16LE	H5T_NATIVE_SHORT	signed integer
Signed32	H5T_STD_I32LE	H5T_NATIVE_INT	signed integer
Signed64	H5T_STD_I64LE	H5T_NATIVE_LLONG	signed integer
Float32	H5T_IEEE_F32LE	H5T_NATIVE_FLOAT	floating point
Float64	H5T_IEEE_F64LE	H5T_NATIVE_DOUBLE	floating point
FixLenStr	H5T_C_S1	H5T_NATIVE_CHAR	character string
VarLenStr	H5T_C_S1, where the length is set to H5T_VARIABLE	H5T_NATIVE_CHAR	character string

SMAP HDF5 files employ two different types of string representation. “VarLenStr” are strings of variable length. “VarLenStr” provides greater flexibility to represent character strings. In an effort to make SMAP HDF5 more interoperable (friendly) for users who wish to use netCDF software, SMAP products restrict the use of “VarLenStr”. “FixLenStr” are strings with a prescribed fixed-length. “FixLenStr” are useful for fixed length strings that are stored in large

multi-dimension arrays. UTC time stamps are an excellent example of the type of data that store well in a "FixLenStr". The L4_C product uses fixed length strings unless otherwise noted.

2.3.4 File Level Metadata

Metadata describing the full content of each granule of the SMAP data product are stored within the explicitly named "/METADATA" Group. SMAP metadata are handled using exactly the same procedures as those that are used to handle SMAP data. The contents of each Attribute that stores metadata conform to one of the SMAP Types. Like data, each metadata element is also assigned a shape. Most metadata elements are stored as scalars. A few metadata elements are stored as arrays.

SMAP data products represent file level metadata in two forms. One form appears in one or more Attributes within the Metadata Group. Combined, those Attributes contain a complete representation of the product metadata. The content conforms to the ISO 19115-2 models in ISO 19139 compliant XML.

The second form of the metadata appears in a set of HDF5 Groups under the "/METADATA" Group. Each of these HDF5 Groups represents one of the major classes in the ISO 19115-2 model. These HDF5 Groups contain a set of HDF5 Attributes. Each HDF5 Attributes represents a specific ISO attribute of the associated ISO class. Although this hierarchical representation inherits its design from the ISO model, it does not completely conform to the model. In many cases, the names of the HDF5 Attributes match those used in the ISO model. In some situations, names were changed to provide greater clarity to SMAP users who are not familiar with the ISO model. Furthermore, to ease metadata searches, the structure of Groups within Groups was limited to four levels.

2.3.5 Local Metadata

SMAP standards incorporate additional metadata that describe each HDF5 Dataset within the HDF5 file. Each of these metadata elements appear in an HDF5 Attribute that is directly associated with the HDF5 Dataset. Where feasible, these HDF5 Attributes employ names that conform to the Climate and Forecast (CF) conventions.

2.4 Data Definition Standards

Section 4.6 of this document specifies the characteristics and definitions of every data element stored in this SMAP data product. Table 5 below defines each of the specific characteristics that are listed in that section of this document. Some of these characteristics correspond with the SMAP HDF5 Attributes that are associated with each Dataset. Data element characteristics that correspond to SMAP HDF5 Attributes bear the same name. The remaining characteristics are descriptive data that help users better understand the data product content.

In some situations, a standard characteristic may not apply to a data element. In those cases, the field contains the character string 'n/a'. Hexadecimal representation sometimes indicates data content more clearly. Numbers represented in hexadecimal begin with the character string '0x'.

Table 4: Definitions of Data element characteristics

Characteristic	Definition
Type	The data representation of the element within the storage medium. The storage class specification must conform to a valid SMAP type. The first column in Table 3 lists all of the valid values that correspond to this characteristic.
Shape	The name of the shape data element that specifies the rank and dimension of a particular data set. Appendix C lists all of the valid shapes that appear in this data product.
Valid_min	The expected minimum value for a data element. In most instances, data element values never fall below this limit. However, some data elements, particularly when they do not reflect normal geophysical conditions, may contain values that fall below this limit.
Valid_max	The expected maximum value for a data element. In most instances, data element values never exceed this limit. However, some data elements, particularly when they do not reflect normal geophysical conditions, may contain values that exceed this limit.
Valid Values	Some data elements may store a restricted set of values (e.g. categorical variables and others). In those instances, this listing specifies the finite list of values that the data element may store.
String Length	This characteristic specifies the length of the data string that represents a single instance of the data element. This characteristic appears exclusively for data elements of FixLenStr type.

Characteristic	Definition
Units	Units of measure. Typical values include “deg”, “degC”, “Kelvins”, “m/s”, “m”, “m**2”, “s” and “counts”. Appendix A and Appendix E include references to important data measurement unit symbols.

2.4.1 Double Precision Time Variables

SMAP double precision time variables contain measurements relative to the J2000 epoch. Thus, these variables represent a real number of Standard International (SI) compatible seconds since 11:58:55.816 on January 1, 2000 UTC.

2.4.2 Array representation

This document employs array notation to demonstrate and clarify the correspondence among data elements in different product data elements. The L4_C is implemented in C/C++ language (c99 standard where supported), so the standard (base0) C/C++ array notation is adopted in this document. All array indices are therefore zero based, and in multidimensional arrays the rightmost subscript index varies most rapidly. This scheme is typically referred to as a “row-major-order” (RMO) indexing scheme, where a linear sequencing through an orthogonal 2D raster extent has the column indices varying fastest and the row indices varying least quickly.

HDF5 is designed to read data seamlessly regardless of the computer language used to write an application. Thus, elements that are contiguous using the dimension notation in this document will appear in contiguous locations in arrays for reading applications in any language with an HDF5 interface.

3 Interface Characteristics

3.1 Coordinate Systems

The SMAP mission uses the Science Orbit Reference Frame (SRF) and the Earth Centered Rotating (ECR) coordinate systems to represent spacecraft attitude, position and relative motion.

The Science Orbit Reference Frame (SRF) is a right-handed coordinate system with its three axes mutually orthogonal. The SRF is defined such that the origin is at the spacecraft center of mass (CM). The +Z axis points toward Geodetic Nadir. Due to the oblateness of the Earth, the vector from the spacecraft to the geometric center of the Earth (Geocentric Nadir) is different from the vector from the spacecraft to the local WGS84 ellipsoid normal (Geodetic Nadir). The +X axis is coplanar with both the +Z axis and the spacecraft inertial velocity vector. The +X axis closely adheres to the direction of the spacecraft inertial velocity vector. The +Y axis completes the right-handed, orthogonal coordinate system. The +Y axis is normal to the orbit plane with positive sense in the direction opposite the orbit angular momentum vector.




Figure 2: The Science Orbit Reference Frame coordinate system.

The Earth Centered Rotating (ECR) or Earth Centered Fixed coordinate system is a right-handed coordinate system with three mutually orthogonal axes. The origin of the system is the Earth's center of mass. The positive x-axis extends from the origin through the intersection of the Equator at 0° latitude and the Greenwich Meridian at 0° longitude. The positive z-axis extends directly North from the origin of the ECR system. Due to a slight wobbling of the Earth, the z-axis does not coincide exactly with the instantaneous rotation axis of the Earth. The y-axis

completes the right- handed coordinate system as a vector from the origin to the intersection of the Equator and 90° East longitude.




Figure 3: Earth Centered Rotating coordinate system.

The SMAP mission adopted the World Reference System WGS84 ellipsoid to define the horizontal Earth reference coordinates. The WGS84 geoid was adopted as the vertical Earth reference coordinates. Geodetic measure is used to define both the spacecraft location and the instrument target location relative to the Earth's surface.

3.2 The Global cylindrical EASE-Grid 2.0

The SMAP L4_C data product is provided in a global cylindrical EASE-Grid 2.0 projection format with a nominal grid spacing of 9 km. Each grid cell has an area of about 81 km², regardless of longitude and latitude; thus grid cells become more elongated in the north-south dimension at higher latitudes relative to grid cells near the equator. Using this projection, all global data arrays have dimensions of 1624 rows and 3856 columns (6,262,144 pixels per layer). Note that the EASE-Grid 2.0 global 1k reference grid is defined as 14616 lines by 34704 samples (507,233,664 pixels per layer). Figure 4 illustrates the region covered by the global cylindrical EASE-Grid 2.0 projection.



Figure 4: Global EASE-Grid 2.0.

The EASE-Grid 2.0 has a flexible formulation. By adjusting a single scaling parameter, a family of multi-resolution grids that “nest” within one another can be generated. The nesting can be made “perfect” so that smaller grid cells can be tessellated to form larger grid cells. Figure 5 shows a schematic of the nesting.




Figure 5: Perfect nesting in EASE-Grid 2.0. Smaller grid cells tessellate into larger grid cells.

The perfect nesting provides SMAP data products with a convenient common projection for both high-resolution radar observations and low-resolution radiometer observations, as well as for their derived geophysical products. Other SMAP data products use 3 km, 9 km and 36 km grid spacing. Figure 6 provides an example of sample Normalized Difference Vegetation Index (NDVI) data on the EASE-Grid 2.0 at these three resolutions.




Figure 6: Sample NDVI data displayed on the SMAP EASE-Grid 2.0 with (left) 36 km, (middle) 9 km, and (right) 3 km nominal grid spacing.

3.3 Fill and Gap Values

SMAP data products employ fill and gap values to indicate when no valid data appear in a particular data element. Fill values ensure that data elements retain the correct shape. Gap values locate portions of a data stream that do not appear in the output data file.

Fill values appear in the L4_C data product over ocean and water surfaces. SMAP data products employ a specific set of data values to connote that an element is fill. The selected values that represent fill are dependent on the data type. Table 5 lists the values that represent fill in SMAP products based on data type:

No *valid* value in the L4_C data product is equal to the values that represent fill. If any exceptions should exist in the future, the L4_C content will provide a means for users to discern between elements that contain fill and elements that contain genuine data values. Operationally, we note that software should not attempt to ever test two floating point values for equality, but instead test two given values using a relational inequality operator and a tolerance based difference in absolute values (e.g. $\text{fabs}(a-b) \leq 1E-14$) method. This document will also contain a description of the method used to ascertain which elements are fill and which elements are genuine.

For discrete categorical variables such as the bit-flags (`carbon_bitflag` and `carbon_qual_flag_{pft}`), all values are defined, therefore no missing data values are expected with these variables. For the analytical variables, the FPAR Climatology provides a fallback input source to help assure there are no spatio-temporal gaps in the modeled data record. Therefore no temporal gaps are expected to occur in the L4_C output data stream. To accommodate potential gaps in the radar Freeze Thaw (F/T) 3KM input variable from L3_SM_A, the L4_C algorithm is designed to fallback and use the F/T derived from surface temperature (`Tsurf`, a modeled LDAS parameter) also carried in the L3_SM_A 3km product. For cases where the entire L3_SM_A daily global granule may be missing, the L4_C algorithm falls back to derive

F/T from a temperature threshold test using the “TS” surface temperature variable from the NASA GEOS5 FP (forward processing) “gph” data product for the day. In all cases, the final F/T state is represented on a EASEGRID v2 3Km grid supplied to the L4_C algorithm.

Table 5: SMAP data product fill values.

Type	Value	Pattern
L4_C : Float32	-9999.0	Negative number outside of valid data range
Float32, Float64	+1e15	Large, negative number
Signed8, NormSigned8	-127	Datatype minimum + 1
Signed16, NormSigned16	-32767	Datatype minimum + 1
Signed24	-8388607	Datatype minimum + 1
Signed32	-2147483647	Datatype minimum + 1
Signed64	-9223372036854775807	Datatype minimum + 1
Unsigned8	254	Datatype maximum - 1
Unsigned16	65534	Datatype maximum - 1
Unsigned24	16777214	Datatype maximum - 1
Unsigned32	4294967294	Datatype maximum - 1
Unsigned64	18446744073709551614	Datatype maximum - 1
FixedLenString, VarLenString	NA	Not available

4 Data Definition

4.1 Product Overview

4.1.1 SMAP L4_C Data Product

The daily global 9km resolution ecosystem productivity variables comprising the L4_C data product include Net Ecosystem Exchange (NEE), gross primary productivity (GPP), heterotrophic respiration (R_h) and surface soil organic carbon (SOC), along with (3) environmental constraint variables (Emult, Tmult, Wmult), each of which are stored as IEEE single precision (4-byte) real values (e.g. "FLOAT32"|"H5T_IEEE_F32LE" HDFv4/HDFv5 data type. Each of these daily model variables, output at 9km resolution on the EASE-Grid 2.0 grid projection, are represented by a "mean" and "standard deviation" dataset named after the variable. Further, the native mapped value of each variable is spatially partitioned by the Plant Function Type (PFT) associated with a given pixel, for PFT classes {1..8}, as summarized in Table 2. This organization is reflected in the HDF5 file using groups named after the ecosystem productivity variable itself.

To avoid repeating key properties (e.g. the datatype, units, and rank/dimensions) for the L4_C output variables in the **L4_C Group Contents** tables appearing later, the **L4_C Data Product Overview** table below summarizes these properties for each variable. The placeholder qualifier notation "**{pft}**" shown in Table 6 (below) is used to indicate that a given ecosystem productivity variable is partitioned into the Plant Function Type (PFT) classifier, for the relevant {1..8} terrestrial PFT's:

Table 6: L4_C Data Product Overview

L4_C Data Product Overview				
Group	Dataset	Datatype	Units	Rank and dimensions
/NEE	nee_{pft:1..8}	Float32	g C m ⁻² day ⁻¹	2: (3856 columns x 1624 rows)
/GPP	gpp_{pft:1..8}	Float32	g C m ⁻² day ⁻¹	2: (3856 columns x 1624 rows)
/Rh	rh_{pft:1..8}	Float32	g C m ⁻² day ⁻¹	2: (3856 columns x 1624 rows)
/SOC	soc_{pft:1..8}	Float32	g C m ⁻²	2: (3856 columns x 1624 rows)
/EC	frozen_area_{pft:1..8}	Float32	percent	2: (3856 columns x 1624 rows)
/EC	emult_av_{pft:1..8}	Float32	percent	2: (3856 columns x 1624 rows)
/EC	tmult_av_{pft:1..8}	Float32	percent	2: (3856 columns x 1624 rows)
/EC	wmult_av_{pft:1..8}	Float32	percent	2: (3856 columns x 1624 rows)
/QA	surface_flag	Uint16	dimensionless	2: (3856 columns x 1624 rows)
/QA	carbon_model_qual_{pft:1..8}	Uint16	dimensionless	2: (3856 columns x 1624 rows)
/QA	carbon_model_bitflag	Uint16	dimensionless	2: (3856 columns x 1624 rows)
/GEO	cell_lat	Float32	decimal deg	2: (3856 columns x 1624 rows)
/GEO	cell_lon	Float32	decimal deg.	2: (3856 columns x 1624 rows)

4.1.2 HDF5 Group GEO Overview

The L4_C product shares certain properties with other SMAP data products, notably its inclusion of the 'latitude' and 'longitude' geolocation fields which play the same role as coordinate variables as defined within the Climate Forecasting (CF-1.x) conventions. Supporting these conventions facilitates additional software tool interoperability, allowing users of L4_C data to use common software tools such as Panoply and Hdfview to visualize and interactively analyze their data. The L4_C geolocation fields are stored in the L4_C granule HDF group named "GEO", and are described in greater detail later in this document.

4.1.3 HDF5 Group QA Overview

The QA group, written to a separate physical file relative to the primary L4_C product granule, stores general product quality information in the form of product bitflag variables (surface_flag, carrying information inherited from L4 and below), the (8) L4_C specific carbon_model_qual_{pft} variable(s), and counts of 1-km pixels simulated within each 9-km grid cell partitioned by Plant Function Class (PFT) as defined from the input 1-km resolution global land cover map and 1-km resolution L4_C processing. The carbon_model_qual_{pft} variable is a 8-bit unsigned integer field binned into discrete categories {0,1,2,3} of lower (0) to larger (3) estimated daily NEE RMSE levels.

4.1.4 L4_C Metadata Overview

The SMAP L4_C metadata are representative of the entire contents of the file. The metadata appear in two forms, that each provide a complete representation of the product metadata.

One form of the metadata appears as HDF5 Attributes that conform to the ISO 19115-2 model in ISO 19139 compliant XML. Metadata in ISO 19139 conformant XML enables users who are familiar with the ISO metadata standards to extract the metadata they need using software that operates with the ISO 19115-2 model and its formal representation.

The second form of the metadata appears in a set of HDF5 groups that contain a set of HDF5 Attributes. The arrangement and names of these groups and their Attribute components approximate major contents of the ISO model. This second form of HDF5 groups and Attributes enable users who are not familiar with the ISO standard to find the particular metadata elements they need to better comprehend product content and format.

4.2 Data Groups and Volume Estimates

L4_C daily 9km data variables are stored in the following HDF root **groups** (/NEE, /GPP, /RH, /SOC, /EC, /QA, and /GEO). Note that the geo-location information for this 9km product includes CF style coordinate variables (separate latitude and longitude bin-center values for each 9km pixel) which reside in the /GEO group. All L4_C quality control information is stored in the /QA group of each L4_C “mdl” output granule, so there is no separate QA file as with some other SMAP products. Metadata unless otherwise noted, is stored in the /METADATA group. Table 7 summarizes the L4_C granule level data volumes.

Table 7: L4_C Data Volume Estimates

L4_C Data Volume Estimates		
Collection	Granule	Expected Data Volumes
L4_C “mdl”	Single granule combining L4C output variables and QA	120 Mb, daily volume
L4_C data product annual total (365 day)		42.78 Gb annual volume
L4_C data product (LOM , 3-yr estimated total **)		128.34 Gb, est Life-of-Mission LOM volume

***Data product total is for the 3-year baseline mission, with a single “mdl”/“qa” product granule per day.*

4.3 File Naming Conventions

Distributable SMAP L4_C data product file names are 45 characters in length. The first 5 characters in the name of all mission distributable products are ‘SMAP_’. These characters identify all products generated by the SMAP mission. The following 5 characters are always ‘L4_C_’. These characters identify the L4_C data product. The subsequent 3 characters identify the file Collection ID (fixed as “MDL”). The following 27 characters uniquely identify the data stored in the file. The final 3 characters of each SMAP Product file name are ‘.h5’. These characters specify the format of the data in the file. More specifically, all SMAP L4_C data product file names must conform to the following convention:

SMAP_L4_C[CollectionID]_[Date/TimeStamp]_[ScienceVersionID]_[Product Counter].[extension]

The outline below describes the content of each field in the file naming convention:

Collection ID – L4_C product and QA granules carry a fixed collection ID of “MDL”.

Date/Time Stamp – The date/time stamp of the data elements that appear in the product. Date/time stamps in SMAP file names are always recorded in Universal Coordinated Time (UTC). Date/time stamps conform to the following convention:

YYYYMMDDThhmmss

where:

YYYY is the calendar year. The full calendar year must appear in the file name.

MM designates the month of the year. The month designator always occupies two digits. Months that can be represented with fewer than two digits must employ a leading zero.

DD designates the day of the month. The day designator always occupies two digits. Days of the month that can be represented with fewer than two digits must employ a leading zero.

T delineates the date from the time, and is a required character in all time stamps in product names.

hh designates the hour of the day on a 24 hour clock in UTC. The hour designator always occupies two digits. Hours that can be represented with fewer than two digits must employ a leading zero.

mm designates the minute of the hour in UTC. The minute designator always occupies two digits. Minutes that can be represented with fewer than two digits must employ leading zeroes.

ss designates the truncated second of the minute in UTC. Fractional second specification is not necessary in file names. The second designator always occupies two digits. Seconds that can be represented with fewer than two digits must employ leading zeroes.

{S} - S{0|1} - data stream ID (0: forward processing (FP), 1: reprocessing (RP))

{M} - major release ID (as defined in the PSD)

{VVV} - L4 version number (e.g. 572 would represent triplet 5.7.2)

{PPP} - product counter (as defined in the PSD)

Extension: the standard .h5 file extension is used.

Science Version ID – The Science Version ID {v(L)(I)(III)} string reflects changes to L4_C algorithm updates that impact the science content of the product. The format of the Science Version ID is as follows:

“V” The character “V” always precedes the version identifier

Launch indicator {L}

Distinguishes between pre-launch and post-launch operating conditions and indicates validation status. One digit launch indicators are assigned as follows:

- 0: pre-launch output (simulated data)
- a: output prior to public beta release (“alpha”)
- b: beta-release output
- v: validated-release output

Major ID {M} One digit that indicates the major version number. Major changes in algorithm or processing approach will generate an update to this identifier.

Minor ID {VVV} Three digits that indicate the minor version number. Any change to any component that impacts the science content of the data product will lead to a change in this identifier.

Note that the data product **Science Version ID** (example: **Vb1003**) is the first part of the data product **Composite Release ID**. The Science Version ID is incremented whenever a change in the L4_C algorithm or its time-invariant ancillary inputs impacts the science content of the L4_C product. The Composite Release ID captures all changes in the L4_C production system, including changes in dynamic ancillary inputs and changes that have no impact on the science content. The Composite Release ID is stored in the metadata (section 4.4) and consists of the Science Version ID plus a four-digit appendix.

Product Counter – The Product Counter tracks the number of times that a particular L4_C data file (or granule) was generated under the same Science Version ID and for the same Collection and time period (or instant). The system assigns a Product Counter of 001 to the first instance of a granule. Subsequent instances of the same granule (that is, representing the same Collection, time period, and Science Version ID) are assigned a Product Counter that represents the next consecutive integer. The Product Counter always occupies three digits. Product Counters that do not require three digits contain leading zeroes

Extension – The extension for all SMAP L4_C data products is “h5”. That extension indicates that the product contents are in HDF5 format. The SMAP SDS will generate a QA file with every data granule. The QA file contains statistical information that will enable users to better assess the quality of the associated granule. QA products bear exactly the same name as the products that they represent. The only difference in names is the extension. The extension for all QA products is “.qa”.

Example File Names – Based on the above standard, the following name describes an operational data beta-release product from science version 1.003 of the L4_C data product for the period 00:00 UTC on June 9, 2015. The file represents the first time (001) an L4_C (“mdl”) product was generated for the date and time interval in question. An example of a full formed SMAP L4_C file name follows:

SMAP_L4_C_mdl_20150609T000000_Vb1003_001.h5

Daily QA information for L4_C is also stored (in the /QA group) within the primary “_mdl_” file documented above.

4.4 Metadata Detail

The L4_C data product contains a significant metadata component, designed to address the needs of users in the discovery, retrieval, and analysis phases of their work, as well as the needs of data managers charged with creating and maintaining data provenance information over time. Unless otherwise noted, all metadata elements conform to the NASA Earth Science Data Model.

The metadata emphasis in this document is on granule level (product specific) metadata, recognizing that some of the granule metadata directly contributes to collection level metadata as well. Structurally, two forms of metadata are included: HDF centric ‘attribute’ name-value pair style metadata, and a form of metadata (stored in nested HDF5 groups) used to help facilitate generation of ISO 19115 / 19139 compliant metadata.

ISO metadata injection for L4_C granules is performed at GSFC/GMAO, as part of a shared L4 metadata post-processing phase. Two phases of metadata injection are performed, where the first is to produce the metadata named in this document as HDF5 hierarchical group form (e.g. “group of groups”). In this treatment, within a given HDF5 group, the nested metadata elements themselves are stored as HDF5 attributes.

The second phase of metadata generation conforms with the ISO 19139 (XML) encoding of the ISO 19115 standard, with its associated ISO 19139 XML representation. This metadata post-processing performed by first extracting the HDF5 group metadata from each L4_C HDF5 granule (using h5dump or equivalent tools) to yield each metadata element’s content. The resulting metadata collection is then run through an XSLT post-process transformation using a Saxon parser, reformatting to be conformal to the ISO 19139 XML schema, for subsequent re-injection back into the granule as conformal XML.

4.4.1 L4_C Mandatory Metadata

In addition to the L4_C product specific metadata, a number of metadata elements (currently 43) are defined as mandatory metadata in order for the granule to conform to the Earth Science Data Model (ESDM).

4.4.2 L4_C Product Specific metadata

The following table defines the L4_C product specific metadata.

Table 8: Granule level metadata in the L4_C data product.

ISO Major Class	SMAP HDF5 Metadata SubGroup	Subgroup/Attribute in SMAP HDF5	Valid Values
DQ_DataQuality	DataQuality/L4_C	Scope	nee, gpp, rh, soc, ec
		CompletenessOmission/evaluationMethodType	directInternal
		CompletenessOmission/measureDescription	Percent of land grid (terrestrial domain) cells, excluding inland water and permanent ice, that lack geophysical data relative to the total number of land grid cells that fall within the global domain (excluding inland water and permanent ice). Metadata values are computed by pooling data from all elements listed in scope.
		CompletenessOmission/nameOfMeasure	Percent of Missing Data
		CompletenessOmission/value	<A measure between 0 and 100>
		CompletenessOmission/unitOfMeasure	Percent
		DomainConsistency/evaluationMethodType	directInternal
		DomainConsistency/measureDescription	Percent of L4_C modeled data that fall within a predefined acceptable range of measure.
		DomainConsistency/nameOfMeasure	Percent of L4_C modeled data that are within the acceptable range.
		DomainConsistency/value	<A measure between 0 and 100>
		DomainConsistency/unitOfMeasure	Percent
EX_Extent	Extent	description	Global land excluding inland water and permanent ice.
		westBoundLongitude	-180 degrees
		eastBoundLongitude	180 degrees
		southBoundLatitude	-90 degrees
		northBoundLatitude	90 degrees

		rangeBeginningDateTime	<Time stamp that indicates the initial time element in the product.>
		rangeEndingDateTime	<Time stamp that indicates the final time of data in the product.>
L_Lineage/LE_ProcessStep	ProcessStep	processor	Soil Moisture Active Passive (SMAP) Mission Science Data System (SDS) Operations Facility
		stepDateTime	<A date time stamp that specifies when the processing of the data product took place.>
		processDescription	L4_C models FPAR 8-DAY composite, GEOS5-FP, and SMAP L4_SM and L3_SM_A data within a global carbon model to generate spatially complete and temporally continuous estimates of NEE (net ecosystem exchange), GPP, Rh, SOC and associated ecosystem variables at 9km.
		documentation	<Software description document title, version, and date stamp.>
		identifier	L4_C_SPS
		runTimeParameters	<Specify any run time parameters if they were used.>
		SWVersionID	<A software version identifier that runs from 001 to 999>
		softwareDate	<A date stamp that specifies when software used to generate this product was released.>
		softwareTitle	Level 4 Surface and Root Zone Soil Moisture SPS
		timeVariableEpoch	J2000
		epochJulianDate	2451545.00
		epochUTCDate	2000-01-01T11:58:55.816Z
		ATBDTitle	Soil Moisture Active Passive (SMAP) Level 4 Carbon (L4_C) Algorithm Theoretical Basis Document (ATBD)

		ATBDDate	<Time stamp that specifies the release date of the ATBD>
		ATBDVersion	<Latest ATBD version number.>
		algorithmDescription	The SMAP L4_C global daily algorithm provides spatially and temporally complete net ecosystem exchange (NEE), gross primary productivity (GPP), heterotrophic respiration (Rh), Soil Organic Carbon (SOC), and related variables at 9km for science and applications users.
		algorithmVersionID	<An algorithm version identifier that runs from 001 to 999>
		version	<The SMAP Composite Release ID(s) of the input data product(s).>
		identifier	SPL2SMDL
		DOI	<Digital object identifier(s) of the input data product(s), if available.>
	Source/L4_SM	description	The surface meteorological data from L4_SM algorithm.
		fileName	<Complete file name(s) of the input data product.>
		creationDate	<Date stamp(s) of the input data product(s).>
		version	<Version ID(s) of the input data product(s).>
		identifier	<Short name(s) of the input data product(s).>
		DOI	<Digital object identifier(s) of the input data product(s), if available.>
	Source/MODIS FPAR 8-DAY	Description	MODIS FPAR 8-day (MOD15A2)
		filename	<Complete file name(s) of the input data product.>
		creationDate	<Date stamp(s) of the input data product(s).>
		Version	<Version ID(s) of the input data product(s).>
		Identifier	<Short name(s) of the input data product(s).>
		DOI	<Digital object identifier(s) of the input data product(s), if available.>

	Source/L4_C_MET	Description	L4_C MET Pre-processor output (L4CMET) to daily L4_C model
		Filename	<Complete file name(s) of the input data product.>
		creationDate	<Date stamp(s) of the input data product(s).>
		Version	<Version ID(s) of the input data product(s).>
		Identifier	<Short name(s) of the input data product(s).>
		DOI	<Digital object identifier(s) of the input data product(s), if available.>
	Source/L4_C_MODIS	Description	L4_C MODIS Pre-processor output (L4CMOD) to daily L4_C model
		Filename	<Complete file name(s) of the input data product.>
		creationDate	<Date stamp(s) of the input data product(s).>
		Version	<Version ID(s) of the input data product(s).>
		Identifier	<Short name(s) of the input data product(s).>
		DOI	<Digital object identifier(s) of the input data product(s), if available.>
	Source/L4_C_SOC	Description	L4_C Soil Organic Carbon (SOC) daily intermediate variables updated daily by L4_C model
		Filename	<Complete file name(s) of the input data product.>
		creationDate	<Date stamp(s) of the input data product(s).>
		Version	<Version ID(s) of the input data product(s).>
		Identifier	<Short name(s) of the input data product(s).>
		DOI	<Digital object identifier(s) of the input data product(s), if available.>

	Source/L4_C_NDVI	Description	L4_C VIIRs NDVI backup source for calculating FPAR
		Filename	<Complete file name(s) of the input data product.>
		creationDate	<Date stamp(s) of the input data product(s).>
		Version	<Version ID(s) of the input data product(s).>
		Identifier	<Short name(s) of the input data product(s).>
		DOI	<Digital object identifier(s) of the input data product(s), if available.>
	Source/L4_C FPAR Climatology	Description	L4_C FPAR 365-day climatology as backup FPAR source for daily L4_C model
		Filename	<Complete file name(s) of the input data product.>
		creationDate	<Date stamp(s) of the input data product(s).>
		Version	<Version ID(s) of the input data product(s).>
		Identifier	<Short name(s) of the input data product(s).>
		DOI	<Digital object identifier(s) of the input data product(s), if available.>
	Source/GEOS5_FP	description	The observation-corrected surface meteorological data from the NASA GEOS-5 system. Precipitation corrected using the gauge-based NOAA Climate Prediction Center Unified precipitation product.
		fileName	<Complete file name(s) of the input data product.>
		creationDate	<Date stamp(s) of the input data product(s).>
		version	<Version ID(s) of the input data product(s).>
		identifier	<Short name(s) of the input data product(s).>

		DOI	<Digital object identifier(s) of the input data product(s), if available.>
DS_Dataset/MD_DataIdentification	DatasetIdentification	creationDate	<Date when the L4_C data product file was created>
		VersionID	<SMAP Science Version ID associated with this data product – See section (4.3) of the L4_C Product Specification Document.>
		CompositeReleaseID	<SMAP Composite Release ID (CRID) associated with this data product. The CRID is the Science Version ID with a four-digit appendix. – See section 4.3 of the L4_SM Product Specification Document.>
		ECSVersionID	<Three-character string that identifies the type of ECS data stream and reflects the major science version of the product, for example: “001” = first operational processing stream; “002” = second operational processing stream (supersedes first stream after major algorithm update and includes reprocessing); “003” = third operational processing stream (supersedes second stream...); “199” = candidate operational stream (a.k.a. OASIS or parallel processing stream) >
		UUID	<Universally Unique identifier (36-character string)>
		fileName	<Name of the L4_C output data file.>
		originatorOrganizationName	NTSG group at the University of Montana, Missoula, Montana, USA
		longName	SMAP Level 4 Carbon <Collection ID> Collection
		shortName	<ECS Short Name>
		SMAPShortName	L4_C <Collection ID> Collection

		abstract	The SMAP L4_C data product provides global, daily net ecosystem exchange, GPP and related variables 9 km resolution. The L4_CM data product consists of one Collection of data granules (or files): the model (MDL) data Collection (shortName=SPL4SMDL).
		characterSet	utf8
		credit	The software that generates the L4_C data product and the data system that automates its production were designed and implemented at the NTSG group at the University of Montana, Missoula, Montana, USA.
		language	eng
		purpose	The SMAP L4_C data product provides spatially and temporally complete net ecosystem exchange, GPP, Rh, Soil Organic Carbon and related variables for science and applications users.
		status	"on-going" ("completed" after final reprocessing)
		topicCategoryCode	geoscientificInformation
		QACreationDate	<The generation date of the QA product that accompanies the L4_C data granule.>
		QAFileName	<The name of QA product.>
		QAAbstract	An HDF5 product that contains statistical information on data product results. These statistics enable data producers and users to assess the quality of the data in the data product granule.
DS_Series/MD_DataIdentificatio n	SeriesIdentification	revisionDate	<Date and time of the software release that was used to generate this data product.>
		VersionID	<SMAP Science Version ID associated with this data product – See section 4.3 of the L4_C Product Specification Document.>
		CompositeReleaseID	<SMAP Composite Release ID (CRID) associated with this data product. The CRID is the Science Version ID with a four-digit appendix. – See section 4.3 of the L4_C Product Specification Document.>

		<p>ECSVersionID</p>	<p><Three-character string that identifies the type of ECS data stream and reflects the major science version of the product, for example: "001" = first operational processing stream; "002" = second operational processing stream (supersedes first stream after major algorithm update and includes reprocessing); "003" = third operational processing stream (supersedes second stream...); "199" = candidate operational stream (a.k.a. OASIS or parallel processing stream)</p> <p>></p>
		<p>longName</p>	<p>SMAP Level 4 Carbon <Collection ID> Collection</p>
		<p>shortName</p>	<p>< ECS Short Name></p>
		<p>identifier_product_DOI</p>	<p><digital object identifier></p>
		<p>resourceProviderOrganization Name</p>	<p>National Aeronautics and Space Administration</p>
		<p>abstract</p>	<p>The SMAP L4_C data product provides global, daily net ecosystem exchange, GPP and related variables 9 km resolution. The L4_CM data product consists of one Collection of data granules (or files): the model (MDL) data Collection (shortName=SPL4SMDL).</p>
		<p>characterSet</p>	<p>utf8</p>
		<p>credit</p>	<p>The software that generates the L4_C data product and the data system that automates its production were designed and implemented at the NTSG group at the University of Montana, Maryland, USA.</p>
		<p>language</p>	<p>Eng</p>
		<p>purpose</p>	<p>The SMAP L4_C data product provides spatially and temporally complete net ecosystem exchange, GPP, Rh, Soil Organic Carbon and related variables for science and applications users.</p>
		<p>status</p>	<p>"on-going" ("completed" after final reprocessing)</p>
		<p>topicCategoryCode</p>	<p>geoscientificInformation</p>

		pointOfContact	NTSG group at the University of Montana, Missoula, Montana, and the National Snow and Ice Data Center, Boulder, Colorado, USA.
		PSDPublicationDate	<Date of publication of the Product Specification Document>
		PSDEdition	<Edition identifier for the Product Specification Document>
		PSDTtitle	Soil Moisture Active Passive Mission Level 4 Carbon (L4_C) Product Specification Document
		SMAPShortName	L4_C
		mission	Soil Moisture Active Passive (SMAP)
		maintenanceAndUpdateFrequency	asNeeded
		maintenanceDate	<Specifies a date when the next update to this product might be anticipated.>
		format	HDF5
		formatVersion	<HDF5 version number>
MD_GridSpatialRepresentation	GridSpatialRepresentation	Latitude/dimensionSize	1624
		Latitude/resolution	9 km on average
		Longitude/dimensionSize	3856
		Longitude/resolution	9 km on average

MD_AcquisitionInformation	AcquisitionInformation	platform/antennaRotationRate	<Antenna rotation rate in rpm>
		platformDocument/publicationDate	<The date of publication of the document that describes the SMAP platform, if available to the general public.>
		platformDocument/edition	<The edition of publication of the document that describes the SMAP platform, if available to the general public.>
		platformDocument/title	<The title of the publication of the document that describes the SMAP platform, if available to the general public.>
		platform/description	The SMAP observatory houses an L-band radiometer that operates at 1.4 GHz and an L-band radar that operates at 1.26-1.29 GHz. The instruments share a rotating reflector antenna with a 6 meter aperture that scans over a 1000 km swath. The bus is a 3 axis stabilized spacecraft that provides momentum compensation for the rotating antenna.
		platform/identifier	SMAP
		radarDocument/publicationDate	<The date of publication of the document that describes the SMAP radar instrument, if available to the general public.>
		radarDocument/edition	<The edition of publication of the document that describes the SMAP radar instrument, if available to the general public.>
		radarDocument/title	<The title of the publication of the document that describes the SMAP radar instrument, if available to the general public.>
		radar/description	The SMAP radar instrument employs an L-band conically scanned system and SAR processing techniques to achieve moderate resolution (1 km) backscatter measurements over a very wide 1000 km swath.
		radar/identifier	SMAP SAR
		radar/type	L-band Synthetic Aperture Radar
		radiometerDocument/publicationDate	<The date of publication of the document that describes the SMAP radiometer instrument, if available to the general public.>
		radiometerDocument/edition	<The edition of publication of the document that describes the SMAP radiometer instrument, if available to the general public.>

		radiometerDocument/title	<The title of the publication of the document that describes the SMAP radiometer instrument, if available to the general public.>
		radiometer/description	The SMAP L-band Radiometer records V-pol, H-pol, 3rd and 4th Stokes brightness temperatures at 40 km resolution at 4.3 Megabits per second with accuracies of 1.3 Kelvin or better.
		radiometer/identifier	SMAP RAD
		radiometer/type	L-band Radiometer

4.5 Data Structure

The data structure of a L4_C granule (HDF5) file is described below, first in its own HDF Group table to show which **group** the indicated dataset is stored within in a HDF5 product granule file. Following these, tables describing a more detailed view of compound (bit-flag) variables and variables with more complex internal organization are included.

4.6 Element definitions

A single L4_C daily global 9km resolution HDF5 product granule is produced each day. Internally, the primary L4_C output HDF5 granule contains (5) HDF5 groups, named after the primary variable indicated e.g. {NEE, GPP, RH, SOC, EC, GEO }, with a final group (GEO) storing geolocation information. Geolocation information in group (GEO) consists of {latitude, longitude} coordinate variables, in decimal degree units, that enable convenient geo-referenced viewing and analysis. All L4_C quality assurance information, in group /QA are stored within the primary L4_C daily HDF5 product file.

Within each of the primary L4_C granule’s HDF5 groups, the indicated variable is partitioned by each of the (8) Plant Function Type classes, such that there are (8) scientific datasets per group. For example, in the NEE group, there are (8) “NEE” output variables, one per PFT class.

The output variables are summarized in Table 6. Note that the primary QA variable is also partitioned by PFT class. Both the QA (partitioned by PFT) and Geolocation information is stored in a companion daily 9km HDF5 granule file, whose name follows the conventions as documented in Section 4.3 above.

4.6.1 NEE Dataset Definition

The following table describes the datasets contained in the HDF group “NEE” (net ecosystem CO₂ exchange) in the L4_C product files:

Table 9: NEE Group Contents

L4_C Contents of HDF Group: NEE		
Group	Dataset	Description
NEE	nee_mean	Mean Net Ecosystem Exchange value, not distinguished by PFT; in gC/m ² units, (valid_min= -30.0,valid_max=20.0)
NEE	nee_std_dev	Standard deviation of Net Ecosystem Exchange, not distinguished by PFT.
NEE	nee_pft_1_mean	Mean value of NEE for pixels classified in PFT class 1 (evergreen needleleaf)
NEE	nee_pft_2_mean	Mean value of NEE for pixels classified in PFT class 2 (evergreen broadleaf)
NEE	nee_pft_3_mean	Mean value of NEE for pixels classified in PFT class 3 (deciduous needleleaf)
NEE	nee_pft_4_mean	Mean value of NEE for pixels classified in PFT class 4 (deciduous broadleaf)
NEE	nee_pft_5_mean	Mean value of NEE for pixels classified in PFT class 5 (shrub)
NEE	nee_pft_6_mean	Mean value of NEE for pixels classified in PFT class 6 (grass)
NEE	nee_pft_7_mean	Mean value of NEE for pixels classified in PFT class 7 (cereal crop)
NEE	nee_pft_8_mean	Mean value of NEE for pixels classified in PFT class 8 (broadleaf crop)

4.6.2 GPP Dataset Definition

The following table describes the datasets contained in the HDF group “GPP” in the L4_C product files:

Table 10: GPP Group Contents

L4_C Contents of HDF Group: GPP		
Group	Dataset	Description
GPP	GPP_mean	Mean Gross Primary Productivity per pixel, not distinguished by PFT; in gC/m ² units, (valid_min=0.0, valid_max =30.0)
GPP	GPP_std_dev	Standard deviation of Gross Primary Productivity per pixel, not distinguished by PFT
GPP	GPP_pft_1_mean	Mean value of GPP for pixels classified as PFT class 1
GPP	GPP_pft_2_mean	Mean value of GPP for pixels classified as PFT class 2
GPP	GPP_pft_3_mean	Mean value of GPP for pixels classified as PFT class 3
GPP	GPP_pft_4_mean	Mean value of GPP for pixels classified as PFT class 4
GPP	GPP_pft_5_mean	Mean value of GPP for pixels classified as PFT class 5
GPP	GPP_pft_6_mean	Mean value of GPP for pixels classified as PFT class 6
GPP	GPP_pft_7_mean	Mean value of GPP for pixels classified as PFT class 7
GPP	GPP_pft_8_mean	Mean value of GPP for pixels classified as PFT class 8

4.6.3 Heterotrophic respiration (R_h) Dataset definition

The following table describes the datasets contained in the HDF group “Rh” in the L4_C product files:

Table 11: Rh Group Contents

L4_C Contents of HDF Group: RH		
Group	Dataset	Description
RH	rh_mean	Mean heterotrophic respiration (Rh) value, not distinguished by PFT; in gC/m ² units, (valid_min=0.0,valid_max=20.0)
RH	rh_std_dev	Standard deviation of heterotrophic (Rh) respiration, not distinguished by PFT
RH	rh_pft_1_mean	Mean value of Rh for pixels classified as PFT class 1
RH	rh_pft_2_mean	Mean value of Rh for pixels classified as PFT class 2
RH	rh_pft_3_mean	Mean value of Rh for pixels classified as PFT class 3
RH	rh_pft_4_mean	Mean value of Rh for pixels classified as PFT class 4
RH	rh_pft_5_mean	Mean value of Rh for pixels classified as PFT class 5
RH	rh_pft_6_mean	Mean value of Rh for pixels classified as PFT class 6
RH	rh_pft_7_mean	Mean value of Rh for pixels classified as PFT class 7
RH	rh_pft_8_mean	Mean value of Rh for pixels classified as PFT class 8

4.6.4 Soil Organic Carbon (SOC) Dataset definition

The following table describes the datasets contained in the HDF group “SOC” (for soil organic carbon) in the L4_C product files:

Table 12: SOC Group Contents

L4_C Contents of HDF Group: Soil Organic Carbon (SOC)		
Group	Dataset	Description
SOC	soc_mean	Mean surface soil organic carbon (SOC) value, not distinguished by PFT; in gC/m ² units (valid_min=0.0, valid_max=25,000)
SOC	soc_std_dev	Standard deviation of surface soil organic carbon (SOC), not distinguished by PFT, in gC/m ² units.
SOC	soc_pft_1_mean	Mean value of SOC for pixels classified as PFT class 1
SOC	soc_pft_2_mean	Mean value of SOC for pixels classified in PFT class 2
SOC	soc_pft_3_mean	Mean value of SOC for pixels classified in PFT class 3
SOC	soc_pft_4_mean	Mean value of SOC for pixels classified in PFT class 4
SOC	soc_pft_5_mean	Mean value of SOC for pixels classified in PFT class 5
SOC	soc_pft_6_mean	Mean value of SOC for pixels classified in PFT class 6
SOC	soc_pft_7_mean	Mean value of SOC for pixels classified in PFT class 7
SOC	soc_pft_8_mean	Mean value of SOC for pixels classified in PFT class 8

The table below describes the datasets contained in the environmental constraints “EC” HDF group in the L4_C product files. Note that these variables are not spatially partitioned by Plant Function Type (PFT) as the other sciences above were.

Table 13: EC -- Environmental Constraints Group Contents

L4_C Contents of HDF Group: EC		
Group	Dataset	Description
EC	frozen_area	Mean frozen_area, per 9x9 km pixel, not distinguished by PFT, units=(0..100 Percent frozen), valid_min = 0.0, valid_max = 100.0
EC	emult_av	Average estimated bulk environmental constraint to PAR conversion efficiency and vegetation productivity, not distinguished by PFT; Units = percent constraint, ranging (min=0 as fully constrained, and max = 100.0 with no constraint)
EC	tmult_av	Average estimated bulk environmental constraint on heterotrophic respiration due to low soil temperature, not distinguished by PFT; Units = percent constraint, ranging (min=0 as fully constrained, and max = 100.0 with no constraint)
EC	wmult_av	Average estimated bulk environmental constraint on heterotrophic respiration due to low soil moisture levels, not distinguished by PFT; Units = percent constraint, with (min=0 as fully constrained, and max = 100.0 with no constraint)

4.6.5 GEO: the Geolocation Group

The following Tables describe the HDF5 datasets present in the GEO HDF5 group, which groups together geo-location fields with the L4_C quality related datasets. Note that for interoperability with common science image data viewers such as Panoply (<http://www.giss.nasa.gov/tools/panoply/>) and HDFView (<http://www.hdfgroup.org/hdf-java-html/hdfview/>), the geolocation variables described here are intended to be functionally equivalent to CF convention “coordinate variables”. In this way, they convey the geographic bin-center coordinate position of each L4_C science variable pixel to the viewer.

Table 14: Overview of Geolocation Group: GEO

L4_C Contents of HDF Group: GEO		
Group	Dataset Name	Description
GEO	latitude	Latitude, in decimal degrees, of the center of the Earth based grid cell.
GEO	longitude	Longitude, in decimal degrees, of the center of the Earth based grid cell.

4.6.6 QA : the Quality Assurance Group

The overall quality of the L4_C daily 9km product, by pixel, is described by several types of quality metrics:

- **carbon_model_bitflag.**
- **carbon_model_qual_pft(1..8)**
- **qa_count_pft{1..8}**
- **qa_count**

Note that the **carbon_model_bitflag** provides ancillary quality related information in the form of a set of decision-tree bit-fields, to assist analysts in performing deeper inquiries describing overall L4_C product quality. The (4) types of L4_C quality metrics are further described in the Table below.

Table 15: Overview of Quality Group: QA

		L4_C Contents of HDF Group: QA	
Group	Dataset Name	Datatype	Description
QA	carbon_model_bitflag	uint16	Bit flag field describing overall L4_C product quality and processing conditions, where a separate QC field is defined for each Plant Function Type (PFT) classifier.
QA	carbon_model_qual_{pft:1..8}	uint8	QA field describing level of estimated daily NEE RMSE averaged by PFT{1..8} and binned to a discrete 4-level score (0=low to 3=high).
QA	qa_count_{pft:1..8}	uint8	Count of 1km pixels simulated (from {0...81} possible) for each PFT present in a 9km cell (excluding non-veg PFTs) for which a valid FPAR or FPAR climatology value was available.
QA	qa_count	uint8	Sum of all 1km pixels simulated (from {0...81} possible) across all PFTs present in a 9km cell (excluding non-veg PFTs) for which a valid FPAR or FPAR climatology value was available. Percent cover for each 1km PFT in a 9km cell calculated as: $\text{pft_qa_count}\{1..8\} / \text{total_qa_count} * 100.$

4.6.7 The carbon_model_bit-flag

The **carbon_model_bitflag** (uint16 bit-flag) variable is represented as a single, spatially explicit quality field output on the 9km grid.

Table 16: L4_C Quality Control Variable (carbon_model_bitflag) Definition

L4_C Quality Control Variable (carbon_model_bitflag) Definition			
Bit field	Bit range (positions)	Value	Description
NEE-bit	0-0 (1-bit)	0 1	When '0', all 1km NEE values within each 9km grid cell lie within their valid range {min, max} and the 9km average is considered 'good'; conversely, when '1', at least one NEE lies outside its valid range and 9 km average is considered 'bad' (or suspect).
GPP-bit	1-1 (1-bit)	0 1	When '0', all 1km GPP values within each 9km grid cell lie within their valid range {min, max} and the 9km average is considered 'good'; conversely, when '1', at least one GPP lies outside its valid range and 9 km average is considered 'bad' (or suspect).
Rh-bit	2-2 (1-bit)	0 1	When '0', all 1km Rh values within each 9km grid cell lie within their valid range {min, max} and the 9km average is considered 'good'; conversely, when '1', at least one Rh lies outside its valid range and 9 km average is considered 'bad' (or suspect).
SOC-bit	3-3 (1-bit)	0 1	When '0', all 1km SOC values within each 9km grid cell lie within their valid range {min, max} and the 9km

			average is considered 'good'; conversely, when '1', at least one SOC lies outside its valid range and 9 km average is considered 'bad' (or suspect).
PFT_dominant	4-7 (4- bits)	Integer value, as PFT class code {1..8} inclusive.	The dominant PFT class code, ranging from {1..8} inclusive, in a 9km cell. The PFT class code is stored in a 4-bit sub-field (capable of representing 16 distinct integer values, of which we use values {1..8}). The dominant PFT class at 9km is defined as the most-frequently occurring PFT class across the 81 input (candidate) 1km pixels from which each 9km output cell is calculated.
QA_score	8-11 (4-bits)	Integer value, 0=best,1=Good, 2=Poor, 3=Inferior	This qa_score (0-3) 4-bit metric represents the overall carbon_model_qual of all PFTs within the 9km grid cell.
GPP_method	12-12 (1-bit)	{0,1}	0=derived GPP using FPAR 8-day input, 1=derived GPP via FPAR-climatology. Indicates where native FPAR vs. the static FPAR climatology is used to compute GPP. If >50% of a 9-km cell uses the ancillary FPAR climatology (defined from 1-km FPAR inputs and L4_C processing), the cell is flagged as 1.
NDVI_method	13-13	{0 1}	0=used either FPAR 8-day or FPAR climatology; 1=used NDVI
FT_method	14-14	{0 1}	0=Radar F/T from L3_SM_A was good and used, or: 1=used Tsurf (surface temperature), as FT from L3_SM_A not available, so used Tsurf to determine F/T state. **
IsFill	15-15	Un-used	0=NO fill value; 1= IsFill value. "IsFill" designated areas indicate L4_C

			simulations not performed due to non-vegetated land or being outside of FPAR or L4_SM domains. Note: a value of 1 indicates that all other bitflags also have a fill value of 1. This is caused by fill value conditions, and NOT by an inferior, inadequate, or incorrect simulation state.
--	--	--	--

** Above as per Scott Hensley, review panel member Sept 12, 2013 Algor. Review.

4.6.8 The carbon_model_qual – partitioned by PFT

The PFT oriented QA partitioning scheme used in L4_C defines a separate 8-bit (uint8) “carbon_model_qual_pft(1..8)” layer is produced for each of the (8) primary Plant Functional Types. In this case, the quality score refers to the PFT-wise average quality. Here quality is defined in terms of the magnitude of estimated NEE RMSE from relative low (low RMSE) to high (higher RMSE) category levels. Note that RMSE is generally proportional to the level of estimated productivity (GPP) so that higher productivity areas (e.g. croplands) will have a higher RMSE QA category than less productive (e.g. shrubland) areas.

Table 17: L4_C Quality Control Variable (carbon_model_qual_{pft:1..8}) Definition

Dataset Variable	Value	Description
Carbon_model_qual_pft1	{0,1,2,3}	For PFT 1, binned to a 4-level score (0=low to 3=high).
Carbon_model_qual_pft2	{0,1,2,3}	For PFT 2, binned to a 4-level score (0=low to 3=high).
Carbon_model_qual_pft3	{0,1,2,3}	For PFT 3, binned to a 4-level score (0=low to 3=high).
Carbon_model_qual_pft4	{0,1,2,3}	For PFT 4, binned to a 4-level score (0=low to 3=high).
Carbon_model_qual_pft5	{0,1,2,3}	For PFT 5, binned to a 4-level score (0=low to 3=high).
Carbon_model_qual_pft6	{0,1,2,3}	For PFT 6, binned to a 4-level score (0=low to 3=high).
Carbon_model_qual_pft7	{0,1,2,3}	For PFT 7, binned to a 4-level score (0=low to 3=high).
Carbon_model_qual_pft8	{0,1,2,3}	For PFT 8, binned to a 4-level score (0=low to 3=high).

4.7 L4_C Data Product Volume Estimates

The table below summarizes early estimates of the L4_C pre-product data volume, with a granule data size of 120 Mb/granule.

Table 18: L4_C Data Product Volume Estimates

L4_C Data Product Volume Estimate			
HDF Group	Description	Bytes	Estimated compressed volume, in bytes
Metadata	Collective granule level Metadata, both ESDM and ISO forms	160000 est	TBD
GEO	Geolocation (latitude, longitude) and all QC, bit-flags	75145728	TBD
QA	(8) QA (carbon_qual) layers, 1 per PFT class.		9.5 Mb /day
QA	1 (uint16) surface_flags, 1 bitflags (uint16) layer		
QA	(8) QA pft_freq 9km uint8 layers, 1 per PFT class and (1)pft_mod_cnt uint8 9km layer		
GPP	Gross Primary productivity datasets for all (8) PFTs	250485760	105 Mb/ day
Rh	Rh for (8) PFTs	250485760	
NEE	NEE for (8) PFTs	250485760	
SOC	SOC for (8) PFTs	250485760	
EC	(3) environmental constraint variables (emult, tmult, wmult)	25048576	
TOTALS	Granule, total Gb	0.12 Gb (= 120Mb)	

5 Appendices

Appendix A: Acronyms and abbreviations

This is the standard Soil Moisture Active Passive (SMAP) Science Data System (SDS) list of acronyms and abbreviations. Not all of these acronyms and abbreviations appear in every SMAP SDS document.

ADT	Algorithm Development Team
AMSR	Advanced Microwave Scanning Radiometer
ANSI	American National Standards Institute
AOS	Acquisition of Signal
APF	Algorithm Parameter File
ARS	Agricultural Research Service
ASF	Alaska Satellite Facility
ATBD	Algorithm Theoretical Basis Document
ATLO	Assembly Test Launch and Operations
BFPQ	Block Floating Point Quantization
BIC	Beam Index Crossing
CARA	Criticality and Risk Assessment
CBE	Current Best Estimate
CCB	Configuration Control Board
CCSDS	Consultative Committee on Space Data Systems
CDR	Critical Design Review
CEOS	Committee on Earth Observing Systems
CF	Climate and Forecast (metadata convention)
CM	Configuration Management
CM	Center of Mass
CONUS	Continental United States
COTS	Commercial Off the Shelf
CR	Change Request
DAAC	Distributed Active Archive Center
DB	Database

DBA	Database Administrator
dB	decibels
deg	degrees
deg/sec	degrees per second
deg C	degrees Celsius
DEM	Digital Elevation Model
DFM	Design File Memorandum
DIU	Digital Interface Unit
DN	Data Number
DOORS	Dynamic Object Oriented Requirements
DQC	Data Quality Control
DSK	Digital Skin Kernel
DVD	Digital Versatile Disc
EASE	Equal Area Scalable Earth
ECMWF	European Centre for Medium Range Weather Forecasts
ECHO	EOS Clearing House
ECI	Earth Centered Inertial Coordinate System
ECR	Earth Centered Rotating Coordinate System
ECR	Engineering Change Request
ECS	EOSDIS Core System
EDOS	EOS Data Operations System
EM	Engineering Model
EOS	Earth Observing System
EOSDIS	Earth Observing System Data and Information System
EPO	Education and Public Outreach
ESDIS	Earth Science Data and Information System Project
ESDT	Earth Science Data Type
ESH	EDOS Service Header
ESSP	Earth Science System Pathfinder
ET	Ephemeris Time
EU	Engineering Units

FOV	Field of View
FRB	Functional Requirements Baseline
FS	Flight System
FSW	Flight Software
F/T	Freeze/Thaw
FTP	File Transfer Protocol
GByte	gigabyte
GDS	Ground Data System
GEOS-5	Goddard Earth Observing System, version 5
GHA	Greenwich Hour Angle
GHz	gigahertz
GLOSIM	Global Simulation
GMAO	Global Modeling and Assimilation Office
GMT	Greenwich Mean Time
GN	Ground Network
GPMC	Governing Program Management Council
GPP	Gross Primary Production
GPS	Global Positioning System
GSE	Ground Support Equipment
GSFC	Goddard Space Flight Center
HDF	Hierarchical Data Format
HK	Housekeeping (telemetry)
Hz	Hertz
HSD	Health and Status Data
ICE	Integrated Control Electronics
ICESat	Ice, Cloud and Land Elevation Satellite
IDL	Interactive Data Language
I&T	Integration and Test
ICD	Interface Control Document
IEEE	Institute of Electrical and Electronics Engineers
IFOV	Instantaneous Field of View

I/O	Input/Output
IOC	In-Orbit Checkout
IRU	Inertial Reference Unit
ISO	International Organization for Standardization
IV&V	Independent Verification and Validation
ITAR	International Traffic in Arms Regulations
I&T	Integration and Test
JPL	Jet Propulsion Laboratory
KHz	kilohertz
km	kilometers
LAN	Local Area Network
LBT	Loopback Trap
LDAS	Land Data Assimilation System
LEO	Low Earth Orbit
LEOP	Launch and Early Operations
LOE	Level Of Effort
LOM	Life Of Mission
LOS	Loss of Signal
LSK	Leap Seconds Kernel
LZPF	Level Zero Processing Facility
m	meters
MHz	megahertz
MIT	Massachusetts Institute of Technology
MMR	Monthly Management Review
MOA	Memorandum of Agreement
MOC	Mission Operations Center
MODIS	Moderate Resolution Imaging Spectroradiometer
MOS	Mission Operations System
m/s	meters per second
ms	milliseconds
MS	Mission System

NAIF	Navigation and Ancillary Information Facility
NASA	National Aeronautics and Space Administration
NCEP	National Centers for Environmental Protection
NCP	North Celestial Pole
NCSA	National Center for Supercomputing Applications
NEDT	Noise Equivalent Diode Temperature
NEE	Net Ecosystem Exchange
NEN	Near Earth Network
netCDF	Network Common Data Form
NFS	Network File System/Server
NISN	NASA Integrated Services Network
NRT	Near Real Time
NOAA	National Oceanic and Atmospheric Administration
NSIDC	National Snow and Ice Data Center
NVM	Non-Volatile Memory
NWP	Numerical Weather Product
n/a	not applicable
OCO	Orbiting Carbon Observatory
OEF	Orbit Events File
ORBNUM	Orbit Number File
ODT	Object Oriented Data Technology
ORR	Operational Readiness Review
ORT	Operational Readiness Test
OSSE	Observing System Simulation Experiment
OSTC	One Second Time Command
PALS	Passive and Active L-Band System
PALSAR	Phased Array L-Band Synthetic Aperture Radar
PcK	Planetary Constants Kernel
PDR	Preliminary Design Review
PPPCS	Pointing, Position, Phasing and Coordinate System
PR	Problem Report

PRF	Pulse Repetition Frequency
PRI	Pulse Repetition Interval
PROM	Programmable Read Only Memory
PSD	Product Specification Document
QA	Quality Assurance
rad	radians
RAM	Random Access Memory
RBA	Reflector Boom Assembly
RBD	Rate Buffered Data
RBE	Radiometer Back End
RDD	Release Description Document
RDE	Radiometer Digital Electronics
RF	Radio Frequency
RFA	Request For Action
RFE	Radiometer Front End
RFI	Radio Frequency Interference
RMS	root mean square
RSS	root sum square
ROM	Read Only Memory
RPM	revolutions per minute
RVI	Radar Vegetation Index
SA	System Administrator
SAR	Synthetic Aperture Radar
S/C	Spacecraft
SCE	Spin Control Electronics
SCLK	Spacecraft Clock
SDP	Software Development Plan
SDS	Science Data System
SDT	Science Definition Team
SI	International System
SITP	System Integration and Test Plan

SMAP	Soil Moisture Active Passive
SMEX	Soil Moisture Experiment
SMOS	Soil Moisture and Ocean Salinity Mission
SMP	Software Management Plan
SNR	signal to noise ratio
SOC	Soil Organic Carbon
SOM	Software Operators Manual
SQA	Software Quality Assurance
SPDM	Science Process and Data Management
SPG	Standards Process Group
SPK	Spacecraft Kernel
SQA	Software Quality Assurance
SPS	Science Production Software
SRF	Science Orbit Reference Frame
SRR	System Requirements Review
SRTM	Shuttle Radar Topography Mission
SSM/I	Special Sensor Microwave/Imager
STP	Software Test Plan
sec	seconds
TAI	International Atomic Time
Tb	Brightness Temperature
TBC	To Be Confirmed
TBD	To Be Determined
TBR	To Be Resolved
TBS	To Be Specified
TCP/IP	Transmission Control Protocol/Internet Protocol
TEC	Total Electron Content
TM	Trademark
TOA	Time of Arrival
TPS	Third Party Software
UML	Unified Modeling Language

U-MT	University of Montana
USDA	United States Department of Agriculture
UTC	Coordinated Universal Time
V&V	Verification and Validation
VWC	Vegetation Water Content

Appendix B: L4_C Dimensions

Table 18 lists all of the Dimensions that are used by data elements in the L4_C data product. The name of each Dimension matches the name given in the Dimension column below. The table also lists the anticipated nominal value and the maximum value for each dimension that appears in the L4_C data product.

Table 18: Dimensions in the SMAP L4_SM data product.

Dimension	Nominal Size	Maximum Size
Latitude	1624	1624
Longitude	3856	3856

Appendix C: L4_C Shapes

Table 19: Shapes in the SMAP L4_SM data product.

Shape	Rank	Nominal Product Dimensions	Maximum Product Dimensions
LatCell_LonCell_Array	2	(1624, 3856)	(1624, 3856)

Appendix D: L4_C Units

Table 20 lists the units that are used by the L4_C data product elements. The SMAP implementation of HDF5 stores unit information for each data element in local metadata. The first column in the Table 17 identifies units that apply to data in the L4_C data product. The second column lists the Common

Symbol used to represent the unit. The third column lists the matching Label that appears in the local metadata in the L4_C data product.

Table 20: Units in the SMAP L4_C data product.

Unit	Common Symbol	L4_C Label	Typical Use
counts	Counts	counts	number of elements in a set
degrees	degrees	degrees	angular measure
dimensionless	n/a	dimensionless	dimensionless quantity
Gigahertz	GHz	GHz	frequency measure
megabytes	Mbytes	Mbytes	computer storage units
meters	m	m	distance measure
kilometers	km	km	distance measure
percent	%	percent	per hundred
seconds	s	sec	time measure
revolutions per minute	rpm	rpm	rotational measure
degrees Celsius	°C	Degrees Celsius	temperature measure
Kelvin	K	Kelvin	temperature measure
Pascal	Pa	Pa	pressure measure
kilograms per kilogram	kg kg ⁻¹	kg kg-1	mass per mass measure
meters per second	m s ⁻¹	m s-1	velocity measure
meters squared per meters squared	m ² m ⁻²	m2 m-2	area per area measure
meters cubed per meters cubed	m ³ m ⁻³	m3 m-3	volume per volume measure
kilograms per meters squared	kg m ⁻²	kg m-2	mass storage per area measure
kilograms per meters squared per second	kg m ⁻² s ⁻¹	kg m-2 s-1	mass flux measure
Watts per meters squared	W m ⁻²	W m-2	energy flux measure