

## Comparison of Surface and Column Variations of CO<sub>2</sub> over Urban Areas for Future Active Remote CO<sub>2</sub> sensors

Yonghoon Choi<sup>1</sup>, Melissa Yang<sup>2</sup>, Susan Kooi<sup>1</sup>, and Edward Browell<sup>3</sup>

<sup>1</sup>Science Systems and Applications, Inc., Hampton, VA

<sup>2</sup>NASA Langley Research Center, Hampton, VA

<sup>3</sup>STARSS-II Affiliate, NASA Langley Research Center, Hampton, VA

High resolution in-situ CO<sub>2</sub> measurements were recorded onboard the NASA P-3B during the DISCOVER-AQ (Deriving Information on Surface Conditions from Column and Vertically Resolved Observations Relevant to Air Quality) Field Campaign, to investigate the ability of space-based observations to accurately assess near surface conditions related to air quality. This campaign includes, Washington DC/Baltimore, MD (July 2011), San Joaquin Valley, CA (January – February 2013), Houston, TX (September 2013), and Denver, CO (July-August 2014). Each of these campaigns consisted of missed approaches and approximately two hundred vertical soundings of CO<sub>2</sub> within the lower troposphere (surface to about 5 km). In this study, surface (0 – 1 km) and column-averaged (0 – 3.5 km) CO<sub>2</sub> mixing ratio values from the vertical soundings in the four geographically different urban areas are used to investigate the temporal and spatial variability of CO<sub>2</sub> within the different urban atmospheric emission environments. Tracers such as CO, CH<sub>2</sub>O, NO<sub>x</sub>, and NMHCs are used to identify the source of CO<sub>2</sub> variations in the urban sites. Additionally, we apply nominal CO<sub>2</sub> column weighting functions for potential future active remote CO<sub>2</sub> sensors operating in the 1.57- $\mu$ m and 2.05- $\mu$ m measurement regions to convert the in situ CO<sub>2</sub> vertical mixing ratio profiles to variations in CO<sub>2</sub> column optical depths, which is what the active remote sensors actually measure. Using statistics calculated from the optical depths at each urban site measured during the DISCOVER-AQ field campaign and for each nominal weighting function, we investigate the natural variability of CO<sub>2</sub> columns in the lower troposphere; relate the CO<sub>2</sub> column variability to the urban surface emissions; and show the measurement requirements for the future ASCENDS (Active Sensing of CO<sub>2</sub> Emissions over Nights, Days, and Seasons) in the continental U.S. urban areas.