

C III] EMISSION IN STAR-FORMING GALAXIES NEAR AND FAR

J. R. RIGBY¹, M. B. BAYLISS^{2,3}, M. D. GLADDERS^{4,5}, K. SHARON⁶, E. WUYTS⁷, H. DAHLE⁸, T. JOHNSON⁶, & M. PEÑA-GUERRERO⁹*Accepted for publication in ApJ Letters*

ABSTRACT

We measure C III] $\lambda\lambda 1907, 1909 \text{ \AA}$ emission lines in eleven gravitationally-lensed star-forming galaxies at $z \sim 1.6\text{--}3$, finding much lower equivalent widths than previously reported for fainter lensed galaxies (Stark et al. 2014). While it is not yet clear what causes some galaxies to be strong C III] emitters, C III] emission is not a universal property of distant star-forming galaxies. We also examine C III] emission in 46 star-forming galaxies in the local universe, using archival spectra from GHRS, FOS, and STIS on *HST*, and *IUE*. Twenty percent of these local galaxies show strong C III] emission, with equivalent widths $< -5 \text{ \AA}$. Three nearby galaxies show C III] emission equivalent widths as large as the most extreme emitters yet observed in the distant universe; all three are Wolf-Rayet galaxies. At all redshifts, strong C III] emission may pick out low-metallicity galaxies experiencing intense bursts of star formation. Such local C III] emitters may shed light on the conditions of star formation in certain extreme high-redshift galaxies.

Subject headings: galaxies: star formation — techniques: spectroscopic — gravitational lensing: strong

1. INTRODUCTION

Rest-frame ultraviolet spectroscopy of star-forming $z \sim 2\text{--}7$ galaxies has revealed strong (rest-frame equivalent width $W_r < -5 \text{ \AA}$) emission lines of C III] $\lambda\lambda 1907, 1909 \text{ \AA}$ (Shapley et al. 2003; Erb et al. 2010; Stark et al. 2014, 2015). Stark et al. (2014) suggest that C III] emission may be a way to spectroscopically confirm the highest-redshift galaxies. Such a tool would be particularly important if Lyman α emission from galaxies at the reionization epoch is suppressed by neutral gas absorption (Treu et al. 2012, 2013; Pentericci et al. 2011, 2014; Faisst et al. 2014; Tilvi et al. 2014; Caruana et al. 2013, 2014). Such studies would need to understand selection effects—namely, the frequency of strong emission among star-forming galaxies.

There is a wide range in observed C III] equivalent width at $z \sim 2$ (see Stark et al. 2014 Figure 3). It is not clear what physical conditions cause galaxies to produce strong equivalent width C III] emission; Stark et al. (2014) find that the emission line ratios of strong C III] emitters at $z \sim 2$ are consistent with low gas metallicity, high ionization parameter, and a hard radiation field.

In addition, the C III] lines, together with lines of sim-

ilar ionization potentials, namely O III], N III], N IV], and Si III], can measure atomic abundances (see study at $z = 3.6$ by Bayliss et al. 2014), and thus may be used to constrain nucleosynthetic yields in the early Universe.

This Letter has three objectives. First, we measure C III] equivalent width in 11 gravitationally-lensed galaxies at $z \sim 2\text{--}3$. This increases the published set of such measurements by a third, and explores a different parameter space than Stark et al. (2014), namely, higher stellar mass and metallicity. Second, we measure C III] equivalent width for star-forming galaxies in the local universe. For nearby samples it is easier to measure physical conditions, spatially resolve ionized regions, and test hypotheses for what produces strong C III] emission. Moreover, if strong C III] emission is a signature of primitive galaxies, then nearby galaxies with such emission should be investigated as analogues of high redshift galaxies. Third, we explore the dependence of C III] equivalent width on gas-phase metallicity.

2. DATA AND METHODS

2.1. Measurements for distant galaxies

We measure the equivalent widths of Ly α and the C III] doublet in the rest-frame UV spectra of 11 gravitationally-lensed galaxies at $1.6 < z < 3.1$. Spectra were obtained using the MagE spectrograph (Marshall et al. 2008) on the Clay Magellan II telescope, as part of a large study of the rest-frame ultraviolet spectral properties of bright lensed galaxies (Rigby et al. in prep.)

Intrinsic (corrected for lensing magnification) stellar masses and star formation rates have been published for three of these galaxies (Wuyts et al. 2012a), and are $3\text{--}7 \times 10^9 M_\odot$ and $20\text{--}100 M_\odot \text{ yr}^{-1}$. Since final lensing models are in development for the MagE sample, we cannot yet quote stellar masses and star formation rates for the others. Metallicities from [N II]/H α have been published for four of the sample galaxies: SGAS 090003.3+223408 (Bian et al. 2010), the Cosmic Horseshoe (Hainline et al. 2009),

¹ Astrophysics Science Division, Goddard Space Flight Center, 8800 Greenbelt Rd., Greenbelt, MD 20771

² Department of Physics, Harvard University, 17 Oxford St., Cambridge, MA 02138

³ Harvard-Smithsonian Center for Astrophysics, 60 Garden St., Cambridge, MA 02138

⁴ Department of Astronomy & Astrophysics, University of Chicago, 5640 S. Ellis Ave., Chicago, IL 60637

⁵ Kavli Institute for Cosmological Physics, University of Chicago, 5640 South Ellis Ave., Chicago, IL 60637

⁶ Department of Astronomy, University of Michigan, 500 Church St., Ann Arbor, MI 48109

⁷ Max Planck Institute for Extraterrestrial Physics, Giessenbachstrasse 1, 85748 Garching, Germany

⁸ Institute of Theoretical Astrophysics, University of Oslo, P.O. Box 1029, Blindern, NO-0315 Oslo, Norway

⁹ Space Telescope Science Institute, 3700 San Martin Dr., Baltimore, MD 21218

SGAS J152745.1+065219 (Wuyts et al. 2012a), and RCGA 032727–132609 (hereafter RCS0327, Whitaker et al. 2014; Wuyts et al. 2014). We measure the metallicity for a fifth galaxy, SGAS J000451.7–010321, from the $[\text{N II}]/\text{H}\alpha$ ratio in FIRE/Magellan spectra. We use the second-order polynomial calibration of Pettini & Pagel (2004).

From the literature, we take C III] equivalent widths for six galaxies from Erb et al. (2010), Christensen et al. (2012), Stark et al. (2014), and Bayliss et al. (2014) with measured metallicities. Four galaxies have metallicities derived by Stark et al. (2014) using a Bayesian approach that relies on the strong emission lines used by the R_{23} method; the R_{23} metallicity measured by Christensen et al. (2012) for one of those galaxies agrees with the Bayesian method. Erb et al. (2010) measured an R_{23} metallicity for BX418 at $z = 2.30$, as did Bayliss et al. (2014) for SGAS J105039.6+001730 at $z = 3.625$.

2.2. Measurements for nearby galaxies

We take spectra of nearby galaxies from three spectral atlases: Leitherer et al. (2011), Peña-Guerrero et al. (in prep.), and Giavalisco et al. (1996).

For 17 galaxies from Leitherer et al. (2011), we take metallicities from their tabulation, and measurements of C III] equivalent width from Bayliss et al. (2014). Four of these galaxies have spectra of multiple regions, totaling 27 measurements of C III]. Thirteen of these galaxies have archival *International Ultraviolet Explorer* (*IUE*) spectra that cover C III], many of which show C III] emission, even in favorable cases lines of moderate equivalent width, example IRAS 08339+6517. For the 18 galaxies of Peña-Guerrero et al. (in prep.), we measure C III] equivalent widths from the *HST* spectra, and take direct gas-phase metallicity measurements from their Table 9. For the 11 galaxies (out of 22) from Giavalisco et al. (1996) that have continuum signal-to-noise ratio $\gtrsim 1$ per pixel, we measure equivalent widths from archival low dispersion (6 Å) *IUE* spectra.

We measure Ly α equivalent widths for the subset of those galaxies that have archival spectra covering Ly α , and redshifts sufficient, given the spectral resolution of each spectrum, to cleanly separate in the galaxy’s Ly α emission from the much brighter geocoronal Ly α emission (Biretta et al. 2015). These criteria are met for: Tol-1214-277 from Leitherer et al. (2011) (*IUE* spectra); two galaxies from Peña-Guerrero et al. (in prep.), IRAS 08339+6517 from *HST* and *IUE* spectra, and NGC 1741 from an *IUE* spectrum; and the 11 galaxies from Giavalisco et al. (1996) (*IUE* spectra).

These are spectra from published atlases, not from a complete galaxy sample. Further, they were obtained using a range of aperture size: $10 \times 20''$ with *IUE*, $1\text{--}2''$ with FOS and GHRIS, and $0.2''$ with STIS. As such, some observations capture bright star-forming regions within a galaxy, whereas others capture the integrated galaxy spectrum. These are necessary limitations of current ultraviolet spectroscopic atlases.

2.3. Uncertainties

When fitting equivalent widths of C III] and Ly α , our uncertainties incorporate statistical uncertainty as well as systematic uncertainty due to continuum estimation.

Each reported equivalent width is the mean of measurements assuming a linear continuum, power law continuum, and (if the spectrum is locally flat) median continuum. We take the standard deviation as the systematic uncertainty due to continuum fitting; it is typically less than half the measurement uncertainty.

The spectra analyzed here vary widely in resolution and signal-to-noise ratio, resulting in emission line equivalent width measurements ranging from strong to weak detections. For sources with strong emission lines and weak or non-detected continuum, the measurement is a lower limit on the equivalent width; the associated uncertainty may be large even though the emission line is very well detected, and reflects the continuum uncertainty.

Following the definition of equivalent width, equivalent widths have positive sign for absorption, and negative sign for emission.

3. RESULTS

3.1. C III] equivalent width

We measure C III] strengths in bright lensed, star forming galaxies at $z \sim 2\text{--}4$, as well as in local starbursting galaxies with archival ultraviolet spectra from *HST* and *IUE*. Figure 1 compares histograms of C III] equivalent width in these samples, as well as to Stark et al. (2014) and the composite spectrum of Shapley et al. (2003). Figure 1 demonstrates that star-forming galaxies in the local universe and at $z \sim 2\text{--}4$ show a large range of C III] equivalent width. The mode for the distant galaxies and the mode for the $z \sim 0$ galaxies are consistent with the equivalent width measured for the Shapley et al. (2003) composite. A significant minority of galaxies show strong C III] emission with $W_r(\text{C III])} < -5 \text{ \AA}$: 22% of the $z \sim 0$ sample, and 45% of the $z \sim 2\text{--}4$ sample.

Indeed, the local sample includes a tail of strong C III] emitters with equivalent widths to -25 \AA , widths that are beyond that yet observed at $z \sim 2$. These results are consistent with a scenario in which strong C III] emission is not a particular feature of $z \gtrsim 2$ galaxies, but rather occurs in a minority of starbursting galaxies independent of redshift.

The MagE sample has systematically lower equivalent widths than the Stark et al. (2014) sample. The MagE sample galaxies were selected as lensed galaxies that are most amenable to deep rest-frame UV spectroscopy; they have typical apparent magnitudes of $g_{AB} \sim 20\text{--}21$, and should be biased toward high surface brightness, high magnification, high luminosity, and compact morphology. Wuyts et al. (2012a) examined in detail the spectral energy distributions of three galaxies from the MagE sample, finding them typical of Lyman Break Galaxies, though younger than average. By contrast, the Stark et al. (2014) sample is selected to be much fainter, with an average apparent V-band magnitude of $m_{AB} = 23.9$.

3.2. C III] equivalent width as a function of metallicity

Figure 2 plots the C III] equivalent width as a function of gas-phase metallicity, for the $z \sim 0$ samples (black symbols) and the $z \sim 2\text{--}4$ samples (blue and purple symbols). Stark et al. (2014) suggest that low metallicity may be largely responsible for large equivalent widths. Figure 2 reveals that the relationship with metallicity is complex. Strong emission ($\lesssim -5 \text{ \AA}$) is observed in galax-

ies over a wide range of metallicity, from 6% to 40% of solar. Strong emission appears not to occur at metallicities above $12 + \log(O/H) = 8.4$. Indeed, only one of the six lowest-metallicity $z \sim 0$ galaxies show strong emission. Thus, low metallicity may be a necessary but not sufficient condition for C III] emission.

3.3. Strong C III] emitters in the local universe

Figure 2 demonstrates that the local universe contains galaxies whose C III] emission is as strong, indeed stronger, than yet observed at $z \sim 2$. The three galaxies from the $z \sim 0$ samples with the most extreme C III], with equivalent widths exceeding -14\AA , are Tololo 1214 – 277, Markarian 71 (a star-forming region within NGC 2366),¹⁰ and Pox 120. We now explore what is known about these three galaxies, and plot their ultraviolet spectra in Figure 3.

Tololo 1214 – 277 is classified as a Wolf-Rayet galaxy; Mrk 71 is classified as a star cluster containing Wolf-Rayet stars within the irregular galaxy NGC 2366, and Pox 120 is classified as a Wolf-Rayet galaxy (Kunth & Joubert 1985; Schaerer et al. 1999). The equivalent width of $H\alpha$ is extremely high for Mrk 71 ($< -1000\text{\AA}$, Buckalew et al. 2005), and for TOL 1214-277 (-1700\AA , Guseva et al. 2011). We could not find a published $W_r(H\alpha)$ for Pox 120. Only 1 (15) of the 417 nearby galaxies of Moustakas & Kennicutt (2006) shows an $H\alpha$ equivalent width of $< -1000\text{\AA}$ ($< -200\text{\AA}$). Simple stellar populations with such high equivalent widths have ages of only a few million years (c.f. Figure 23 of Leitherer et al. 2014.) Another indication of the extreme star formation occurring in these galaxies is the fact that Tololo 1214 – 277 is one of very few galaxies in the local universe that show indications of leaking Lyman continuum photons (Verhamme et al. 2015).

In addition, 7 $z \sim 0$ galaxies show strong C III] emission with $-14 < W_r < -5 \text{\AA}$. These are: NGC 5253 (regions HIIR-1 and HIIR-2), NGC 4861, NGC 5457 (region NGC 5471), Tololo 1345-420, Sbs 1319+579, III Zw 107, and Arp 252.

3.4. Lyman α

Shapley et al. (2003) noted a correlation between the equivalent widths of C III] and Lyman α emission in their quartile composite spectra. Stark et al. (2014) and Stark et al. (2015) further examined this correlation. In Figure 4, we plot $W_r(C\ III])$ versus $W_r(Ly\alpha)$ for galaxies at $1.6 < z < 7$ as well as galaxies at $z \sim 0$. A correlation is apparent for both redshift ranges, though there is considerable scatter. The correlation is dominated by the strongest emitters, with $W(Ly\alpha) \lesssim -50 \text{\AA}$, $W(C\ III]) \lesssim -5 \text{\AA}$. Within the dynamic range of the MagE sample, no correlation is apparent.

3.5. Strongly star-forming galaxies that are **not** strong C III] emitters

Since the physical conditions of the lensed galaxy RCS0327 at $z = 1.70$ have been measured precisely, we

compare them to the derived physical parameters of the Stark et al. (2014) sample. RCS0327 has a measured ionization parameter $\log U = -2.84 \pm 0.06$ (Rigby et al. 2011, using the average extinction from Whitaker et al. 2014.) This value is consistent with ionization parameters measured for other lensed Lyman Break galaxies: SGAS J122651.3+215220 and SGAS J152745.1+065219 in the MagE sample (Wuyts et al. 2012a), the Cosmic Horseshoe and the Clone (Hainline et al. 2009), and cB58 (Pettini et al. 2002). The ionization parameters of these LBGs are 0.6 to 1 dex lower than those inferred by Stark et al. (2014) for four galaxies. RCS0327 has a measured stellar mass of $6.3 \pm 0.7 \times 10^9 M_\odot$, and a star formation rate of $30\text{--}50 M_\odot \text{yr}^{-1}$ (Wuyts et al. 2012a). Its specific star formation rate (sSFR) is $5 \times 10^{-9} \text{yr}^{-1}$, such that it lies a factor of 3 above the correlation between star formation rate and stellar mass at that redshift. Three objects in Stark et al. (2014) have lower sSFRs than this; all have very strong C III] emission with $W_r \leq -10\text{\AA}$. Of the four galaxies in Stark et al. (2014) with highest stellar mass, $8.9 < \log[M_*/M_\odot] < 9.2$, three have C III] equivalent widths $< -4\text{\AA}$. Thus, RCS0327 is similar to the Stark et al. (2014) sample in terms of sSFR, but has a higher metallicity and lower ionization parameter than inferred for that sample. This comparison provides some support for the hypothesis of Stark et al. (2014), that both low metallicity and high ionization parameter are required to create strong C III] emission.

Perhaps more surprising is the fact that the vigorously star-forming, very low metallicity starburst galaxy 1 Zw 18 is *not* a strong C III] emitter; its equivalent width in three different regions varies from -1.3 to -4.4\AA . Its ionization parameter is high for the local universe ($\log U \sim -2.5$ from Dufour et al. (1988); inferred as -2.2 as the lowest-metallicity object in Morales-Luis et al. 2014). Any hypothesis that explains strong C III] emission will need to explain why 1 Zw 18 shows relatively weak C III] emission.

4. LOOKING FORWARD

C III] emission has emerged as a practical method to spectroscopically confirm galaxies at the reionization epoch, should Ly α emission prove to be suppressed. Our MagE spectra of lensed $1.6 < z < 3$ star-forming galaxies show that C III] emission typically has modest equivalent width in bright, high surface brightness galaxies at that epoch; this result is consistent with the composite spectra of Shapley et al. (2003). Strong C III] emission is clearly not a universal feature of galaxies at that epoch; equivalent widths exceeding -5\AA are seen in 45% of the $z \sim 2\text{--}4$ galaxies examined here.

We have examined C III] emission in local star-forming galaxies with archival *HST* spectra; a minority (22%) show strong C III] emission. Three of these nearby galaxies show C III] emission as strong or stronger ($< -14 \text{\AA}$) than yet reported for any galaxy at $z \sim 2$. The distribution function of equivalent width of C III] emission at any redshift may contain a tail toward very high equivalent widths; if so, by compiling spectra for 46 nearby galaxies, we have sampled this tail. The high equivalent width tail may be more populated at $z \sim 2$ than at $z \sim 0$, but determining this would require matched samples at both redshifts, which has not yet been done. The sample

¹⁰ Many studies incorrectly name this object; see <http://cdsarc.u-strasbg.fr/viz-bin/vizExec/.getpuz?7239&N%202363>

of Stark et al. (2014) clearly samples this tail of strong emission, while the MagE galaxies reported here do not.

The question then becomes, what physical conditions cause galaxies to be strong C III] emitters? For local galaxies, we see that metallicity may set an envelope on maximum C III] equivalent width. While low metallicity galaxies show a wide range on C III] equivalent width, strong emission is only seen at low metallicity. Additional physical conditions must be required to produce strong C III] emission. The three $z \sim 0$ galaxies with extreme C III] emission, as strong as anything seen in the distant universe, are Wolf-Rayet galaxies with extremely high H α equivalent widths. Such extreme emission calls to mind the candidate $z \sim 8$ dropout galaxies whose *Spitzer*/IRAC colors are best explained by extreme equivalent widths of [O III] $\lambda\lambda 4959, 5007$ and H β (Labbé et al. 2013). Follow-up investigations of these nearby, low-metallicity extreme C III] emitters may thus reveal the physical conditions that foster such strong emission from star-forming galaxies, with consequences

for our understanding of star formation at the highest redshifts.

This paper includes data gathered with the 6.5 meter Magellan Telescopes located at Las Campanas Observatory, Chile. This research has made use of the NASA/IPAC Extragalactic Database (NED) which is operated by the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration. This paper includes observations made with the NASA/ESA Hubble Space Telescope and with the International Ultraviolet Explorer, obtained from the Data Archive at the Space Telescope Science Institute, which is operated by the Association of Universities for Research in Astronomy, Inc., under NASA contract NAS 5-26555. JR thanks the “First Carnegie Symposium in honor of Leonard Searle”. We thank G. Sonneborn and S. Heap for advice regarding *IUE* spectra. HD acknowledges support from the Research Council of Norway.

REFERENCES

- Abazajian, K. N., Adelman-McCarthy, J. K., Agüeros, M. A., et al. 2009, *ApJS*, 182, 543
- Asplund, M., Grevesse, N., Sauval, A. J., & Scott, P. 2009, *Annual Review of Astronomy and Astrophysics*, 47, 481
- Bayliss, M. B., Rigby, J. R., Sharon, K., Wuyts, E., Florian, M., Gladders, M. D., Johnson, T., & Oguri, M. 2014, *ApJ*, 790, 144
- Belokurov, V., Evans, N. W., Moiseev, A., King, L. J., Hewett, P. C., Pettini, M., Wyrzykowski, L., McMahon, R. G., Smith, M. C., Gilmore, G., Sánchez, S. F., Udalski, A., Kozlov, S., Zucker, D. B., & Walcher, C. J. 2007, *ApJ*, 671, L9
- Bian, F., Fan, X., Bechtold, J., McGreer, I. D., Just, D. W., Sand, D. J., Green, R. F., Thompson, D., Peng, C. Y., Seifert, W., Ageorges, N., Juette, M., Knierim, V., & Buschkamp, P. 2010, *ApJ*, 725, 1877
- Biretta, J., et al. 2015, *STIS Instrument Handbook*, Version 14.0, (Baltimore: STScI).
- Buckalew, B. A., Kobulnicky, H. A., & Dufour, R. J. 2005, *The Astrophysical Journal Supplement Series*, 157, 30
- Caruana, J., Bunker, A. J., Wilkins, S. M., Stanway, E. R., Lacy, M., Jarvis, M. J., Lorenzoni, S., & Hickey, S. 2013, *Monthly Notices of the Royal Astronomical Society*, 427, 3055
- Caruana, J., Bunker, A. J., Wilkins, S. M., Stanway, E. R., Lorenzoni, S., Jarvis, M. J., & Ebert, H. 2014, *Monthly Notices of the Royal Astronomical Society*, 443, 2831
- Christensen, L., Richard, J., Hjorth, J., Milvang-Jensen, B., Laursen, P., Limousin, M., Dessauges-Zavadsky, M., Grillo, C., & Ebeling, H. 2012, *Monthly Notices of the Royal Astronomical Society*, 427, 1953
- Dufour, R. J., Garnett, D. R., & Shields, G. A. 1988, *ApJ*, 332, 752
- Erb, D. K., Pettini, M., Shapley, A. E., Steidel, C. C., Law, D. R., & Reddy, N. A. 2010, *ApJ*, 719, 1168
- Faisst, A. L., Capak, P., Carollo, C. M., Scarlata, C., & Scoville, N. 2014, *ApJ*, 788, 87
- Giavalisco, M., Koratkar, A., & Calzetti, D. 1996, *ApJ*, 466, 831
- Guseva, N. G., Izotov, Y. I., Stasińska, G., Fricke, K. J., Henkel, C., & Papaderos, P. 2011, *Astronomy and Astrophysics*, 529, 149
- Hainline, K. N., Shapley, A. E., Kornei, K. A., Pettini, M., Buckley-Geer, E., Allam, S. S., & Tucker, D. L. 2009, *ApJ*, 701, 52
- Koester, B. P., Gladders, M. D., Hennawi, J. F., Sharon, K., Wuyts, E., Rigby, J. R., Bayliss, M. B., & Dahle, H. 2010, *The Astrophysical Journal Letters*, 723, L73
- Kunth, D. & Joubert, M. 1985, *Astronomy and Astrophysics*, 142, 411
- Labbé, I., Oesch, P. A., Bouwens, R. J., Illingworth, G. D., Magee, D., González, V., Carollo, C. M., Franx, M., Trenti, M., van Dokkum, P. G., & Stiavelli, M. 2013, *ApJ*, 777, L19
- Leitherer, C., Ekstrom, S., Meynet, G., Schaerer, D., Agienko, K. B., & Levesque, E. M. 2014, *The Astrophysical Journal Supplement Series*, 212, 14
- Leitherer, C., Tremonti, C. A., Heckman, T. M., & Calzetti, D. 2011, *The Astronomical Journal*, 141, 37
- Marshall, J. L., Burles, S., Thompson, I. B., Shectman, S. A., Bigelow, B. C., Burley, G., Birk, C., Estrada, J., Jones, P., Smith, M., Kowal, V., Castillo, J., Storts, R., & Ortiz, G. 2008, *Ground-based and Airborne Instrumentation for Astronomy II*. Edited by McLean, 7014, 169
- Morales-Luis, A. B., Perez-Montero, E., Sánchez Almeida, J., & Muñoz-Tuñón, C. 2014, *ApJ*, 797, 81
- Moustakas, J. & Kennicutt, R. C. 2006, *The Astrophysical Journal Supplement Series*, 164, 81
- Pentericci, L., Fontana, A., Vanzella, E., Castellano, M., Grazian, A., Dijkstra, M., Boutsia, K., Cristiani, S., Dickinson, M., Giallongo, E., Giavalisco, M., Maiolino, R., Moorwood, A., Paris, D., & Santini, P. 2011, *ApJ*, 743, 132
- Pentericci, L., Vanzella, E., Fontana, A., Castellano, M., Treu, T., Mesinger, A., Dijkstra, M., Grazian, A., Bradač, M., Conselice, C., Cristiani, S., Dunlop, J., Galametz, A., Giavalisco, M., Giallongo, E., Koekemoer, A., McLure, R., Maiolino, R., Paris, D., & Santini, P. 2014, *ApJ*, 793, 113
- Pettini, M. & Pagel, B. E. J. 2004, *Monthly Notices of the Royal Astronomical Society*, 348, L59
- Pettini, M., Rix, S. A., Steidel, C. C., Adelberger, K. L., Hunt, M. P., & Shapley, A. E. 2002, *ApJ*, 569, 742
- Rigby, J. R., Bayliss, M. B., Gladders, M. D., Sharon, K., Wuyts, E., & Dahle, H. 2014, *ApJ*, 790, 44
- Rigby, J. R., Wuyts, E., Gladders, M. D., Sharon, K., & Becker, G. D. 2011, *ApJ*, 732, 59
- Schaerer, D., Contini, T., & Pindao, M. 1999, *Astronomy and Astrophysics Supplement Series*, 136, 35
- Shapley, A. E., Steidel, C. C., Pettini, M., & Adelberger, K. L. 2003, *ApJ*, 588, 65
- Small, I., Swinbank, A. M., Richard, J., Ebeling, H., Kneib, J.-P., Edge, A. C., Stark, D., Ellis, R. S., Dye, S., Smith, G. P., & Mullis, C. 2007, *ApJ*, 654, L33
- Stark, D. P., Richard, J., Charlot, S., Clément, B., Ellis, R., Siana, B., Robertson, B., Schenker, M., Gutkin, J., & Wofford, A. 2015, *Monthly Notices of the Royal Astronomical Society*, 450, 1846
- Stark, D. P., Richard, J., Siana, B., Charlot, S., Freeman, W. R., Gutkin, J., Wofford, A., Robertson, B., Amanullah, R., Watson, D., & Milvang-Jensen, B. 2014, *Monthly Notices of the Royal Astronomical Society*, 445, 3200
- Tilvi, V., Papovich, C., Finkelstein, S. L., Long, J., Song, M., Dickinson, M., Ferguson, H. C., Koekemoer, A. M., Giavalisco, M., & Mobasher, B. 2014, *ApJ*, 794, 5

TABLE 1

name	z	Z	ref	$W_r(Ly\ \alpha)$ (Å)	σ (Å)	ref	$W_r(C\ III])$ (Å)	σ (Å)	ref	sample	m_{AB}	ref
RCSGA 032727–13260 knotE	1.703745	8.34 ± 0.02	W14	> -1.2	-99	R14	-2.0	0.14	R14	MagE	$g = 19.15$	W10
SGAS J0004–0103	1.6811	$8.30^{+0.06}_{-0.08}$	FIRE	-3.3	0.4	R14	-0.45	0.14	R14	MagE	$g = 19.91$	A09
Pox 120	0.020748	7.83	G96	-70.2	4.11	IUE	-14.4	1.8	IUE	G96		
IRAS 08339+6517	0.019113	8.31	PG15	-4.45	0.5	STIS	-0.7	0.1	STIS	PG15		
NGC 4861	0.002785	7.9	L11	—	—	—	-8.1	1.0	FOS	L11		

NOTE. — Columns: 1) source name, 2) redshift, 3) $12 + \log(O/H)$ metallicity, 4) reference; 5) Rest-frame Ly α equivalent width; 6) uncertainty; 7) reference; 8) Rest-frame C III] equivalent width. Sum of both transitions when resolved; 9) uncertainty; 10) reference; 11) sample; 12) magnitude for MagE sample; 13) reference. Redshifts for MagE galaxies are from Bordoloi et al. 2015 (submitted to MNRAS) for RCS0327, else from Rigby et al. (2014) and in prep. Redshifts for $z \sim 0$ galaxies are from NED. Reference code: A09 Abazajian et al. (2009); B07 Belokurov et al. (2007); B10 Bian et al. (2010); G96 Gialalisco et al. (1996); H09 Hainline et al. (2009); K10 Koester et al. (2010); L11 Leitherer et al. (2011); PG15 Peña-Guerrero et al. (in prep.), R14 Rigby et al. (2014); S07 Smail et al. (2007); W10 Wuyts et al. (2010); W12 Wuyts et al. (2012b); W14 Wuyts et al. (2014). Table 1 is published in its entirety in the electronic edition of ApJ Letters. A portion is shown here for guidance regarding its form and content.

Treu, T., Schmidt, K. B., Trenti, M., Bradley, L. D., & Stiavelli, M. 2013, ApJ, 775, L29

Treu, T., Trenti, M., Stiavelli, M., Auger, M. W., & Bradley, L. D. 2012, ApJ, 747, 27

Verhamme, A., Orlitová, I., Schaerer, D., & Hayes, M. 2015, Astronomy and Astrophysics, 578, A7

Whitaker, K. E., Rigby, J. R., Brammer, G. B., Gladders, M. D., Sharon, K., Teng, S. H., & Wuyts, E. 2014, ApJ, 790, 143

Wuyts, E., Barrientos, L. F., Gladders, M. D., Sharon, K., Bayliss, M. B., Carrasco, M., Gilbank, D., Yee, H. K. C., Koester, B. P., & Muñoz, R. 2010, ApJ, 724, 1182

Wuyts, E., Rigby, J. R., Gladders, M. D., Gilbank, D. G., Sharon, K., Gralla, M. B., & Bayliss, M. B. 2012a, ApJ, 745, 86

Wuyts, E., Rigby, J. R., Gladders, M. D., & Sharon, K. 2014, ApJ, 781, 61

Wuyts, E., Rigby, J. R., Sharon, K., & Gladders, M. D. 2012b, ApJ, 73




FIG. 1.— Histograms of C III] $\lambda\lambda 1907, 1909$ Å rest-frame equivalent width. The $z \sim 0$ sample (grey filled histogram) combines galaxies from Leitherer et al. (2011), Peña-Guerrero et al. (in prep.), and Giavalisco et al. (1996). Two $z \sim 2$ samples are shown: the MagE spectral atlas (solid black line), including the $z = 3.6$ galaxy from Bayliss et al. (2014); and the galaxies from Stark et al. (2014) (dashed black line.) Asterisks shows equivalent widths for the composite spectrum and composite quartile spectra of Shapley et al. (2003), with 1σ uncertainty, at arbitrary y values. At both $z \sim 0$ and $z \sim 2$, the mode is small, and there is a tail to large equivalent widths.




FIG. 2.— Rest-frame C III] equivalent width as a function of gas-phase metallicity. Nearby galaxies are plotted in black: Leitherer et al. (2011) (black squares); the STIS sample of Peña-Guerrero et al. (in prep.) (black triangles); and Giavalisco et al. (1996) (black diamonds). Redshift $z \sim 2-4$ galaxies are plotted with colored symbols: SGAS J105039.6+001730 at $z = 3.6$ (Bayliss et al. 2014) (purple triangle); four physically-distinct star-forming regions within lensed galaxy RCS0327 at $z = 1.70$ (light blue squares); four additional galaxies at $1.6 < z < 2.8$ with MagE spectra and measured metallicities (purple squares and upper limit); four galaxies from Stark et al. (2014) with measured metallicity, and one galaxy (BX418) from Erb et al. (2010) (blue squares). A dashed line marks solar metallicity (Asplund et al. 2009). At large equivalent width, systematic uncertainty in the continuum level dominates. Metallicity appears to set an envelope; low metallicity may be a necessary but not sufficient condition for high C III] equivalent width.




FIG. 3.— Spectra of C III] emitters at $z \sim 0$. The three $z = 0$ emitters with largest $W_r(\text{C III]})$ are plotted: Tololo 1214 – 277 (top left), the region Mrk 71 within galaxy NGC 2366 (top right), and Pox 120 (bottom left). For comparison, we also plot a spectrum with moderate emission, IRAS 08339+6517 (bottom right). The $W_r(\text{C III]})$ for Tol 1214–277 is larger in *HST*-FOS compared to *IUE*, which we attribute to aperture effects. Clearly, low-redshift galaxies can produce strong C III] emission.




FIG. 4.— *Left panel*: Comparison of rest-frame equivalent widths of Ly α and C III]. Nearby galaxies are plotted in black, and $1.6 < z < 7$ galaxies are plotted with other colors. Symbols are coded as in Figure 2, with additional values from Stark et al. (2014) (orange circles), Stark et al. (2015) (red asterisks), Erb et al. (2010) (cyan triangle), and Shapley et al. (2003) (green squares). Two *IUE* spectra were available for each of IRAS 08339+6517 and NGC 1741; we plot a point for each spectrum. *Right panel*: Zoom of the region outlined in grey, containing most of our $z \sim 2$ MagE and $z \sim 0$ measurements. The equivalent widths are correlated, but the correlation is not obvious within the MagE sample.