




Trajectory Simulation of Meteors Assuming Mass Loss and Fragmentation

Gary A. Allen, Jr., Dinesh K. Prabhu, and David A. Saunders
ERC, Inc. at NASA Ames Research Center, Moffett Field, CA 94035, USA

1st International Workshop on PHA Characterization, Atmospheric Entry and Risk Assessment
7 – 9 July 2015, NASA Ames Research Center, California



Introduction and Objective

TRAJ Features:

- Program used to simulate atmospheric flight trajectories of entry capsules [1]
- Includes models of atmospheres of different planetary destinations – Earth, Mars, Venus, Jupiter, Saturn, Uranus, Titan, ...
- Solves 3-degrees of freedom (3DoF) equations for a single body treated as a point mass
- Also supports 6-DoF trajectory simulation and Monte Carlo analyses
- Uses Fehlberg-Runge-Kutta (4th-5th order) time integration with automatic step size control
- Includes rotating spheroidal planet with gravitational field having a J₂ harmonic
- Includes a variety of engineering aerodynamic and heat flux models
- Capable of specifying events – heatshield jettison, parachute deployment, etc. – at predefined altitudes or Mach number
- Has material thermal response models of typical aerospace materials integrated

Modify trajectory simulation tool, TRAJ, to make it suitable for meteor entries including mass loss & fragmentation

Modifications Made to TRAJ for Meteor Simulation

- NASA's Galileo probe to Jupiter only one that experienced significant mass loss
- Entry capsule was a 45° sphere-cone with fully-dense carbon phenolic as heatshield material
- M. Tauber *et al.* [2] developed JAE code for simulation of Galileo probe (Jupiter entry)
- JAE logic incorporated into Traj
 - Sphere-cone shape replaced by sphere
 - Mass loss equation of meteor physics used
 - Allow input specification of heat of ablation, Q
 - Allow heat transfer coefficient to vary in time
 - Time-varying heat transfer coefficients from detailed flow computations curve fit as a function of altitude, velocity, and size

Test Case: Chelyabinsk [3]

Basic Assumptions:

Hyperbolic excess velocity: 15.0 km/s
Altitude at entry: 95.0 km
Relative velocity at entry: 19.0 km/s
Relative entry angle: -18.5 deg
Relative heading angle: -76.6 deg
Geographic latitude at entry: 54.5 deg
Oblate rotating Earth
Gravitational model includes J₂ term
US-1976 atmospheric model

Meteoroid Assumptions:

Shape: Sphere
Density of meteoric material: 3300 kg/m³
Aerodynamic model: Sphere


Sensitivity study to entry mass, heat transfer coefficient, heat of ablation, and fragmentation

Heat Transfer Coefficient, C_H, Model


$$C_H(z) = \left[a + b(z-c)^d \right] \exp\left(-\frac{z-c}{f}\right)$$

- Curve fit expressions are to be used for z > 15.5 km
- C_H for different velocities and diameters obtained through linear interpolation

An example "quality of fit" plot generated with curve fit.



Trajectory Simulation Process with Meteor Physics Equations



Mass Change Equation

$$\frac{dm_m}{dt} = -\frac{1}{2} \rho_a u_m^3 \frac{C_H A_m}{Q}$$


t = Time
A₀ = Cross sectional area at entry
m_m = Mass at time t
u_m = Relative velocity at time t
ρ_a = Atmospheric density time t
Q = Heat of ablation
C_H = Heat transfer coefficient

Shape Change Equation (assuming spherical meteor)

$$\frac{A_m}{A_0} = \left(\frac{m_m}{m_0} \right)^{2/3}$$


Sensitivity to Basic Assumptions: Entry Mass, Fragmentation, Heat Transfer Coefficient and Heat-of-Ablation

Altitude versus Velocity Chelyabinsk Single Body Meteor Compared to Heat Transfer Coefficient Data Points



- Case 1_A: Single body with C_H = 0.1
 - Case 1_B: Single body with time varying C_H
 - Case 1_C: Time varying C_H for a single body. Entry mass and heat of ablation tuned to overlay Chelyabinsk observations
- Entry mass: 13 x 10⁶ kg
Meteor radius at entry: 9.8 m
Heat-of-Ablation: 8 MJ/kg
- Entry mass: 6.8 x 10⁴ kg
Meteor radius at entry: 1.7 m
Heat-of-Ablation: 0.85 MJ/kg


Altitude versus Velocity for Chelyabinsk Fragmented Meteor Compared to Heat Transfer Coefficient Data Points



- Case 2_A: Single fragmentation event at 40 km, and C_H = 0.1
 - Case 2_B: Two fragmentation events at 40 & 30 km, and C_H = 0.1
 - Case 2_C: Single fragmentation event at 40 km, and C_H time varying
 - Case 2_D: Two fragmentation events at 40 and 30 km, and C_H time varying
- For Cases 2_A – 2_D: Meteor radius: 3.5 m at 40 km alt. Revised mass: 5.93 x 10⁵ kg
- For Cases 2_B & 2_D: Meteor radius: 0.7 m at 30 km alt. Revised mass: 4.74 x 10³ kg

- Myriad ways to fit observations by choice of model parameters and fragmentation events
- Problem compounded by the fact that exo-atmospheric dynamical mass not known precisely

Basic Plots for Variable C_H and Double Fragmentation (Case 2_D)



- For Case 2_D simulation fragmentation at 40 & 30 km altitudes assumed to occur instantly
- Fragment masses tuned to overlay simulated trajectory on Chelyabinsk observations.
- On a scale of 40 to 90 km altitude, mass vs altitude trace appears to be a straight line over the entire mass range
- Trace is actually parabolic when mass scale is expanded
- Influence of C_H model is insignificant if large changes occur in meteor mass due to fragmentation.

Conclusions, Future Work and References

- TRAJ, an established trajectory simulation tool successfully modified for meteor entries
- Improvements include:
 - Simple mass loss equation of meteor physics
 - Time-varying heat transfer coefficient based on detailed flow computations
 - Ability to specify fragmentation events
- Updated version of TRAJ tested against Chelyabinsk observations
- TRAJ can now be used to establish sensitivity of trajectories to various meteor parameters
- Leaves open the issue of verification/validation of TRAJ and additional test cases are needed
- Could tektites [4] be used as additional test cases?
- Advantages of simulating tektite entries into Earth's atmosphere
 - Exo-atmospheric shapes are definitely spherical
 - Small sizes and (sub)orbital entry velocities
 - Problem is dominated by convective heating and melting
 - Melted shapes are aerodynamically stable
 - Chemical composition of australite tektites is statistically well defined
 - Serve as a good foundation for the tougher meteor entry problem



Recovered australite tektite

References:

1. Gary A. Allen, Jr., Michael J. Wright, and Peter Gage, "The Trajectory Program (Traj): Reference Manual and User's Guide," NASA TM-2004-212847, 2005.
2. Michael E. Tauber, Paul Wercinski, Lily Yang, and Yih-Kanq Chen, "A Fast Code for Jupiter Atmospheric Entry Analysis," NASA/TM-1999-208796, September 1999.
3. Jiri Borovicka, *et al.*, "The trajectory, structure and origin of the Chelyabinsk asteroidal impactor," *Nature*, **503**, 235-237, 14 November 2013.
4. George Baker, "Structures of Well Preserved Australite Buttons from Port Campbell, Victoria, Australia," *Meteoritics*, **3**(4), December 1967.