National Aeronautics and Space Administration



Observing strategies for focused, faint orbital debris surveys using the Magellan telescope

James Frith¹ September 2015

Patrick Seitzer², Phillip Anz-Meador³, Heather Cowardin¹, Brent Buckalew³, Susan Lederer⁴

¹Univ. Texas El Paso ²University of Michigan ³JETS/Jacobs ⁴NASA Johnson Space Center







Magellan Baade Telescope





6.5 Meter Telescope located at LasCampanas Observatory


- 2 half nights granted for orbital debris studies
- Attempted a more focus debris search than previously


→ Searching specifically for small debris (<10cm) that may have come about from recent Titan Transtage breakup

Titan Transtage Breakup



- 4 June at approximately 02:38 UT, a provisional breakup of a Titan 3C Transtage rocket body (SSN# 3692, International Designator 1969-013B) occurred
- >45 years after its launch
- The breakup has reportedly five observed fragments.
- This is the fourth breakup of this class





Focused Search Strategy




- Debris cloud produced using models
- Fragment orbits were propagated forward in time
- Expected debris cloud density was calculated for sky above Magellan over observing window

Magellan Tasking Strategy




3692 Predicted Debris Cloud Density Over Magellan for Night of UT July 27th - 28th and Background Stars



National Aeronautics and Space Administration






3692 Predicted Debris Cloud Density > 5cm Over Magellan for Night of UT July 26th - 27th and Background Stars

Magellan Tasking Strategy




3692 Debris Cloud Ra Dec Rates



National Aeronautics and Space Administration





Observations



• 26-28 July

- Magellan run very successful. 0.5 night of 3 night run shared with another project
- Observations 5 seconds through r filter until moonset then broad 4800-7800 filter using the IMACS f/2 instrument (0.5 degree diameter field)
- Conditions clear and probably photometric, seeing < 0.8 arcsec FWHM.
- 20 sequences of 16 images each rate tracked at expected positions and rates of maximum of debris population. Take 2 images while sidereal tracked to establish focus, then 16 at rate track.
- 15 sequences at first initial RA, DEC; 5 sequences at second initial RA, DEC.


• 30 Oct – 1 Nov

• Next observing run using Baade

Eyeball results



- Object ~16-17th mag
- Moving 1 arc-sec/sec in EW direction with respect to predicted debris motion
- Object at upper edge of frame in first two images, at extreme right of frame in last image.
- Visually detected object at telescope in between in-between attempting to set personal paddle ball record.



Rest of processing in progress...