

## Note

# Low-temperature thermal reactions between SO<sub>2</sub> and H<sub>2</sub>O<sub>2</sub> and their relevance to the jovian icy satellites

Mark J. Loeffler\*, Reggie L. Hudson

NASA Goddard Space Flight Center, Astrochemistry Branch, Code 691, Greenbelt, MD 20771, USA

## ARTICLE INFO

### Article history:

Received 20 December 2012

Revised 11 February 2013

Accepted 11 February 2013

Available online 21 February 2013

### Keywords:

Europa

Ices, IR spectroscopy

Jupiter, Satellites

## ABSTRACT

Here we present first results on a non-radiolytic, thermally-driven reaction sequence in solid H<sub>2</sub>O + SO<sub>2</sub> + H<sub>2</sub>O<sub>2</sub> mixtures at 50–130 K, which produces sulfate (SO<sub>4</sub><sup>2-</sup>), and has an activation energy of 53 kJ/mole. We suspect that these results may explain some of the observations related to the presence and distribution of H<sub>2</sub>O<sub>2</sub> across Europa's surface as well as the lack of H<sub>2</sub>O<sub>2</sub> on Ganymede and Callisto.

Published by Elsevier Inc.

## 1. Introduction

The radiation-driven chemistry of surface and near-surface ices of Europa has been studied by several research groups in recent years. In contrast, thermally-driven reactions in outer Solar System ices have remained relatively unexplored. In this note we report new laboratory experiments demonstrating that a straightforward sequence of hydrolysis, protonation, and redox chemistry can convert SO<sub>2</sub> into SO<sub>4</sub><sup>2-</sup> (sulfate) at the temperatures of Europa's surface and in the presence of H<sub>2</sub>O<sub>2</sub> (hydrogen peroxide), a well-known radiation product of H<sub>2</sub>O-ice. These reactions suggest that sub-surface SO<sub>4</sub><sup>2-</sup> formation will occur in the absence of the direct reach of jovian magnetospheric radiation.

Laboratory studies of H<sub>2</sub>O-ice, one of the more abundant substances on cold surfaces in outer space, have shown that radiation can cause both amorphization of crystalline H<sub>2</sub>O-ice (Hudson and Moore, 1995; Strazzulla et al., 1992) and the production of new molecules, such as H<sub>2</sub>O<sub>2</sub>, H<sub>2</sub>, and O<sub>2</sub>. Of these three species, H<sub>2</sub>O<sub>2</sub> (hydrogen peroxide) formation has been observed *in situ* by laboratory infrared (IR) spectroscopy in H<sub>2</sub>O-ice irradiated at temperatures relevant to extraterrestrial surfaces (Loeffler et al., 2006a; Moore and Hudson, 2000; Zheng et al., 2006). Although all of these prior studies showed that the H<sub>2</sub>O<sub>2</sub> abundance is relatively low in irradiated H<sub>2</sub>O-ice, the H<sub>2</sub>O<sub>2</sub> absorption at 3.5 μm is well-separated from the stronger OH stretching vibrations of water, making it an attractive candidate for observation on planetary surfaces if both H<sub>2</sub>O-ice and radiation are present.

Given the many extraterrestrial objects with surficial H<sub>2</sub>O-ice exposed to radiation, one might expect to find H<sub>2</sub>O<sub>2</sub> on a great number of icy bodies. However, H<sub>2</sub>O<sub>2</sub> has been detected unequivocally only on Europa (Carlson et al., 1999), with IR spectra of Ganymede and Callisto failing to show the 3.5-μm peroxide band. Interestingly, ultraviolet spectra of Ganymede do show O<sub>3</sub> (Noll et al., 1996), a reliable indicator of radiation processing of H<sub>2</sub>O-ice containing O<sub>2</sub>. Differences exist among surface environments of Europa, Ganymede, and Callisto, such as concerning radiation flux, average surface temperature, and composition, but so far none of these factors readily explains why Europa would be the only place where H<sub>2</sub>O<sub>2</sub> is present.

In recent years we have used IR spectroscopy to investigate the solid-phase radiation chemistry of H<sub>2</sub>O + SO<sub>2</sub> ices (Moore et al., 2007), solid sulfuric acid, and sulfuric acid hydrates (Loeffler and Hudson, 2012; Loeffler et al., 2011), all at temperatures relevant to the icy jovian satellites. With H<sub>2</sub>O + SO<sub>2</sub> ices, it was observed that radiation readily produces SO<sub>4</sub><sup>2-</sup> (sulfate ion), presumably by the oxidizing action of radiolytically-generated H<sub>2</sub>O<sub>2</sub>. This assumption has been tested in the present paper by studying IR spectra of *unirradiated* H<sub>2</sub>O + SO<sub>2</sub> + H<sub>2</sub>O<sub>2</sub> ices as they are warmed. We have observed that even at relatively low temperatures, H<sub>2</sub>O<sub>2</sub> and SO<sub>2</sub> react thermally in the presence of H<sub>2</sub>O-ice. The main products of the reaction have been identified and the reaction's rate has been measured at 110–122 K, temperatures found on the jovian icy satellites mentioned above.

## 2. Experimental methods

Experiments were performed with a cryostat ( $T_{\min} \sim 10$  K) operating in a stainless steel high-vacuum chamber ( $P \sim 1 \times 10^{-7}$  Torr). Ice films were prepared by co-deposition of H<sub>2</sub>O, H<sub>2</sub>O<sub>2</sub>, and SO<sub>2</sub> onto a pre-cooled (50–100 K) gold-coated aluminum mirror using three separate pre-calibrated gas lines. Pure H<sub>2</sub>O<sub>2</sub> was prepared in a glass manifold, using the technique described previously (Loeffler and Baragiola, 2011). During deposition, the sample's thickness was monitored with interferometry using a diode laser (670 nm). In all experiments the thickness of the resulting H<sub>2</sub>O + SO<sub>2</sub> + H<sub>2</sub>O<sub>2</sub> (80:14:6) ice was  $1.3 \pm 0.1$  μm, assuming indices of refraction at 670 nm are similar to those measured in the visible region, 1.31 for H<sub>2</sub>O (Merwin, 1930), 1.4 for H<sub>2</sub>O<sub>2</sub> (Giguère, 1943) and 1.36 for SO<sub>2</sub> (Musso et al., 2000), uniform mixing, and an ice density of 0.96 g/cm<sup>3</sup> (0.82 g/cm<sup>3</sup> for H<sub>2</sub>O (Westley et al., 1998), 1.6 g/cm<sup>3</sup> for H<sub>2</sub>O<sub>2</sub> (Loeffler et al., 2006b) and 1.49 g/cm<sup>3</sup> for SO<sub>2</sub> (Weast, 1983).

Each sample's IR spectrum was recorded before, during, and after warming at 1 K/min to the annealing temperature (110–130 K). Spectra of ices were measured from 7000 to 400 cm<sup>-1</sup> with a Bruker Vector Fourier Transform infrared spectrometer at 2-cm<sup>-1</sup> resolution and with 200-scan accumulations. To obtain a spectrum, the reflectance from the ice-coated substrate was ratioed against the reflectance of the bare substrate, taken before ice formation, and then converted to absorbance units.

\* Corresponding author.

E-mail address: [mark.loeffler@nasa.gov](mailto:mark.loeffler@nasa.gov) (M.J. Loeffler).

To quantify the kinetics of the thermal reactions occurring in our samples, we first fit the baseline of the  $2850\text{ cm}^{-1}$  overtone band of  $\text{H}_2\text{O}_2$  with a non-linear curve and then integrated this same  $\text{H}_2\text{O}_2$  IR feature. We chose to examine  $\text{H}_2\text{O}_2$  and not  $\text{SO}_2$  because the sublimation of pure  $\text{H}_2\text{O}_2$  is negligible below 130 K, whereas the sublimation of pure  $\text{SO}_2$  can become significant above 100 K, the temperature region of interest. We note that when  $\text{H}_2\text{O}_2$  is dispersed in water the strength of the  $2850\text{ cm}^{-1}$  band depends slightly on temperature (Loeffler et al., 2006a). This dependence was confirmed in  $\text{H}_2\text{O}$ – $\text{H}_2\text{O}_2$  calibration experiments and we made use of it in our calculations; the strength of the  $\text{H}_2\text{O}_2$  band in question drops linearly by about 20% between 50 and 130 K.

### 3. Results

Fig. 1 shows the IR spectrum of an  $\text{H}_2\text{O} + \text{SO}_2 + \text{H}_2\text{O}_2$  (80:14:6) ice after deposition at 50 K, during warming to 118 K, and while held at 118 K. At 50 K, each of the three molecules of the sample is easily identified in this spectral region:  $\text{H}_2\text{O}_2$  ( $2845\text{ cm}^{-1}$ ),  $\text{H}_2\text{O}$  ( $1655\text{ cm}^{-1}$ ), and  $\text{SO}_2$  ( $1329$  and  $1151\text{ cm}^{-1}$ ). On warming to 118 K, new absorptions, belonging to  $\text{HSO}_3^-$  ( $1035\text{ cm}^{-1}$ ),  $\text{SO}_4^{2-}$  ( $1092\text{ cm}^{-1}$ ), and  $\text{S}_2\text{O}_5^{2-}$  ( $954\text{ cm}^{-1}$ ) appeared, as did a broadening on the high-wavenumber side of the  $\text{H}_2\text{O}$  band at  $1655\text{ cm}^{-1}$ , indicating  $\text{H}_3\text{O}^+$  formation. As the sample was annealed at 118 K, the main  $\text{SO}_4^{2-}$  band continued to increase, other weaker bands of  $\text{SO}_4^{2-}$  ( $606\text{ cm}^{-1}$  and  $980\text{ cm}^{-1}$ ) appeared, and the  $\text{H}_2\text{O}_2$  and  $\text{SO}_2$  bands decreased significantly. Fig. 2 shows the  $\text{H}_2\text{O}_2$  abundance in the sample as a function of time (50 K;  $t = 0$ ) for annealing temperatures of 110, 118, and 130 K. In all cases the  $\text{H}_2\text{O}_2$  abundance began to decrease around 100 K and continued to fall as the sample was an-

nealed. For the samples held at 110 and 118 K, the  $\text{H}_2\text{O}_2$  abundance dropped by a factor of two and a factor of eight, respectively, after 1200 min, while only  $\sim 20$  min were needed at 130 K for the  $\text{H}_2\text{O}_2$  abundance to drop below the noise level. Blank experiments with  $\text{H}_2\text{O}$ – $\text{H}_2\text{O}_2$  showed that over these same time periods there was no  $\text{H}_2\text{O}_2$  loss, as expected from the negligible sublimation rate of  $\text{H}_2\text{O}_2$  at 130 K.

The inset of Fig. 2 shows an Arrhenius plot derived from six different annealing temperatures, which together yielded an activation energy of  $53 \pm 5\text{ kJ/mole}$ . The rate constant,  $k$ , was derived from  $N = N_0 \exp(-kt)$ , since we observed that, to a first approximation, the  $\text{H}_2\text{O}_2$  abundance ( $N$ ) dropped exponentially with time ( $t$ ) at each temperature studied. In the higher temperature experiments, the reaction began before we reached the annealing temperature, so in all cases we calculated  $k$  by equating it to  $\ln(2)/t_{1/2}$ , where  $t_{1/2}$  is the time at the annealing temperature needed for the  $\text{H}_2\text{O}_2$  abundance to drop to half of its original value. We note that as the abundance of  $\text{H}_2\text{O}_2$  has already dropped by a factor of two by the time we warm to  $\sim 125\text{ K}$  (see 130 K experiment in Fig. 2), we only used annealing temperatures below 122 K to calculate the activation energy. The error that we give in the activation energy is a combination of the difficulty of fitting the  $\text{H}_2\text{O}_2$  baseline, the narrow temperature range (12 K) for which we can obtain a value for  $t_{1/2}$ , and the fact that at the upper temperatures the reactions already have proceeded to some extent by the time the sample attained those temperatures. Future studies will focus on reducing the uncertainty in these measurements by extending the experiments to lower temperatures so that the activation energy can be more-confidently used below 100 K.

### 4. Discussion

#### 4.1. Reaction chemistry

Thermal reactions between  $\text{SO}_2$  and  $\text{H}_2\text{O}_2$  have been studied extensively within the atmospheric-chemistry community due to the importance of such chemistry in removing  $\text{SO}_2$ , in liquid, gas, and solid phases, from Earth's atmosphere. The primary reaction sequence is believed to begin with the formation of bisulfite ( $\text{HSO}_3^-$ ) through (1)



where the  $\text{H}^+$  either rapidly or in concerted fashion attaches to a second  $\text{H}_2\text{O}$  molecule to form hydronium:



In our previous studies with  $\text{H}_2\text{O} + \text{SO}_2$  ices, we found that these thermal reactions occur in the solid state even at temperatures below 100 K (Loeffler and Hudson, 2010). We now find, in addition, that in the presence of  $\text{H}_2\text{O}_2$  the bisulfite is oxidized to form the sulfate ( $\text{SO}_4^{2-}$ ) ion (Martin and Damschen, 1981):



Fig. 1 provides evidence that these same thermally-induced reactions occur in our  $\text{H}_2\text{O} + \text{SO}_2 + \text{H}_2\text{O}_2$  ices. Beginning with the 50-K spectrum and working toward those for higher temperatures, IR absorptions of both  $\text{HSO}_3^-$  ( $1035\text{ cm}^{-1}$ ) and  $\text{SO}_4^{2-}$  ( $1092\text{ cm}^{-1}$ ) products can be seen starting at  $\sim 80\text{ K}$  during warming. Both bands continued to increase with temperature until the  $\text{SO}_4^{2-}$  feature was one of the more-prominent remaining IR absorptions. The increase in abundances for the  $\text{HSO}_3^-$  and  $\text{SO}_4^{2-}$  ions correlated with the decrease in absorption band areas of reactants  $\text{SO}_2$  and  $\text{H}_2\text{O}_2$  (Fig. 1), supporting the simple reaction sequence already described. Comparing the  $\text{H}_2\text{O}_2$  loss for our different annealing temperatures, we see that not surprisingly the reaction proceeds the fastest at the highest temperature. However, even at the lower temperatures, the  $\text{H}_2\text{O}_2$  abundance is still decreasing after the longest times studied ( $\sim 1200$  min), indicating that given enough time all of the  $\text{H}_2\text{O}_2$  will convert to  $\text{SO}_4^{2-}$  via reaction (3).

#### 4.2. Icy satellite implications

Infrared spectra of the surface of Europa, Ganymede, and Callisto all contain an absorption at  $4.05\text{ }\mu\text{m}$ , which usually is assigned to  $\text{SO}_2$  (Hansen and McCord, 2008; Hibbitts et al., 2000; McCord et al., 1998). In contrast, the  $3.5\text{-}\mu\text{m}$  band of  $\text{H}_2\text{O}_2$  has been found only in Europa's spectra (Carlson et al., 1999; Hansen and McCord, 2008). Our experiments show that the abundance of  $\text{H}_2\text{O}_2$  in these surface ices depends on the presence of  $\text{SO}_2$ , and that, conversely, sulfur dioxide's abundance will be influenced by the presence of  $\text{H}_2\text{O}_2$ . Given these new laboratory results, we now turn to some of the previous observations of the jovian icy satellites.

On Europa, although both  $\text{H}_2\text{O}_2$  and  $\text{SO}_2$  have been detected they do not appear to be uniformly distributed. Hansen and McCord (2008) determined that the  $\text{SO}_2$  infrared feature was present on Europa's trailing hemisphere. This agrees with previous measurements of Europa's ultraviolet reflectance spectrum (Domingue and Lane, 1998; Hendrix et al., 2011; Hendrix and Johnson, 2008; Lane et al., 1981), which showed a strong slope on the moon's trailing side, attributed to  $\text{SO}_2$  ( $0.28\text{ }\mu\text{m}$  band center). It also is consistent with the sulfur implantation rate being an order of magnitude higher on Europa's trailing hemisphere (Johnson et al., 2004).




Fig. 1. Evolution of a  $\text{H}_2\text{O} + \text{SO}_2 + \text{H}_2\text{O}_2$  sample (80:14:6) during heating to and annealing at 118 K. The bottom four spectra are labeled by the time elapsed after the sample had reached 118 K.




Fig. 2. Evolution of a  $\text{H}_2\text{O} + \text{SO}_2 + \text{H}_2\text{O}_2$  sample (80:14:6) during heating to and isothermal annealing at 110, 118, and 130 K. All samples were deposited at 50 K and warmed at 1 K/min. Inset: Arrhenius plot for the  $\text{H}_2\text{O} + \text{SO}_2 + \text{H}_2\text{O}_2$  reaction. The line's slope yields an activation energy of  $53\text{ kJ/mole}$ .

Interestingly, the more-recent ultraviolet measurements indicate that SO<sub>2</sub> is not only more abundant on the trailing side but that it is actually absent on the leading side, where the majority of the H<sub>2</sub>O<sub>2</sub> detections have been made (Carlson et al., 1999). Thus, based on our results it seems plausible that the distribution of SO<sub>2</sub> and H<sub>2</sub>O<sub>2</sub> may be a result of excess SO<sub>2</sub> consuming any H<sub>2</sub>O<sub>2</sub> produced by radiolysis in the trailing hemisphere and excess H<sub>2</sub>O<sub>2</sub> consuming the SO<sub>2</sub> formed by implantation in the leading hemisphere. Finally, we point out that one recent study did detect both H<sub>2</sub>O<sub>2</sub> and SO<sub>2</sub> in a spectrum of an ice-rich region on Europa's trailing side (Hansen and McCord, 2008). However, it is possible that these two molecules may be spatially separated, but sufficiently close to lie within the same pixel area of the detector.

On Callisto and Ganymede the 3.5- $\mu$ m absorption diagnostic of H<sub>2</sub>O<sub>2</sub> is absent, which could indicate that surficial SO<sub>2</sub> is widespread or that the H<sub>2</sub>O<sub>2</sub> abundance is simply much lower than on Europa (Hendrix et al., 1999; Hendrix and Johnson, 2008). The former possibility is consistent with IR measurements of Ganymede and Callisto (McCord et al., 1998), which showed that each satellite's spectrum contained the 4.05- $\mu$ m absorption band attributed to SO<sub>2</sub>. However, we note that there has been some discussion as to whether another species, such as H<sub>2</sub>CO<sub>3</sub> or other carbonates (Johnson et al., 2004), could provide an adequate match for the 4.05- $\mu$ m feature in Callisto's spectrum, as this band does not seem to be correlated with magnetospheric bombardment (Hibbitts et al., 2000). Also, the most recent ultraviolet measurements show little variation across Callisto's surface (Hendrix and Johnson, 2008).

As the H<sub>2</sub>O + SO<sub>2</sub> + H<sub>2</sub>O<sub>2</sub> reaction we studied occurs quickly at temperatures relevant to these icy satellites ( $t_{1/2} \sim 1$  yr at 100 K and  $\sim 0.3$  h at 120 K), we expect that it plays an important role in the evolution of the jovian icy satellites' surface chemistry and may explain observations of Europa related to the presence and distribution of H<sub>2</sub>O<sub>2</sub> as well as the lack of H<sub>2</sub>O<sub>2</sub> on Ganymede and Callisto. If other molecules prove to be reactive with H<sub>2</sub>O<sub>2</sub> at these or at even lower temperatures, then similar thermal chemistry may explain why H<sub>2</sub>O<sub>2</sub> has not been detected on most Solar System icy bodies exposed to radiation. Future laboratory studies will focus on extending the reported measurements to other temperatures and concentrations, identifying and quantifying other thermal reactions that may occur between H<sub>2</sub>O<sub>2</sub> and other astrochemically-relevant molecules, and establishing whether the reaction products contain IR absorptions which could be detected by remote sensing. Finally, we note that the reactions we have described are not restricted to the surface of Europa and other worlds. To the extent that vertical transport of radiolytic products occurs (Greenburg, 2010), our reactions (1)–(3) also will take place beneath the IR-sensed layer of Europa, providing a thermally-driven source of SO<sub>4</sub><sup>2-</sup>.

#### Acknowledgments

The support of NASA's Planetary Geology and Geophysics program is gratefully acknowledged. R.L.H. also acknowledges support from the NASA Astrobiology Institute through the Goddard Center for Astrobiology.

#### References

- Carlson, R.W. et al., 1999. Hydrogen peroxide on the surface of Europa. *Science* 283, 2062–2064.
- Domingue, D.L., Lane, A.L., 1998. IUE views Europa: Temporal variations in the UV. *Geophys. Res. Lett.* 25, 4421–4424.
- Giguère, P.A., 1943. The refractive indices of hydrogen peroxide and its aqueous solutions. *Can. J. Res.* 21b, 156–162.
- Greenburg, R., 2010. Transport rates of radiolytic substances into Europa's ocean: Implications for the potential origin and maintenance of life. *Astrobiology* 10, 275–283.
- Hansen, G.B., McCord, T., 2008. Widespread CO<sub>2</sub> and other non-ice compounds on the anti-jovian and trailing sides of Europa from Galileo/NIMS observations. *Geophys. Res. Lett.* 35, L01202.
- Hendrix, A.R., Johnson, R.E., 2008. Callisto: New insights from Galileo disk-resolved UV measurements. *Astrophys. J.* 687, 706–713.
- Hendrix, A.R., Barth, C.A., Stewart, A.I.F., Hord, C.W., Lane, A.L., 1999. Hydrogen peroxide on the icy Galilean satellites. *Lunar Planet. Sci.* 30, 2043 (Abstract).
- Hendrix, A.R., Cassidy, T.A., Johnson, R.E., Paranicas, C., Carlson, R.W., 2011. Europa's disk-resolved ultraviolet spectra: Relationships with plasma flux and surface terrains. *Icarus* 212, 736–743.
- Hibbitts, C.A., McCord, T.B., Hansen, G.B., 2000. Distributions of CO<sub>2</sub> and SO<sub>2</sub> on the surface of Callisto. *J. Geophys. Res.* 105, 22541–22557.
- Hudson, R.L., Moore, M.H., 1995. Far-IR spectral changes accompanying proton irradiation of solids of astrochemical interest. *Radiat. Phys. Chem.* 45, 779–789.
- Johnson, R.E., Carlson, R.W., Cooper, J.F., Paranicas, C., Moore, M.H., Wong, M.C., 2004. Radiation effects on the surface of the Galilean satellites. In: Bagenal, F., Dowling, T., McKinnon, W.B. (Eds.), *Jupiter: The Planet, Satellites and Magnetosphere*. Cambridge University Press, Cambridge, pp. 485–512.
- Lane, A.L., Nelson, R.M., Matson, D.L., 1981. Evidence for sulphur implantation in Europa's UV absorption band. *Nature* 292, 38–39.
- Loeffler, M.J., Baragiola, R.A., 2011. Isothermal decomposition of hydrogen peroxide dihydrate. *J. Phys. Chem. A* 15, 5324–5328.
- Loeffler, M.J., Hudson, R.L., 2010. Thermally-induced chemistry and the jovian icy satellites: A laboratory study of the formation of sulfur oxyanions. *Geophys. Res. Lett.* 37, 19201.
- Loeffler, M.J., Hudson, R.L., 2012. Thermal regeneration of sulfuric acid hydrates after irradiation. *Icarus* 219, 561–566.
- Loeffler, M.J., Raut, U., Vidal, R.A., Baragiola, R.A., Carlson, R.W., 2006a. Synthesis of hydrogen peroxide in water ice by ion irradiation. *Icarus* 180, 265–273.
- Loeffler, M.J., Teolis, B.D., Baragiola, R.A., 2006b. Decomposition of solid amorphous hydrogen peroxide by ion irradiation. *J. Chem. Phys.* 124, 104702.
- Loeffler, M.J., Hudson, R.L., Moore, M.H., Carlson, R.W., 2011. Radiolysis of sulfuric acid, sulfuric acid monohydrate, and sulfuric acid tetrahydrate and its relevance to Europa. *Icarus* 215, 370–380.
- Martin, L.R., Damschen, D.E., 1981. Aqueous oxidation of sulfur dioxide by hydrogen peroxide at low pH. *Atmos. Environ.* 15, 1615–1621.
- McCord, T.B. et al., 1998. Non-water-ice constituents in the surface material of the icy Galilean satellites from the Galileo near-infrared mapping spectrometer investigation. *J. Geophys. Res.* 103, 8603–8626.
- Merwin, H.E., 1930. Refractivity of birefringent crystals. In: *International Critical Tables*. McGraw-Hill, New York, pp. 16–33.
- Moore, M.H., Hudson, R.L., 2000. IR detection of H<sub>2</sub>O<sub>2</sub> at 80 K in ion-irradiated laboratory ices relevant to Europa. *Icarus* 145, 282–288.
- Moore, M.H., Hudson, R.L., Carlson, R.W., 2007. The radiolysis of SO<sub>2</sub> and H<sub>2</sub>S in water ice: Implications for the icy jovian satellites. *Icarus* 189, 409–423.
- Musso, M., Aschauer, R., Asenbaum, A., Vasi, C., Wilhelm, E., 2000. Interferometric determination of the refractive index of liquid sulphur dioxide. *Meas. Sci. Technol.* 11, 1714–1720.
- Noll, K.S., Johnson, R.E., Lane, A.L., Domingue, D.L., Weaver, H.A., 1996. Detection of ozone on Ganymede. *Science* 273, 341–343.
- Strazzulla, G., Baratta, G.A., Leto, G., Foti, G., 1992. Ion-beam-induced amorphization of crystalline water ice. *Europhys. Lett.* 18, 517–522.
- Weast, R.E. (Ed.), 1983. *Handbook of Chemistry and Physics*. CRC Press, Boca Raton.
- Westley, M.S., Baratta, G.A., Baragiola, R.A., 1998. Density and index of refraction of water ice films vapor deposited at low temperatures. *J. Chem. Phys.* 108, 3321–3326.
- Zheng, W., Jewitt, D., Kaiser, R.I., 2006. Formation of hydrogen, oxygen, and hydrogen peroxide in electron-irradiated crystalline water ice. *Astrophys. J.* 639, 534–548.