Speed of CMEs and the Magnetic Non-potentiality of their Source A ve Regions

Sanjiv K. Tiwari[1*], David A. Falconer[1,2], Ronald L. Moore[1,2], P. Venkatakrishnan[3], Amy R. Winebarg or G. Khazanov[2]

[1]NASA Marshall Space Flight Center, ZP 13, Huntsville, AL 35812, USA

[2]Center for Space Plasma and Aeronomic Research, University of Alabama in Huntsville, Huntsville, AL 358

[3]Udaipur Solar Observatory, Physical Research Laboratory, Udaipr 313001, India

Abstract

Most fast coronal mass ejections (CMEs) originate from solar active regions (ARs). Non-potentiality of ARs plausibly determines the speed of CMEs in the rona. Several other unexplored parameters might be important as well. To find out the relation between the initial speed of CMEs and the non-potentiality of source ARs, we identified over a hundred of CMEs rce ARs via their co-produced flares. The speed of the CMEs are collected from the SOHO LASCO CME catalog. We have used vector magnetograms obtained with HMI/SDO, to evaluate various magnetic non-pot parameters, e.g. magnetic free-energy proxies, twist, shear angle, signed shear angle, net current etc. We have also included several other parameters e.g. total unsigned flux, magnetic area of ARs, area of su to investigate their correlation, if any, with the initial speeds of CMEs. Our preliminary results show that the ARs with larger non-potentiality and area produce faster CMEs but they can also produce slow ones. Th ith lesser non-potentiality and area generally produce only slower CMEs.

CME & Active Region Selection Criterion

An Example: An X-class flare occurred in the upper-left at 00:02:00 UT; 22 minutes later a CME appeared. The largest AR in that area is 11429, which produced flare/CME eruption.







To confirm the source AR of the CME, first we look at STEREO-A & B movies to make sure that the CME is Earth-directed. We then look at AIA movies, and SolarMonitor.org to find flare activity & timing, and NOAA AR number & position of the AR on the solar disk.





AR Magnetic Non-potentiality Parameters:

Signed Shear Angle (SSA): $SSA = \tan^{-1}\left(\frac{B_{yo}B_{xp} - B_{yp}B_{xo}}{B_{xo}B_{xp} + B_{yo}B_{yp}}\right)$: Tiwari et al 2009, ApJL;

Force-free parameter (α_g): $\alpha_g = \frac{\sum\left(\frac{\partial B_y}{\partial z} - \frac{\partial B_z}{\partial y}\right)B_z}{\sum B_z^2}$: Tiwari et al 2009, ApJ;

Gradient weighted neutral line length (WL_{SG}): $WL_{SG} = \int |\nabla Bz| dl$: Falconer et al 2008, ApJ;

R

CME Speed vs Various Magnetic Non



(a free-energy proxy)

SASSA: Spatially Averaged SSA

MSSA : Median of SSA: both SASSA and MSSA measure twist at a given height irrespective of shape and force-free nature of spot

α_g : measures gradient of twist per unit axial length; a proxy for global twist

ϕ : total unsigned magnetic flux, this is proportional to area of the AR

WL_{SG} is a free energy proxy

iality Parameters of the Source AR



Discussion and Conclusions

A right-angled triangle shape of the scatter plots of CMEs speed vs free energy parameters is most noticeable for MSSA × ϕ, giving the sharpest CME speed lim sing with AR free energy. Each of the other three free-energy proxies shown here shows the CME speed limit better than its corresponding twist proxy. Evidently the fastest CMEs are produced by ARs having the ee energy. Such ARs also produce many slower CMEs. Least non-potential ARs do not produce fast CMEs. We are collecting more data to reduce scatter and improve statistics. One caution is that the LASCO CME s used here are linear fit plane-of-the-sky values, and therefore underestimated.

*sanjiv.k.tiwari@nasa.gov

SKT is supported by an appointment to the NASA Postdoctoral Program at the NASA MSFC, administered by ORAU through a contract with NASA. RLM and ARW is supported by funding LWS TRT Program of the Heliophysics Division of NASA's SMD. Support for MAG4 development comes from NASA's Game Changing Development Program, and Johnson Space Center's Space Radiation Analysis Group (SRAG). Use of SOHO LASCO CME nd data from AIA+HMI (SDO), STEREO, SOHO is acknowledged.