

Hydrogen (H) Isotope Composition of Type II Kerogen Extracted by Pyrolysis-GC-MS-IRMS: Terrestrial Shale deposits as Martian Analogs

To be submitted to the AGU Fall Meeting, session VO36, Shale Reservoir Science & Technology, session ID#2918, Abstract ID#26264

Richard A. Socki, NASA-JSC Jacobs-JETS, 2224 Bay Area Blvd., Houston, TX 77058, **Denet Pernia**, University of Houston, Center for Petroleum Geochemistry, Houston, TX 77204, **Michael Evans**, Texas A&M University, Dept of Oceanography, College Station, TX 77834; **Qi Fu**, University of Houston, Dept. of Earth & Atmospheric Sciences, Houston, TX 77058, **Kadry K. Bissada**, University of Houston, Center for Petroleum Geochemistry, Houston, TX 77204, **Joseph A. Curiale**, Chevron, Houston, TX 77002, and **Paul B. Niles**, NASA-JSC, Code KR, Houston, TX 77058.

Described here is a technique for H isotope analysis of organic compounds pyrolyzed from kerogens isolated from gas- and liquids-rich shales. Application of this technique will progress the understanding of the use of H isotopes not only in potential kerogen occurrences on Mars, but also in terrestrial oil and gas resource plays. H isotope extraction and analyses were carried out utilizing a CDS 5000 Pyroprobe connected to a Thermo Trace GC interfaced with a Thermo MAT 253 IRMS. Also, a split of GC-separated products was sent to a DSQ II quadrupole MS to make qualitative and semi-quantitative compositional measurements of these products.

Kerogen samples from five different basins (type II and II-S) were dehydrated (heated to 80°C overnight under vacuum) and analyzed for their H isotope compositions by Pyrolysis-GC-MS-TC-IRMS. This technique takes pyrolysis products separated via GC and reacts them in a high temperature conversion furnace (1450°C), which quantitatively forms H₂. Samples ranging from ~0.5 to 1.0mg in size, were pyrolyzed at 800°C for 30s. and separated on a Poraplot Q GC column.

H isotope data from all kerogen samples typically show enrichment in D from low to high molecular weight. H₂O average $\delta D = -215.2\text{‰}$ (V-SMOW), ranging from -271.8‰ for the Marcellus Shale to -51.9‰ for a Polish shale. Higher molecular weight compounds like toluene (C₇H₈) have an average δD of -89.7‰, ranging from -156.0‰ for the Barnett Shale to -50.0‰ for the Monterey Shale. We interpret these data as representative of potential H isotope exchange between hydrocarbons and sediment pore water during basin formation. Since hydrocarbon H isotopes readily exchange with water, these data may provide some useful information on gas-water or oil-water interaction in resource plays, and further as a possible indicator of paleo-environmental conditions. Alternatively, our data may be an indication of H isotope exchange with water and/or acid during the kerogen isolation process. Either of these interpretations will prove useful when deciphering H isotope data derived from kerogen analyses. Understanding the role that these H-bearing compounds play in terrestrial shale paleo-environmental reconstruction may also prove useful as analogs for understanding the interactions of water and potential kerogen/organic compounds on the planet Mars.