

Validation of the Two-Layer Model for Correcting Clear Sky Reflectance Near Clouds

Guoyong Wen^{1,2}, Alexander Marshak¹, Frank Evans⁴, Tamás Várnai³
¹NASA/Goddard Space Flight Center, ²Morgan State University, ³JCET/UMBC, ⁴University of Colorado



Executive Summary

A two-layer model was developed in our earlier studies to estimate the clear sky reflectance enhancement near clouds. This simple model accounts for the radiative interaction between boundary layer clouds and molecular layer above, the major contribution to the reflectance enhancement near clouds for short wavelengths. We use LES/SHDOM simulated 3D radiation fields to validate the two-layer model for reflectance enhancement at 0.47 μm. We find:

- The simple model captures the viewing angle dependence of the reflectance enhancement near cloud, suggesting the physics of this model is correct.
- The magnitude of the 2-layer modeled enhancement agree reasonably well with the “truth” with some expected underestimation.

We further extend our model to include cloud-surface interaction using the Poisson model for broken clouds. We found that including cloud-surface interaction improves the correction, though it can introduced some over corrections for large cloud albedo, large cloud optical depth, large cloud fraction, large cloud aspect ratio. This over correction can be reduced by excluding scenes (10 km x 10km) with large cloud fraction for which the Poisson model is not designed for. Further research is underway to account for the contribution of cloud-aerosol radiative interaction to the enhancement.

Two-Layer Model



$$\rho = \Delta R = \frac{\alpha_c(\lambda)T_m(\tau(\lambda), \Omega_0)}{1 - \alpha_c(\lambda)R_{m,diff}(\tau(\lambda))} [T_{m,diff}(\tau(\lambda), \Omega) - T_{m,beam}(\tau(\lambda), \Omega)]$$

Where α_c is cloud albedo, Ω_0 and Ω are the direction of the Sun and the direction of the viewing direction of the satellite.


Expression of transmittance and reflectance for molecular layer with optical depth of $\tau(\lambda)$.

Name	Input Flux (source)	Expression
$T_m(\tau, \Omega_0)$	Flux Transmittance F_0 (Beam flux)	$F(\tau = \tau_0) / F_0 \cos(\Omega_0)$
$T_{m,diff}(\tau, \Omega)$	Radiance Transmittance F_0 (Isotropic flux)	$\pi F(\tau = \tau_0) / F_0$
$R_{m,diff}(\tau_0)$	Reflectance F_0 (Isotropic flux)	$F(\tau = 0) / F_0$
$T_{m,beam}(\tau, \Omega)$	Beam Transmittance I_0 (Beam radiance)	$\exp(-\tau_0 / \cos(\Omega))$


SHDOM Simulations

- 26 fields (cloud liquid water content and relative humidity) were simulated using the UCLA large eddy simulation (LES) model
- Combined 26 LES cumulus fields with 40 GEOS-5 aerosol profiles to make 80 cloud/aerosol scene (20 km x 20 km x 15 km).
- Radiances were simulated using SHDOM at 500 m MODIS resolution with 23 viewing direction appropriate for MODIS Aqua.
- MOD04 cloud masking procedure is applied to select “good” pixels. There are 100188 “good” pixels in all 80 cloud scenes, and number of “good” 10 km x 10 km boxes is 3154.


Compare 2-Layer Model and SHDOM Truth



Compare Statistics Between 2-Layer Model and SHDOM Truth




Poisson Stochastic Model for Broken Cloud



Reflectance Near Clouds

With and Without Cloud-Surface



View metadata, citation and similar papers at core.ac.uk
provided by NASA Technical Reports Server
brought to you by CORE