

Geosci. Model Dev. Discuss., 6, 2517–2549, 2013
www.geosci-model-dev-discuss.net/6/2517/2013/
doi:10.5194/gmdd-6-2517-2013
© Author(s) 2013. CC Attribution 3.0 License.



This discussion paper is/has been under review for the journal Geoscientific Model Development (GMD). Please refer to the corresponding final paper in GMD if available.

Downscaling a global climate model to simulate climate change impacts on US regional and urban air quality

M. Trail¹, A. P. Tsimpidi¹, P. Liu¹, K. Tsigaridis^{2,3}, Y. Hu¹, A. Nenes^{4,5}, and A. G. Russell¹

¹School of Civil & Environmental Engineering, Georgia Institute of Technology, Atlanta, GA 30332, USA
²Center for Climate Systems Research, Columbia University, New York, NY 10025, USA
³NASA Goddard Institute for Space Studies, New York, NY 10025, USA
⁴School of Earth & Atmospheric Sciences, Georgia Institute of Technology, Atlanta, GA 30332, USA
⁵School of Chemical and Biomolecular Engineering, Georgia Inst. Technology, Atlanta, GA 30332, USA

Received: 25 February 2013 – Accepted: 3 April 2013 – Published: 15 April 2013

Correspondence to: M. Trail (mcus2rail@gmail.com)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper

GMDD
6, 2517–2549, 2013

Downscaling a global climate model
model

M. Trail et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Abstract

Climate change can exacerbate future regional air pollution events by making conditions more favorable to form high levels of ozone. In this study, we use spectral nudging with WRF to downscale NASA earth system GISS modelE2 results during the years 5 2006 to 2010 and 2048 to 2052 over the continental United States in order to compare the resulting meteorological fields from the air quality perspective during the four seasons of five-year historic and future climatological periods. GISS results are used as initial and boundary conditions by the WRF RCM to produce hourly meteorological fields. The downscaling technique and choice of physics parameterizations used are 10 evaluated by comparing them with in situ observations. This study investigates changes of similar regional climate conditions down to a 12 km by 12 km resolution, as well as the effect of evolving climate conditions on the air quality at major US cities. The high resolution simulations produce somewhat different results than the coarse resolution simulations in some regions. Also, through the analysis of the meteorological variables 15 that most strongly influence air quality, we find consistent changes in regional climate that would enhance ozone levels in four regions of the US during fall (Western US, Texas, Northeastern, and Southeastern US), one region during summer (Texas), and one region where changes potentially would lead to better air quality during spring (northeast). We also find that daily peak temperatures tend to increase in most major 20 cities in the US which would increase the risk of health problems associated with heat stress. Future work will address a more comprehensive assessment of emissions and chemistry involved in the formation and removal of air pollutants.

1 Introduction

25 Changes in climate, emissions, population, technologies, and land-use can impact air quality in a variety of ways. Studies suggest that climate change can exacerbate future regional air pollution events (e.g. (Liao et al., 2006; Mickley et al., 2004; Stevenson et

GMDD

6, 2517–2549, 2013

Downscaling
a global climate
model

M. Trail et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



(CO) and black carbon (BC) tracers to simulate the impact of climate change on air quality in the US. They found that increased severity of future pollutant episodes in the northeast and midwest during the summer was due to a decrease in frequency of surface cyclone tracking in southern Canada.

5 Dawson et al. (2008) developed a global-regional climate-air pollution modeling system (GRE-CAPS) by coupling GISS II GCM, MM5 regional meteorological model, and the PMCAMx regional CTM. This system was intended to enable studies of the effects of changes in climate, intercontinental transport, and emissions on regional and urban air quality. Leung and Gustafson (2005) applied a similar approach to assess the potential effects of climate change in the United States. They developed meteorological fields by downscaling the NASA GISS GCM simulations using an MM5-based RCM (Grell, 1994). Their analyses were based on changes in surface air temperature and downward solar radiation, precipitation frequency, stagnation events, and ventilation. They defined a time to be stagnant when for four consecutive days the following criteria are met: (a) the 10 m wind speed is less than 4 ms^{-1} (b) the 500 mb wind speed is less than 13 ms^{-1} at 07:00 LST, and (c) the total rainfall is less than 0.001 cm for the 4 day period (Korshover and Angell, 1982). They also compared the daily average number of unvented hours, which are hours when the product of the mean wind speed within the boundary layer and the boundary layer height is less than $6000 \text{ m}^2 \text{ s}^{-1}$ (Pielke et al., 1991). In Tagaris et al. (2007), meteorological inputs to the Community Multi-scale Air Quality (CMAQ) Chemistry/Transport Model (CTM) were developed using the meteorological fields of Leung and Gustafson (2005) to investigate the potential impacts of global climate change and emissions on regional air quality using CMAQ (Byun and Schere, 2006). Similarly, Nolte et al. (2008) investigated the impact of climate change on future air quality in the United States by dynamically downscaling outputs from the GISS GCM with the MM5 RCM and predicted an increase in O_3 over Texas and large portions of the southeast using CMAQ model.

25 These studies have illustrated the value of using the regional downscaling approach in order to better understand the impact of climate change on regional air quality. With

GMDD

6, 2517–2549, 2013

Downscaling a global climate model

M. Trail et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the strong dependence on localized flow patterns, air quality models benefit from the higher-resolution wind, temperature, precipitation, and boundary layer structures produced by a RCM (Leung and Gustafson, 2005). Weaver, et al. (2009) stresses that the science of coupling global climate and regional air quality models is still at a young state and that there are particular questions as to which climate metrics and statistics are most relevant to air quality and how sensitive simulation results are to downscaling methodologies.

In our previous recent work (Liu et al., 2012) we examined the performance of two nudging techniques, grid and spectral nudging, by downscaling NCEP/NCAR data using the Weather Research and Forecasting (WRF) Model and identified benefits of spectral nudging at producing small scale features while preserving the large scale forcings. Following these findings, in this study, we use spectral nudging to downscale the NASA earth system GISS modelE2 results during the years 2006 to 2010 and 2048 to 2052 over the continental United States (CONUS) in order to compare the resulting meteorological fields from the air quality perspective during the four seasons of five year historic and future climatological periods. GISS results are used as initial and boundary conditions by the WRF RCM to produce hourly meteorological fields. The downscaling technique and choice of physics parameterizations used are evaluated by comparing them with in situ observations. This study investigates changes of similar regional climate conditions down to a 12 km by 12 km resolution, as well as the effect of evolving climate conditions on the air quality at major US cities.

2 Approach

In this study a regional climate model is used to downscale a global climate model to develop meteorological fields for the present and future. Each component of the modeling system is described below along with the downscaling and evaluation methods used.

GMDD

6, 2517–2549, 2013

Downscaling a global climate model

M. Trail et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2.1 Model descriptions

Global model

Lateral boundary and initial conditions for the regional forecast modeling are taken from the GISS ModelE2. The model has a horizontal resolution of $2^\circ \times 2.5^\circ$ latitude by longitude. The vertical discretization has 40 layers and follows a sigma coordinate up to 150 hPa, with constant pressure layers between 150 and 0.1 hPa. The surface is split into four types: open water (including lakes, rivers and oceans), ice-covered water (including lake ice and sea ice), ground (including bare soil and vegetated regions) and glaciers.

Simulations are carried out for the calendar years 2006–2010 and 2048–2052, driven by future atmospheric conditions over the 21st century and follow the scenario development process for IPCC AR5. The specific scenario used for this study is the “Representative Concentration Pathway” (RCP) 4.5 (Lamarque et al., 2011; Moss et al., 2010), that is a scenario of decadal global emissions of greenhouse gases, short-lived species, and land-use-land-cover which produces an anthropogenic radiative forcing at 4.5 W m^{-2} (approximately 650 ppm CO_2 -equivalent) in the year 2100. The detailed characteristics of this scenario are enumerated in Moss et al. (2010). The atmosphere/terrestrial biosphere-only version of the GISS modelE2 was driven by sea-ice and sea-surface temperature conditions calculated by the coupled earth system model version that is submitted to the CMIP5 archive. The model spinup time was 3 years, starting either from 2003 or 2045, which is sufficient for the dynamic equilibration of the model’s climate and chemically active tracers. SST and sea-ice boundary conditions vary both seasonally and interannually, GHG concentrations change annually, and emissions change annually by linearly interpolating the decadal CMIP5 emission datasets. 6-hourly instantaneous outputs of physical and chemical parameters were produced for regional downscaling by WRF (Sect. 2.2).

GMDD

6, 2517–2549, 2013

Downscaling a global climate model

M. Trail et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Regional model

The regional climate model used is the non-hydrostatic Weather Research and Forecasting (WRF) Model (Skamarock and Klemp, 2008) version 3.4. The simulation domain covers the CONUS and portions of southern Canada and northern Mexico and is centered at 40° N and 97° W with dimensions of 164 × 138 horizontal grids cells with a grid-spacing of 36 km. It contains 35 vertical levels, with the top pressure of 50 hPa. The configuration of physics schemes is as follows: the long-wave Rapid Radiative Transfer Model (RRTM) (Mlawer et al., 1997) and Dudhia scheme (Dudhia, 1989) are used for longwave and shortwave radiation respectively; the Yonsei University (YSU) (Hong et al., 2006) scheme is used for the planetary boundary layer; the Noah scheme (Ek et al., 2003) is used for land surface model (LSM); a revised version of Kain-Fritsch scheme (Kain, 1993) is used to represent the effects of both deep and shallow cumulus clouds; Lin et al. (1983) is chosen for cloud microphysics.

2.2 Dynamical downscale of global results

The GISS ModelE2 fields include temperature, relative humidity, horizontal wind velocities, soil temperature and moisture at different soil depths, sea surface temperature, surface pressure, ice fraction and snow water equivalent. The WRF Preprocessing System (WPS), which reads in this global data and interpolates it to the WRF grid points, does not process GISS data directly. Therefore, an interface program was developed to link the GISS output with WPS. 3-D variables, such as wind and temperature, are interpolated to 21 fixed pressure levels; the lowest level of these 3-D variables and surface level properties were vertically interpolated to produce 2 m temperature, 2 m humidity, and 10 m wind fields. The soil-related variables were also interpolated to the depths defined from the LSM.

Global model results are used as initial and boundary conditions for the regional climate simulations. Spectral nudging with a wave number of 3 in both zonal and meridional directions is used, i.e. all waves with wave numbers greater than 3 are filtered (Liu

GMDD

6, 2517–2549, 2013

Downscaling
a global climate
model

M. Trail et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



et al., 2012); no nudging is conducted for shorter waves to provide similarity with the large scale GCM simulation but allow small scale features to freely develop (Liu et al., 2012). Spectral nudging is applied to temperature, horizontal winds, and geopotential height. Only horizontal winds are nudged at all vertical levels, while no nudging is conducted for other variables within the planetary boundary layer (PBL). The nudging coefficient for all variables was set to $3 \times 10^{-4} \text{ s}^{-1}$ (Stauffer and Seaman, 1990). During the simulation, nudging is conducted every 6 h, consistent with the frequency of the GISS data.

3 Model application and evaluation

WRF is applied here using a nested grid approach. The modeling domain uses a Lambert Conformal Projection centered at 40° N, 97° W with true latitudes of 33° N and 45° N. The outer domain uses a 36 km horizontal grid-spacing that covers the entire continental US as well as portions of Canada and Mexico (5940 × 5004 km). Two innermost domains cover 984 × 1020 km, and, 948 × 948 km regions with 12 km horizontal grid-spacing and focusing on the northeast and southeast US respectively (Fig. 1). The periods modeled are 2006 through 2010 (historic) and 2048 to 2052 (future). The simulated coarse-grid hourly meteorology is used as initial and boundary conditions for the finer grids.

Observations are used to evaluate the ability of GISS-WRF to reproduce the long-term yearly climatic means, and the meteorological fields that strongly impact air quality. The model performance is evaluated by using statistical measures. This is a common analysis that is proposed by Emery and Tai Emery (2001) and has been adopted by the meteorological modeling community. Statistics such as mean bias (MB), mean absolute gross error (IMAGE_Team) and root mean square error (RMSE) are calculated:

GMDD

6, 2517–2549, 2013

Downscaling
a global climate
model

M. Trail et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



$$\text{MAGE} = \frac{1}{N} \sum_{i=1}^N |P_i - O_i| \quad (1)$$

$$\text{MB} = \frac{1}{N} \sum_{i=1}^N (P_i - O_i) \quad (2)$$

$$\text{RMSE} = \sqrt{\frac{1}{N} \sum_{i=1}^N (P_i - O_i)^2} \quad (3)$$

5 where P_i is the predicted value of the tested parameter (i.e. temperature), O_i is the corresponding observed value, and N is the total number of the predictions used for the comparison. MAGE gives an estimation of the overall discrepancy between predictions and observations, while MB is sensitive to systematic errors. The root mean square error (RMSE) incorporates both the variance of the prediction and its bias.

10 Additional details for the above evaluation metrics can be found in Yu et al. (2006). The observations used for the statistical analysis are TDL (Techniques Development Laboratory) data from the Research Data Archive (RDA) <http://dss.ucar.edu> in dataset number ds472.0), which is maintained by the Computational and Information Systems Laboratory (CISL) at the National Center for Atmospheric Research (NCAR). These

15 are hourly surface observations for wind speed, wind direction, and temperature during the four seasons over a five year period (2006–2010). In the statistical analysis, the continental US domain has been divided in 4 sub regions, the western (W), the midwest (MW), the south (S) and the northeast (NE) USA (Fig. 1).

4 Results

20 Table 1 summarizes the comparison of the GISS-WRF modeling system predictions for wind speed and direction against TDL hourly surface observations during the four

GMDD

6, 2517–2549, 2013

Downscaling a global climate model

M. Trail et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



as O₃ and secondary PM, but warming could lead to decreased emissions of PM from heating processes such as wood burning (e.g. from wood stoves). The GCM simulations predict a similar warming pattern during the winter and spring, but only up to 3 degrees K (Fig. S1). During the summer months, Texas and northeastern Canada
5 experience a warming of 2 and 4 degrees respectively, although the GCM predicts up to one degree more warming over western Texas and the p value associated with the downscaled temperature changes over western Texas is between 0.05 and 0.10. An average warming of 3 degrees occurs over the Midwest (p value < 0.05) and a warming of around 2 degrees also occurs over most of Texas and Eastern US (the p value is
10 between 0.05 and 0.15 which is not significant) during the fall. The eastern US states, on the other hand, are cooler during the winter and spring months with the southeastern states and Texas cooling up to just less than 2 degrees, however the cooling here is not statistically significant (p value greater than 0.05).

The smaller, more highly resolved, 12 km domain over the northeast simulates similar magnitudes of temperature change to the 36 km domain. The root mean square difference of the future temperature change between the 36 km and 12 km domains is very small (less than 0.004 K); indicating the similarity between the two simulations. The standard deviations of the simulated temperature changes in the northeast for the 12 km (standard deviation of 0.25 K²) and 36 km (0.24 K²) domains show that the fine
20 resolution simulation introduces slightly more variability than the coarse resolution domain, especially during the winter (0.49 K² for the 36 km and 0.52 K² for 12 km) and spring (0.27 K² for 36 km and 0.29 K² for 12 km). The northeast sees cooling of less than 1 degree during the spring and warming of up to 2 degrees during the summer (Fig. 2b). During fall, large warming between 2–3 degrees is simulated over much of
25 New York State. Over the southeastern 12 km domain, similar warming occurs as the 36 km domain, ranging between 1 and 3 degrees during the summer and fall with the greatest warming occurring during the fall over North Carolina and Tennessee (Fig. 2c).

GMDD

6, 2517–2549, 2013

Downscaling a global climate model

M. Trail et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



4.2 Insolation and precipitation

A change in downward solar radiation at the surface, or insolation, is an indicator of changes in cloudiness. For this reason, spatial distributions of the change in insolation at the surface are similar in structure to average daily precipitation, but not identical.

5 Spatial distributions of surface temperature and insolation changes have similar structures in some cases. Weaver et al. (2009) explain that these meteorological conditions can have either competing or reinforcing effects on air quality. When temperature and insolation change in the same direction, O₃ concentrations tend to change in the same corresponding direction, whereas temperature and insolation varying in opposite directions correspond with mixed changes in O₃.

10 Decreases in daily mean precipitation are found over the Pacific coast where some regions receive 2 mm less rain per day (or 30 % less rain), on average, and some decreases were simulated over the southeastern region (Fig. 3a). Reduced rain along the Pacific coast occurs mostly during the winter, as a major portion of Western US sees greater than a 2 mm per day decrease. Correspondingly, insolation over the Pacific coast increases during the winter by up to 15 W m⁻² (Fig. 4a). The southeast experiences a similar magnitude of drying, but mainly during the fall. Both the 36 and the 12 km simulations over the southeast predicted greater than 2 mm less rain per day during the fall (Figs. 3c and 4c), which is also consistent with insolation changes in the region (increase of up to 10 W m⁻²). Interestingly, the high-resolution simulation predicts that the southeast receives up to 2 mm per day more rain during the summer, which is not apparent in the 36 km domain. The 36 and 12 km resolution simulations over the northeast on the other hand, predict more rain over most of Vermont, New Hampshire and Maine during the summer, while most of Connecticut and New York receive less rain (Fig. 3b). The precipitation trend in the northeast reverses during the fall when Connecticut and New York receive more rain and the states farther north are dryer. There is a decrease in insolation of 5–15 W m⁻² during the spring and fall and an increase during the fall of 10–20 W m⁻² over the northeast (Fig. 4b). Correspondingly,

GMDD

6, 2517–2549, 2013

Downscaling a global climate model

M. Trail et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the temperature only decreases slightly during spring, while it increases during summer. However, contrary to the insolation trend, the temperature in the northeast sees large increases during fall.

4.3 Stagnation events

5 Stagnation events occur when wind speeds are low and little precipitation occurs over an extended period. Since transport and deposition of pollutants is decreased during a stagnation period these events promote poor air quality. During the winter and spring months, the spatial distribution of the number of stagnation days per season does not change significantly over the US (Fig. 5a). Over southern Texas, the number of
10 stagnation days during fall increases in some small areas by 10 to 15 days per season, which correlates with the increase in temperature in the region. Large portions of this region already see over 30 days of stagnation per season. Stagnation decreases over Texas during the summer. Over most of the southeast stagnation days also decrease by up to 10 days per season corresponding to the increase in precipitation (Fig. 5c) which
15 is large compared to the average number of stagnation days during the summer of the historic simulation (between 15 and 30). While the 36 km domain shows little change in stagnation in the northeast, the high resolution simulation shows stagnation increases of up to 5 days per season during the summer over parts of the northeast which is large compared to the average 5 to 10 stagnation days per season that occur in this region
20 (Fig. 5b). During fall, a large increase (over 15 days) in the number of stagnation days is found along the Gulf coast and the California coast. Along the coast of California during fall, the increase in stagnation days leads to increased concentrations of pollutants, reinforcing the negative impact that increased temperature and insolation have on air quality in the area. Similarly, the decreased precipitation along the Gulf coast may
25 reinforce higher concentrations of pollutants due to increased stagnation in the area.

GMDD

6, 2517–2549, 2013

Downscaling a global climate model

M. Trail et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Los Angeles, Philadelphia, Phoenix, and Seattle. Further, land use changes may exacerbate meteorological changes in cities. Philadelphia and Phoenix are chosen here, rather than more populated cities such as New York and Houston, because future studies are planned to address the impact of land use changes on regional climate in those cities. Extremes in meteorological variables are important because there are adverse health effects associated with short term exposure to poor air quality. Air pollution is highly variable in time and temperature extremes are also associated with adverse health outcomes (McMichael et al., 2006). Cumulative distribution function (CDF) plots show the percentage of hourly temperature and daily precipitation that exceed a given value in major US cities for each simulation year (Fig. 7a–e). The hourly temperatures within any given percentile range tend to shift a few degrees warmer in the future, except for the lower 20th percentile range in Atlanta and Philadelphia and the upper 90th percentile range in Seattle. Most warming in Seattle occurs at the lower 75 percentile range, where high O₃ concentrations are not likely, which reflects the increase in temperature mentioned earlier during the winter in the Pacific Northwest. This can also decrease emissions related to domestic heating, including PM from wood burning. The cumulative distribution of the maximum daily temperature in Seattle follows a similar trend, as the hourly CDF in the lower range and temperature are nearly the same in the 60 to 90 percentile range (Fig. 8a–e). The upper 95th percentile in daily maximum temperatures in Seattle are actually around a degree cooler in the future, decreasing the chance of high ozone during hot days. Los Angeles and Phoenix have similar hourly temperature and maximum temperature CDF structures and shifts from present to future. These cities are warmed by 1 to 3 degrees regardless of the percentile range. Lin et al. (2001) have developed estimates of the probability that the maximum daily 8 h average O₃ will exceed 80 ppb given the maximum daily temperature in a given region (including Los Angeles, the southeast, and the northeast). Given that the upper 95th percentile in daily maximum temperatures in Phoenix increases from around 308 K to 312 K, the probability that O₃ will exceed 80 ppb on these days increases. Similarly, a shift in the upper 95th percentile in daily maximum temperatures in Los Angeles,

GMDD

6, 2517–2549, 2013

Downscaling a global climate model

M. Trail et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



from around 305 K to 308 K, also increases the probability of high O₃. The coolest days in Atlanta and Philadelphia are similar to, if not cooler than the future simulations, reflecting the cooling that occurs during the winter and spring in the surrounding regions. While very little, if any, warming occurs in the lower 50th percentile of daily maximum temperature, the upper 95th percentile maximum temperatures in Philadelphia and Atlanta increase from around 302 K to 303 K and 304 K to about 307 K respectively, with a corresponding increase in probability of high O₃ concentrations on those hot days. The 12 km simulations produce similar, but not identical, cumulative distribution of total hourly temperatures and maximum temperatures in Atlanta and Philadelphia (Fig. 9). The 12 km domains show a shift in daily maximum temperature at the upper 95th percentile of 303 to 305 K and 304 to 305 K in Philadelphia and Atlanta respectively, implying a higher probability of high ozone on those days than would be derived from the coarse resolution. The hottest days in Philadelphia are simulated by the future 12 km domain where temperatures reach 310 K. Maximum temperatures and high O₃ probability in New York exhibit similar changes to that of Philadelphia. Temperature distributions in Chicago were also analyzed, however the only difference from present to future is a shift in the upper 95th percentile from 300 to 301 K while the distribution below the 90th percentile does not change.

Rain can improve air quality, and although the seasonal mean precipitation has already been examined, it is important to also understand how the frequency and distribution of rainfall can shift over time. Rainfall frequency shifts are especially important in cities, where frequent light rains will clean the air more than infrequent heavy rains. The most notable change in daily precipitation distribution occurs in Los Angeles, where the driest year is simulated in the future and the wettest year is simulated in the historic simulations (Fig. 10). Seattle does not seem to receive more or less rain in the future but does have a more consistent distribution of daily rainfall from year to year in the future. The precipitation distributions of the remaining cities do not appear to be affected. Precipitation distributions produced from the 12 km simulations over Atlanta and Philadelphia are similar to the distribution produced by the coarse simulation (Fig. S2).

GMDD

6, 2517–2549, 2013

Downscaling a global climate model

M. Trail et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



5 Conclusions

The high resolution simulations produce somewhat different results than the coarse resolution simulations in some regions. Also, through the analysis of the meteorological variables that most strongly influence air quality, we find consistent changes in regional climate that would enhance ozone levels in four regions of the US during fall, one region during summer, and one region where changes potentially would lead to better air quality during spring.

During summer and fall, all air quality indicators, with the exception of insolation and precipitation, suggest an increase in air pollutant concentrations, including increased production of secondary PM and O₃, in most of Texas. Consistent with Leung and Gustafson (Leung and Gustafson, 2005), when comparing 2048–2052 to 2006–2010, we find warmer temperatures (2–3 K), less ventilation (1–2 h per day) and more stagnation (10–15 days per season) during summer in Texas and the same for fall, though slightly less intense.

The West Coast is marked by warmer temperatures (ranging from 1–4 K), slightly less rainfall (less than 1 mm per day), and more stagnation (10–15 days) during fall, while there is no significant change in ventilation and insolation. Similarly during fall, the southeast shows little change in ventilation and stagnation but is warmer (2–3 K), dryer (up to 2 mm per day in some areas) and with slightly higher insolation (5–10 W m⁻²). Also, the shift in the distribution of maximum daily temperatures in Atlanta increases the probability of high O₃ concentrations for days when the maximum temperature is in the 95th percentile.

The northeast is also warmer during fall (2–3 K) and less ventilated (1–2 h per day) but increased rainfall (up to 2 mm per day) and decreased insolation (~ 5 W m⁻²) compete for better air quality. Regardless, as in Atlanta, the shift in the distribution of maximum daily temperature in New York and Philadelphia increases the probability of high O₃ concentrations (0.06–0.12) for days when the maximum temperature is in the 95th percentile. During spring, however, increased rain (1–2 mm per day), more ventilation

GMDD

6, 2517–2549, 2013

**Downscaling
a global climate
model**

M. Trail et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



(~ 1 h per day) and decreased temperatures (~ 1 K) could promote better air quality in the northeast.

While climate conditions strongly impact air quality, emissions and chemistry also play a vital and complex role in the formation and removal of atmospheric pollutants.

5 A more comprehensive assessment of emissions and chemistry will be addressed in the future.

Supplementary material related to this article is available online at:

<http://www.geosci-model-dev-discuss.net/6/2517/2013/gmdd-6-2517-2013-supplement.pdf>.

10 References

Byun, D. and Schere, K.: Review of the governing equations, computational algorithms, and other components of the models-3 Community Multiscale Air Quality (CMAQ) modeling system, *Appl. Mech. Rev.*, 59, 51–77, 2006.

15 Dawson, J. P., Racherla, P. N., Lynn, B. H., Adams, P. J., and Pandis, S. N.: Simulating present-day and future air quality as climate changes: model evaluation, *Atmos. Environ.*, 42, 4551–4566, doi:10.1016/j.atmosenv.2008.01.058, 2008.

Dudhia, J.: Numerical study of convection observed during the winter monsoon experiment using a mesoscale two-dimensional model, *J. Atmos. Sci.*, 46, 3077–3107, 1989.

20 Ek, M. B., Mitchell, K. E., Lin, Y., Rogers, E., Grunmann, P., Koren, V., Gayno, G., and Tarp-ley, J. D.: Implementation of Noah land surface model advances in the National Centers for Environmental Prediction operational mesoscale Eta model, *J. Geophys. Res.-Atmos.*, 108, 8851, doi:10.1029/2002jd003296, 2003.

Emery, C. and Tai, E.: Meteorological modeling and performance evaluation of the 13–20 September 1999 ozone episode, prepared for the Texas near non-attainment areas through the Alamo Area Council of Governments, by ENVIRON International Corp. Navato, CA., 2001.

25 EPA, US: The potential effects of global climate change on the United States, EPA, EPA-230-05-89-050, 457, 1989.

GMDD

6, 2517–2549, 2013

Downscaling a global climate model

M. Trail et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Giorgi, F.: Regional climate modeling: status and perspectives, *J. Phys.* IV, 139, 101–118, doi:10.1051/jp4:2006139008, 2006.
- Grell, G. A., Dudhia, J., and Stauffer, D. R. A.: A Description of the Fifth Generation Penn State/NCAR Mesoscale Model (MM5), National Center for Atmospheric Research, Boulder, CO, 1994.
- 5 Hong, S. Y., Noh, Y., and Dudhia, J.: A new vertical diffusion package with an explicit treatment of entrainment processes, *Mon. Weather Rev.*, 134, 2318–2341, 2006.
- IMAGE_Team: The IMAGE 2.2 implementation of the SRES scenarios. A comprehensive analysis of emissions, climate change and impacts in the 21st century, National Institute for Public Health and the Environment, Bilthoven, the Netherlands, 2001.
- 10 Jacob, D. J. and Winner, D. A.: Effect of climate change on air quality, *Atmos. Environ.*, 43, 51–63, doi:10.1016/j.atmosenv.2008.09.051, 2009.
- Kain, J. S. and Fritsch, J. M.: Convective parameterization models: the Kain–Fritsch scheme, cumulus convection in numerical models, *Am. Meteorol. Soc.*, 46, 165–170, 1993.
- 15 Korshover, J. and Angell, J. K.: A review of air-stagnation cases in the Eastern-United-States during 1981 – annual summary, *Mon. Wea. Rev.*, 110, 1515–1518, doi:10.1175/1520-0493(1982)110<1515:aroasc>2.0.co;2, 1982.
- Lamarque, J. F., Kyle, G. P., Meinshausen, M., Riahi, K., Smith, S. J., van Vuuren, D. P., Conley, A. J., and Vitt, F.: Global and regional evolution of short-lived radiatively-active gases and aerosols in the representative concentration pathways, *Climatic Change*, 109, 191–212, doi:10.1007/s10584-011-0155-0, 2011.
- 20 Leung, L. R. and Gustafson, W. I.: Potential regional climate change and implications to US air quality, *Geophys. Res. Lett.*, 32, 4, L16711, doi:10.1029/2005gl022911, 2005.
- Liao, H., Chen, W. T., and Seinfeld, J. H.: Role of climate change in global predictions of future tropospheric ozone and aerosols, *J. Geophys. Res.-Atmos.*, 111, D12304, doi:10.1029/2005jd006852, 2006.
- 25 Liao, K. J., Tagaris, E., Manomaiphiboon, K., Napelenok, S. L., Woo, J. H., He, S., Amar, P., and Russell, A. G.: Sensitivities of ozone and fine particulate matter formation to emissions under the impact of potential future climate change, *Environ. Sci. Tech.*, 41, 8355–8361, 2007.
- 30 Lin, Y. L., Farley, R. D., and Orville, H. D.: Bulk parameterization of the snow field in a cloud model, *J. Clim. Appl. Meteorol.*, 22, 1065–1092, doi:10.1175/1520-0450(1983)022<1065:bpotsf>2.0.co;2, 1983.

GMDD

6, 2517–2549, 2013

**Downscaling
a global climate
model**

M. Trail et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Liu, P., Tsimpidi, A. P., Hu, Y., Stone, B., Russell, A. G., and Nenes, A.: Differences between downscaling with spectral and grid nudging using WRF, *Atmos. Chem. Phys.*, 12, 3601–3610, doi:10.5194/acp-12-3601-2012, 2012.
- McMichael, A. J., Woodruff, R. E., and Hales, S.: Climate change and human health: present and future risks, *Lancet*, 367, 859–869, doi:10.1016/s0140-6736(06)68079-3, 2006.
- Mickley, L. J., Jacob, D. J., Field, B. D., and Rind, D.: Effects of future climate change on regional air pollution episodes in the United States, *Geophys. Res. Lett.*, 31, L24103, doi:10.1029/2004gl021216, 2004.
- Mlawer, E. J., Taubman, S. J., Brown, P. D., Iacono, M. J., and Clough, S. A.: Radiative transfer for inhomogeneous atmospheres: RRTM, a validated correlated-k model for the longwave, *J. Geophys. Res.-Atmos.*, 102, 16663–16682, doi:10.1029/97jd00237, 1997.
- Moss, R. H., Edmonds, J. A., Hibbard, K. A., Manning, M. R., Rose, S. K., van Vuuren, D. P., Carter, T. R., Emori, S., Kainuma, M., Kram, T., Meehl, G. A., Mitchell, J. F. B., Nakicenovic, N., Riahi, K., Smith, S. J., Stouffer, R. J., Thomson, A. M., Weyant, J. P., and Wilbanks, T. J.: The next generation of scenarios for climate change research and assessment, *Nature*, 463, 747–756, doi:10.1038/nature08823, 2010.
- Nolte, C. G., Gilliland, A. B., Hogrefe, C., and Mickley, L. J.: Linking global to regional models to assess future climate impacts on surface ozone levels in the United States, *J. Geophys. Res.-Atmos.*, 113, D14307, doi:10.1029/2007JD008497, 2008.
- Pielke, R. A., Stocker, R. A., Arritt, R. W., and McNider, R. T.: A procedure to estimate worst-case air-quality in complex terrain, *Environ. Int.*, 17, 559–574, doi:10.1016/0160-4120(91)90168-p, 1991.
- Rind, D. and Lerner, J.: Use of on-line tracers as a diagnostic tool in general circulation model development, 1. Horizontal and vertical transport in the troposphere, *J. Geophys. Res.-Atmos.*, 101, 12667–12683, doi:10.1029/96jd00551, 1996.
- Rind, D., Lerner, J., Shah, K., and Suozzo, R.: Use of on-line tracers as a diagnostic tool in general circulation model development 2. Transport between the troposphere and stratosphere, *J. Geophys. Res.-Atmos.*, 104, 9151–9167, doi:10.1029/1999jd900006, 1999.
- Skamarock, W. C. and Klemp, J. B.: A time-split nonhydrostatic atmospheric model for weather research and forecasting applications, *J. Computat. Phys.*, 227, 3465–3485, doi:10.1016/j.jcp.2007.01.037, 2008.

GMDD

6, 2517–2549, 2013

**Downscaling
a global climate
model**

M. Trail et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 1. GISS-WRF modeling system performance for wind speed and direction against TDL observations for US regions and seasons.

Wind Speed Direction	West				Midwest				South				Northeast				Total Domain			
	Winter	Fall	Spring	Summer	Winter	Fall	Spring	Summer	Winter	Fall	Spring	Summer	Winter	Fall	Spring	Summer	Winter	Fall	Spring	Summer
Spd Mean OBS (ms ⁻¹)	4.3	3.9	4.6	4.0	4.7	4.6	4.8	3.7	4.4	3.8	4.4	3.6	4.4	4.4	4.1	3.4	4.4	4.3	4.5	3.6
Spd Mean PRD (ms ⁻¹)	5.1	4.0	4.5	3.6	4.3	4.5	5.0	3.5	3.7	4.2	4.2	3.5	4.1	3.6	5.0	3.5	4.4	4.4	4.7	3.5
Spd Bias (ms ⁻¹)	0.8	0.1	-0.1	-0.4	-0.4	-0.2	0.3	-0.1	-0.7	0.2	-0.2	-0.1	-0.3	-0.8	1.0	0.2	-0.0	0.1	0.2	-0.1
Spd Error (ms ⁻¹)	3.0	2.3	2.5	1.9	2.3	2.6	2.6	1.4	2.1	2.2	2.2	1.7	2.5	2.5	2.4	1.7	2.6	2.5	2.5	1.9
Spd RMSE (ms ⁻¹)	3.9	3.0	3.3	2.5	3.0	3.2	3.2	2.3	2.7	2.8	2.8	2.2	3.4	3.4	3.2	2.4	3.4	3.3	3.2	2.4
Dir Mean OBS (degrees)	213	256	264	248	231	242	179	219	223	183	168	145	209	218	192	216	248	265	192	214
Dir Mean PRD (degrees)	213	221	245	262	220	212	229	167	219	111	156	172	240	246	198	193	254	238	233	200
Dir Bias (degrees)	5.6	-4.7	-1.0	3.8	0.3	-2.3	-5.2	1.9	-4.5	-1.8	4.6	6.5	8.2	0.9	1.1	4.1	2.2	-2.3	-1.7	2.2
Dir Error (degrees)	78	77	73	72	91	86	86	81	89	80	81	68	83	80	85	216	86	82	82	76

GMDD

6, 2517–2549, 2013

Downscaling a global climate model

M. Trail et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 2. GISS-WRF modeling system performance for temperature against TDL observations for US regions and seasons.

Temperature	West				Midwest				South				Northeast				Total Domain			
	Winter	Fall	Spring	Summer	Winter	Fall	Spring	Summer	Winter	Fall	Spring	Summer	Winter	Fall	Spring	Summer	Winter	Fall	Spring	Summer
Mean OBS (K)	278	285	286	298	274	279	286	297	285	290	294	301	276	283	285	297	276	282	287	297
Mean PRD (K)	273	286	281	293	264	286	284	296	280	292	292	301	267	285	284	295	269	285	284	295
Bias (K)	-4.7	1.1	-4.7	-4.2	-10	6.8	-1.9	-1.4	-4.5	2.4	-2.7	-0.2	-9.7	2.2	-1.4	-2.0	-7.5	2.9	-2.7	-1.9
Error (K)	6.7	4.7	5.9	5.4	11	7.5	5.8	3.7	7.4	5.3	4.7	2.7	11	5.2	4.7	3.2	9.2	5.7	5.4	3.8
RMSE (K)	9.1	5.9	7.8	6.7	14	8.8	7.1	4.6	9.6	6.6	6.0	3.5	13	6.4	6.0	4.1	12	7.1	6.8	5.0

GMDD

6, 2517–2549, 2013

Downscaling a global climate model

M. Trail et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 1. Modelling domains with horizontal grid-spacing resolutions of 36 km and 12 km northeast (NE) and southeast (SE).

GMDD

6, 2517–2549, 2013

Downscaling a global climate model

M. Trail et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 2. Predicted average yearly and seasonal 2 m atmospheric temperature change (future minus historic) for **(a)** the 36 × 36 km resolution modeling domain, **(b)** the 12 × 12 km resolution sub-domain over northeast and **(c)** the 12 × 12 km resolution sub-domain over southeast.

Downscaling
a global climate
model

M. Trail et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 3. (a) Predicted average yearly and seasonal precipitation (mm day^{-1}) change (future minus historic) for (a) the 36×36 km resolution modeling domain, (b) the 12×12 km resolution sub-domain over northeast and (c) the 12×12 km resolution sub-domain over southeast.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 4. Predicted average yearly and seasonal downward short wave radiative flux at the surface ($W m^{-2}$) change (future minus historic) for (a) the 36×36 km resolution modeling domain, (b) the 12×12 km resolution sub-domain over northeast and (c) the 12×12 km resolution sub-domain over southeast.




Fig. 5. Predicted total seasonal change in the number of stagnation days (days per season) (future minus historic) for **(a)** the 36×36 km resolution modeling domain, **(b)** the 12×12 km resolution sub-domain over northeast and **(c)** the 12×12 km resolution sub-domain over south-east.




Fig. 6. Predicted total seasonal change in average unvented hours (hours per day) (future minus historic) for (a) the 36 × 36 km resolution modeling domain, (b) the 12 × 12 km resolution sub-domain over northeast and (c) the 12 × 12 km resolution sub-domain over southeast.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Downscaling
a global climate
model**

M. Trail et al.




Fig. 7. Empirically determined cumulative distribution of 36 km historic (2006–2010) and future (2048–2050) hourly temperatures at major US cities: **(a)** Atlanta, **(b)** Los Angeles, **(c)** Philadelphia, **(d)** Phoenix and **(e)** Seattle.



Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Downscaling
a global climate
model

M. Trail et al.




Fig. 8. Empirically determined cumulative distribution of 36 km historic (2006–2010) and future (2048–2050) maximum daily 1 h average temperature at major US cities: (a) Atlanta, (b) Los Angeles, (c) Philadelphia, (d) Phoenix and (e) Seattle.




Fig. 9. Empirically determined cumulative distribution of 12 km historic (2006–2010) and future (2048–2050) hourly temperature **(a)** Atlanta, **(b)** Philadelphia and maximum daily 1 h average temperature at **(c)** Atlanta, **(d)** Philadelphia.




Fig. 10. Cumulative distribution of 36 km historic (dark) and future (light) daily precipitation at major US cities: **(a)** Atlanta, **(b)** Los Angeles, **(c)** Philadelphia, **(d)** Phoenix and **(e)** Seattle.

**Downscaling
a global climate
model**

M. Trail et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

