

# P162 Expansion of the Real-Time SPoRT-Land Information System for NOAA/National Weather Service Situational Awareness and Local Modeling Applications



Jonathan L. Case; ENSCO, Inc./NASA Short-term Prediction Research and Transition (SPoRT) Center; Huntsville, AL, and  
Kristopher D. White; NOAA/NWS Huntsville Weather Forecast Office, Huntsville, AL  
American Meteorological Society 94<sup>th</sup> annual meeting; Atlanta, GA; 26<sup>th</sup> Conf. on Weather Analysis and Forecasting / 22<sup>nd</sup> Conf. on Numerical Weather Prediction  
Poster Session 2; 4 February 2014

## Background

SPoRT runs the NASA Land Information System (LIS) in real-time to support local modeling and diagnostics at NOAA/National Weather Service (NWS) weather forecast offices (WFOs)

- Current domain covers the Southeastern half of the Continental U.S. (CONUS) due to limitations in the Stage IV precipitation grids driving the Noah land surface model (LSM) integration in LIS
- This past year, SPoRT added a new real-time run over a full CONUS domain
  - Enables expansion of LIS applications to NOAA/NWS partners outside Southern Region
  - Sets stage for future soil moisture data assimilation activities


## Poster objectives

- Provide summary of real-time LIS activities at SPoRT
- Compare/contrast real-time LIS over SE U.S. with full CONUS-LIS
- Map out future direction of LIS applications

## Current Applications of SPoRT-LIS

Initializing LSM fields in local modeling applications (i.e., WRF model)

- Supported option in the NWS SDO Science Training and Resource Center's Environmental Modeling System (EMS; <http://strc.comet.ucar.edu/software/newrems>)
- LIS GRIB output files uploaded to ftp server in real-time
- EMS users over SE U.S. can initialize with LIS LSM fields in place of coarser-resolution, large-scale model fields



## Situational Awareness

- Drought Monitoring
- Assessing flood potential
- LIS data ingested and displayed in AWIPS II at NWS Huntsville, AL
- Refer to training examples below

## Vegetation Stress during Growing Season

- University of Alabama - Huntsville acquires SPoRT-LIS and MODIS vegetation products
- Manages in-house crop-stress model over SE U.S.
- Distributes reports to end-users


## Comparison Between SE U.S. and CONUS SPoRT-LIS Configurations

Table 1. Summary of configuration details for the real-time SPoRT-LIS runs over the Southeastern U.S. domain and new CONUS domain.


Configuration detail	Common to both domains	Southeastern U.S. domain	CONUS domain
Land surface model	Noah version 3.2		
Horizontal grid spacing	0.03 degrees		
Grid dimensions		1064 x 672	2200 x 934
Atmospheric Forcing	NLDAS-2, GDAS, GFS SW/LW Rad, sfc P, 2-m T, 2-m q, 10-m wind	Stage IV hourly Precipitation <sup>1</sup>	NSSL's Multi-Radar Multi-Sensor (MRMS) hourly gauge-adjusted radar QPE <sup>2</sup>
Static soil database	STATSGO		
Static land use database	MODIS/IGBP		
Green Vegetation Fraction	Daily SPoRT-MODIS <sup>3</sup>		
Cold start initialization	1 June 2010		
History restart interval	6 hours		

<sup>1</sup>Lin and Mitchell (2005), Liu et al. (2006); <sup>2</sup>Zhang et al. (2011); Zhang et al. (2014, this meeting, 28<sup>th</sup> Conf. Hydrology); <sup>3</sup>Case et al. (2014)


## Current SE CONUS domain with Stage IV



## New full CONUS domain with MRMS



## Sample Results / Comparison between SE U.S. LIS and CONUS LIS



Summer 2013 had distinct dipole pattern in total column soil moisture change: SW U.S. moistening; Midwest drying (above) Better/worse USDM classification over SW/Midwest (below)

- MRMS-driven LIS soil moisture tends to be drier, esp. in Ohio Valley / Appalachians
- Drier tendency accentuated by late Autumn as precipitation increases across this region

## Issues Documented with MRMS Precipitation Dataset

Comparison between sample soil moist and WSR-88D radar coverage (left)

- Note similar patterns in soil moisture to radar coverage patterns in Western U.S.
- Moist soils within radar coverage; dry soils outside radar coverage gaps
- MRMS product very dependent on available input radar data

Beam blockage due to terrain / physical impediments

- Not just a concern in Rocky Mountains
- Columbus, MS radar: Rapidly-growing trees have blocked beam over time at certain azimuths
- Pattern particularly discernable in integrated soil moisture fields
- LIS is a good tool to identify problems in QPE datasets through long-term integrations

Edge of radar networks and non-overlapping radars

- Especially a problem in S. Canada / N. Mexico
- Recommend that end-users do not utilize CONUS-LIS output in these problem regions
- Better blending of precipitation forcing and/or soil data assimilation needed to improve spatial continuity

Periodic drop-outs of regional tiles (fixed Oct 2013)

- Numerous days in late Summer / Autumn 2013
- Instead of precip assigned as missing, entire regional tile was filled with "0" values
- Led to artificial gradients in soil moisture in portions of domain when active precipitation occurs at tile boundary

Points with no input data assigned "0" (fixed Feb 2013)

- Instead of a missing data flag, grid points with no input radar or gauge data were set to "0" (a legitimate value)
- Led to artificially dry soil, esp. in Canada and Mexico

(top-left) Examples of Beam Blockage problems in MRMS Columbus, MS and Columbia radars, manifested in locally drier moisture in distinct radial patterns


(bottom-left & above) Example regional tile "drop-out" caused by artificial soil moisture gradient edge of missing MRMS tiles

## Modeling System and Capabilities

### NASA Land Information System (LIS)

- High-performance land surface modeling & data assimilation framework
- Can run a variety of LSMs (Noah, SLM, Catchment, etc.)
- Supports several static databases for land use and soil classification
- Able to run up to global domains at 1-km grid spacing
- Land surface data assimilation
  - Ensemble Kalman Filter (EnKF) or Direct Insertion (DI)
  - Soil moisture, skin temperature, snow fraction/depth/SWE
  - Kumar et al. 2009 (J. Hydromet); soil moisture); Liu et al. 2013 (Adv. Water Res.; snow)
- Optimization and Uncertainty Estimation (Santanello et al. 2013, J. Hydromet)

## LIS modes of operation



## Development of LIS Training Module for Situational Awareness Applications

### SPoRT-LIS for Drought Monitoring

Examining LIS volumetric soil moisture during a dry-down period can help quantify the amount of soil-moisture degradation on sub-county scales

- Daily temperatures reached 95-100 F in Muscle Shoals and Russellville during this period with very little rainfall
- Highlighted regions had varying degrees of drying on local scales, which can help refine decisions about drought classification

### SPoRT-LIS for Assessing Flood Potential

Early March 2011 Event: Prevailing wet soils

High antecedent soil moisture + Moderate-heavy precipitation → Moderate river flooding and numerous flooding reports

September 2011 Event: Tropical Storm Lee with dry soil

Low antecedent soil moisture + Heavy precipitation → Isolated minor flooding

Contrasting antecedent soil moisture conditions likely played a strong role in the different outcomes

Local, subjective analysis of several events suggests typical moderate-heavy synoptic rainfall events over deep-layer relative soil moisture values exceeding 60% will lead to more substantial moderate or heavier flooding events

## Poster References

Case, J. L., F. J. LaFontaine, J. R. Bell, G. J. Jedlovec, S. V. Kumar, and C. D. Peters-Lidard, 2014: A real-time MODIS vegetation product for land surface and numerical weather prediction models. *Trans. Geost. Remote Sens.*, **52**(3), 1772-1786.

Kumar, S. V., R. H. Reichle, R. D. Koster, W. T. Crow, and C. D. Peters-Lidard, 2009: Role of subsurface physics in the assimilation of surface soil moisture observations. *J. Hydrometeorol.*, **10**, 1534-1547.

Liu, Y., and K. E. Mitchell, 2006: The NCEP Stage IV hourly precipitation analysis: Development and applications. *Preprints, 15th Conf. on Hydrology, San Diego, CA, Amer. Meteor. Soc.*, 1.2. [Available online at <http://www.confex.com/ams/pdfpapers/33847.pdf>].

\_\_\_\_\_, E. Rogers, and G. J. DiMego, 2005: Using hourly and daily precipitation analyses to improve model budget. *Hydrometeorol.*, **6**, 124-138.

\_\_\_\_\_, and K. E. Mitchell, 2006: The NCEP Stage IV hourly precipitation analysis: Development and applications. *Preprints, 15th Conf. on Hydrology, San Diego, CA, Amer. Meteor. Soc.*, 1.2. [Available online at <http://www.confex.com/ams/pdfpapers/33847.pdf>].

Liu, Y., C. D. Peters-Lidard, S. Kumar, J. L. Foster, M. Shaw, Y. Tian, and G. M. Fall, 2013: Assimilating satellite-based snow depth and snow cover products for improving snow predictions in Alaska. *Adv. Water Res.*, **54**, 208-227.

Santanello Jr., J. A., S. V. Kumar, C. D. Peters-Lidard, K. Harrison, S. Zhou, 2013: Impact of land model calibration on coupled land-atmosphere prediction. *J. Hydrometeorol.*, **14**, 1373-1400.

Vargas, M., Z. Jiang, J. Ju, and A. A. Castar, 2013: EVI based green vegetation fraction derived from Suomi NPP-VIIRS. *Preprints, 15th Symp. Future Operational Env. Sat. Systems, Austin, TX, Amer. Meteor. Soc.*, 16B8.

Zhang, J., and Coauthors, 2011: National Mosaic and multi-sensor QPE (NMQ) system: Description, results, and future plans. *Bull. Amer. Meteor. Soc.*, **92**, 1321-1338.

\_\_\_\_\_, 2014: Initial operating capabilities of quantitative precipitation estimation in the Multi-Radar Multi-Sensor system. *28<sup>th</sup> Conf. on Hydrology, Atlanta, GA, Amer. Meteor. Soc.*, 5.3 [Available online at <http://www.confex.com/ams/pdfpapers/34467.pdf>].

## Future Direction

- Upgrade to LISv7 and utilize LIS Validation Toolkit
    - Validate LIS against soil moisture observations and field campaign
    - Identify regional errors/biases, and causes of those errors
  - Assimilate satellite-based soil moisture from SMOS and SSM/T
  - Incorporate NESDIS global real-time VIIRS green vegetation fraction (Vargas et al. 2013; *9th Future Op. Env. Sat. Conf.*)
- 
- SMOS DA in LIS: P53 (Blankenship et al.), 28<sup>th</sup> Hydro., 4 Feb 2014, this meeting