

HYDROTHERMAL PROCESSES AND MOBILE ELEMENT TRANSPORT IN MARTIAN IMPACT CRATERS — EVIDENCE FROM TERRESTRIAL ANALOGUE CRATERS. H. E. Newsom¹, M. J. Nelson¹, C. K. Shearer¹, and B. O. Dressler², ¹Univ. of New Mexico, Institute of Meteoritics, Dept. of Earth & Planetary Sci., Albuquerque, NM 87131, USA newsom@unm.edu, ²185 Romfield Circuit, Thornhill, Ontario, L3T 3H7, Canada.

Introduction: Hydrothermal alteration and chemical transport involving impact craters probably occurred on Mars throughout its history. [1]. Our studies of alteration products and mobile element transport in ejecta blanket and drill core samples from impact craters show that these processes may have contributed to the surface composition of Mars [2]. Recent work on the Chicxulub Yaxcopoil-1 drill core has provided important information on the relative mobility of many elements that may be relevant to Mars [3].

The Chicxulub impact structure in the Yucatan Peninsula of Mexico and offshore in the Gulf of Mexico is one of the largest impact craters identified on the Earth, has a diameter of 180-200 km, and is associated with the mass extinctions at the K/T boundary. The Yax-1 hole was drilled in 2001 and 2002 on the Yaxcopoil hacienda near Merida on the Yucatan Peninsula. Yax-1 is located just outside of the transient cavity, which explains some of the unusual characteristics of the core stratigraphy. No typical impact melt sheet was encountered in the hole and most of the Yax-1 impactites are breccias. In particular, the impact melt and breccias are only 100 m thick which is surprising taking into account the considerably thicker breccia accumulations towards the center of the structure and farther outside the transient crater encountered by other drill holes.

SIMS studies of mobile element transport: Clays in thin sections from crater drill cores were imaged and analyzed by microprobe for major elements. Trace elements Li, B, Be, and Ba were measured with the Cameca IMS 4f ion probe, using primary O^+ ions, a 10 kV potential; a primary beam current of 10 nA and spot diameter of 8 to 10 μ m. Numerous standards were analyzed to verify the ability of the SIMS to analyze water-bearing mineral phases, and to show the absence of matrix effects.

Results: The altered groundmass found between clasts of target rocks contain materials with a desiccation texture characteristic of clays. The chemistry of the clays in thin section and preliminary XRD data on separated clays are consistent with smectite at all depths [4, 5]. The chemistry of these alteration materials range from that of an average montmorillonite composition in the uppermost units (from 800.68m to about 836m), to that of a magnesium rich saponite in the lower units (846.7m to 861.72m).

The clay analytical data show interesting trends as a function of depth in the drill core (e.g. **Figs.**

1-3). The elements Li, B, and Be, FeO generally increase upwards in the samples, with $Li > B > Be > FeO$ while the Ba concentration in the clays decreases upwards. The abundance of Al_2O_3 in the clay material is also significantly higher in the suevite than in lower units, even though Al_2O_3 is not usually considered a fluid mobile component. Li in Unit 2 is enriched by factors of 1.1 to 3.5 relative to the lower units. There is a strong correlation between the abundances of Li, Be, and B (e.g. **Fig. 1**). The correlation coefficients among these elements are $Li-B = 0.89$, $B-Be = 0.71$, $Li-Be = 0.50$.




Fig. 1. Boron and lithium abundances in Chicxulub Yaxcopoil-1 matrix clays.

The concentrations of mobile elements determined by SIMS in clays in the Yaxcopoil drill core can be compared with bulk analyses of the impact breccias by Tuchscherer et al., [5]. Although, the types of samples from the core that were analyzed are not identical, a consistent behavior can be seen in the data for a number of mobile elements. For example, in the bulk analyses of the drill core, Tuchscherer et al., [5], documented upward enrichments in the concentration of the trace elements in the relative order $Cs > Au > Rb = Zn$ in stratigraphic units 1 to 4. The greatest upward increase in this data set is for Cs, which increases a factor of up to 3.7 relative to the lower units, similar to the SIMS data for Li.

Mobile element deposition in sediments above the impactites: Evidence for hydrothermal transport has also been presented by Rowe et al. [6], who studied mobile element concentrations in the sediments overlying the impact breccias. Rowe et al., [6] found that samples of Tertiary biomicrites from depths of 794.01 to 777.02 m have higher concentrations of Mn, Fe, P, Ti, and Al relative samples from higher up in the stratigraphic section. They attribute the observed en-

