

NASA/CN-97-

206475

1-1011  
7N-51-12  
OCIT  
12592-1

Final Report for  
"Investigation of Isotopic and Geochemical Evidence for an  
Active Planktonic Biota in the Precambrian"

Grant No. NAGW-5184

July 1, 1996 - June 30, 1997

Submitted by

Lee R. Kump  
Earth System Science Center/Dept. of Geosciences  
Pennsylvania State University  
210 Deike Building  
University Park, PA 16802  
814-863-1274  
[kump@geosc.psu.edu](mailto:kump@geosc.psu.edu)

## Final Report, Contract No. NAGW-5184

**Title of Cooperative Agreement:** "Investigation of Isotopic and Geochemical Evidence for an Active Planktonic Biota in the Precambrian"

**Type of Report:** Final

**Period Covered by Report:** July 1, 1996 - June 30, 1997

**Principal Investigator:** Lee R. Kump  
Earth System Science Center  
The Pennsylvania State University  
248 Deike Building  
University Park, PA 16802

### Report Narrative:

The funded research was motivated by the earlier study of Burdett *et al.* (1990), who collected carbon and oxygen isotopic data from Paleoproterozoic rocks of the Northwest Territories from deep- and shallow-water facies of the Rocknest Platform. Their results (Fig. 1a) displayed a possible decrease in  $\delta^{13}\text{C}$  with depth when arranged by increasing distance from the paleoshore. The most  $^{13}\text{C}$ -depleted samples were seafloor cements and fans from the underlying siliciclastic Odjick Formation, and slope carbonates of the Rocknest platform..




Figure 1. Preliminary data on the carbon and oxygen isotopic composition of carbonates from the Rocknest (**a**; Burdett *et al.*, 1990) and Pethei (**b**; this study) Platforms.

Intrigued by these data, and in collaboration with John Grotzinger and Michael Pope (MIT), Roberta Hotinski (Penn State Graduate student) and I collected samples in the summer of 1996 from the Pethei platform (with funding from this project). The isotopic data we generated represent seafloor and/or early diagenetic cements from shallow (filled diamonds) and deep water (filled circles) facies (Fig. 1b). The  $\delta^{18}\text{O}$  values of our shallow water carbonates agree with previous estimates of a lighter ocean composition of about -7 to -10 ‰ in the Precambrian (Burdett *et al.*, 1990; Veizer *et al.*, 1992). The data do, however, show a wide spread in  $\delta^{18}\text{O}$  which might be the result of differential diagenesis of the deep-water facies. Deeper water samples likely have experienced a decrease rather than an increase in  $\delta^{13}\text{C}$  due to diagenesis, so we tentatively conclude that the water column in the vicinity of the Pethei platform had a carbon isotopic gradient of at most 1‰.

We presented a paper at the CSPG-SEPM Joint Convention (Hotinski and Kump, 1997) in which we argued that this small gradient is consistent with the idea that atmospheric  $p\text{CO}_2$  was elevated in the Paleoproterozoic, compensating for a less luminous sun. As a result, oceanic inorganic carbon contents were high, and the ability of organisms to stratify the isotopic composition of the oceans was diminished. Large inferred gradients in the Neoproterozoic indicate low atmospheric  $p\text{CO}_2$ , which is consistent with the occurrence of glaciation at the time.

**References:**

- Burdett, J.W., and Grotzinger, J.P., and Arthur, M.A., 1990. Did major changes in the stable-isotope composition of Proterozoic seawater occur? *Geology*, 18:227-230.
- Hotinski, R. and Kump, L.R. A geochemical investigation of the Pethei Group, Northwest Territories, Canada: Implications for Paleoproterozoic ocean chemistry. *Programs with Abstracts, CSPG-SEPM Joint Convention*, 1997, p. 132.
- Veizer, J., Plumb, K.A., Clayton, R.N., Hinton, R.W., and Grotzinger, J.P., 1992. Geochemistry of Precambrian carbonates: V. Late Paleoproterozoic seawater. *Geochim. Cosmochim. Acta*, 56: 2487-2501.