

NASA-CR-204739

Reprinted from

11-51-21
94112-
03247

HEARING RESEARCH

Hearing Research 91 (1995) 63-71

Damage and recovery of hair cells in fish canal (but not superficial) neuromasts after gentamicin exposure

Jiakun Song ^{*}, Hong Young Yan ¹, Arthur N. Popper

Department of Zoology, University of Maryland, College Park, MD 20742-4415, USA

Received 13 May 1995; revised 5 July 1995; accepted 24 July 1995



ELSEVIER

AIMS AND SCOPE

The aim of the journal is to provide a forum for papers concerned with basic auditory mechanisms. Emphasis is on experimental studies, but theoretical papers will also be considered.

The editor of the journal is prepared to accept original research papers in the form of full-length papers, short communications, letters to the Editor, and reviews. Papers submitted should deal with auditory neurophysiology, ultrastructure, psychoacoustics and behavioural studies of hearing in animals, and models of auditory functions. Papers on comparative aspects of hearing in animals and man, and on effects of drugs and environmental contaminants on hearing function will also be considered. Clinical papers will not be accepted unless they contribute to the understanding of normal hearing functions.

TYPES OF PAPERS

1. Full-length research papers should include a summary not exceeding 200 words and 3-6 key words and should be divided into sections.
2. Short communications should provide a brief but complete account of a particular piece of work, and will be limited to 4 printed pages, including 1 figure. A summary of not more than 50 words should be included.
3. Letters to the Editor should be comments and clarifications on articles that have been published in *Hearing Research*, and should be limited to 4 printed pages.
4. Review articles should give a survey, evaluation and critical interpretation of recent research data and concepts in the fields covered by the journal.
5. The inclusion of announcements, book reviews, etc., is at the discretion of the Editor-in-Chief and the Publisher, and subject to space availability.

EDITOR-IN-CHIEF: Professor Aage R. Møller, Pittsburgh, PA, USA.

EDITORIAL SECRETARY: Clet Szczępaniak

EDITORIAL BOARD

L. Aitkin, Clayton, Australia
R. Altschuler, Ann Arbor, MI, USA
R.C. Bilger, Champaign, IL, USA
R.P. Bobbin, New Orleans, LA, USA
B.A. Bohne, St. Louis, MO, USA
W.E. Brownell, Houston, TX, USA
D.A. Cotanche, Boston, MA, USA
D.G. Drescher, Detroit, MI, USA
J.J. Eggermont, Calgary, Canada
D.M. Green, Gainesville, FL, USA
R. Klinke, Frankfurt/M., FRG
M.C. Liberman, Boston, MA, USA
D.J. Lim, Rockville, MD, USA
B. Lonsbury-Martin, Miami, FL, USA
N. Manuyama, Suzuka, Japan
K. Morest, Farmington, CT, USA
A.L. Nuttall, Ann Arbor, MI, USA
A.N. Popper, College Park, MD, USA

R. Pujol, Montpellier, France
W.S. Rhode, Madison, WI, USA
E.W. Rubel, Seattle, WA, USA
A.F. Ryan, San Diego, CA, USA
A.N. Salt, St. Louis, MO, USA
R.J. Salvi, Buffalo, NY, USA
P.A. Santi, Minneapolis, MN, USA
J. Santos-Sacchi, New Haven, CT, USA
J.C. Saunders, Philadelphia, PA, USA
J. Schacht, Ann Arbor, MI, USA
N. Slepceky, Syracuse, NY, USA
G. Smoorenburg, Soesterberg, the Netherlands
K. Steel, Nottingham, UK
J. Syka, Prague, Czech Republic
T. Takasaka, Sendai, Japan
I.E. Thalmann, St. Louis, MO, USA
T.F. Weiss, Cambridge, MA, USA
J. Zwislocki, Syracuse, NY, USA

SUBSCRIPTION INFORMATION

Subscription for 1995: Volumes 82-92 (11 volumes). Subscriptions are accepted on a prepaid basis only, unless different terms have been previously agreed upon. For this journal a personal subscription rate and a special rate for ARO members are available. Prices are available upon request. Personal subscriptions are issued on the assumption that these are for personal use only and will not replace an institutional subscription. Journals are sent by surface delivery to all countries, except the following countries where SAL air delivery (Surface Airlifted Mail) is ensured: USA, Canada, Japan, Australia, New Zealand, PR China, India, Israel, South Africa, Malaysia, Singapore, South Korea, Taiwan, Pakistan, Hong Kong, Brazil, Argentina, Mexico and Thailand. Airmail rates for other countries are available upon request.

Subscription orders can only be entered by calendar year (Jan-Dec) and should be sent to: Elsevier Science B.V., Journals Department, P.O. Box 211, 1000 AE Amsterdam, the Netherlands (Tel: 31 20 4853642; Fax: 31 20 4853598), or to your usual subscription agent. Claims for missing issues should be made within three months of publication, otherwise they cannot be honoured free of charge.

In the United States and Canada: All questions arising after acceptance of a manuscript by the editor, especially those relating to proofs, publication and reprints, should be directed to the publishers, Elsevier Science B.V., P.O. Box 1527, 1000 BM Amsterdam, the Netherlands. For further information concerning this or any other Elsevier Science journal, contact: Elsevier Science, Inc., Attn: Journal Information Center, 655 Avenue of the Americas, New York, NY 10010, USA. Tel. (212) 633-3750; Telefax (212) 633-3990; Telex 420-643 AEP UI.

Claims and editorial enquiries concerning accepted manuscripts should be addressed to: Hearing Research, Elsevier Science B.V., P.O. Box 1527, 1000 BM Amsterdam, the Netherlands (Telex 18582 ESPA NL; Fax No. (20)485-3266) stating editor's code and manuscript number.

Advertising information: Advertising orders and enquiries may be sent to: International: Elsevier Science, Advertising Department, The Boulevard, Langford Lane, Kidlington, Oxford, OX5 1GB, UK. Tel.: (+44) (0) 1865-843565; Fax: (+44) (0) 1865-843952.

USA and Canada: Weston Media Associates, Dan Lipner, P.O. Box 1110, Greens Farms, CT 06436-1110, USA. Tel.: (203) 261-2500; Fax: (203) 261-0101.

Japan: Elsevier Science Japan, Ms. Noriko Kodama, 20-12 Yushima, 3 chome, Bunkyo-ku, Tokyo 113, Japan. Tel.: (+81) 3-3836-0810; Fax: (+81) 3-3839-4344.

US mailing notice - *Hearing Research* (ISSN 0378-5955) is published monthly by Elsevier Science (Molenwerf 1, P.O. Box 211, 1000 AE Amsterdam, the Netherlands). Annual subscription price in the USA US\$ 1990 (valid in North, Central and South America only) (subject to change) air speed delivery. Second class postage rate is paid at Jamaica, NY 11431.

USA POSTMASTERS: Send address changes to *Hearing Research* Publication Expediting, Inc. 200 Meacham Avenue, Elmont, NY 11003.

Airfreight and mailing in the USA by Publications Expediting.

© 1995 Elsevier Science B.V. All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, without the prior written permission of the publisher. Copyright and Permissions Department Elsevier Science B.V., P.O. Box 1527, 1000 BM, the Netherlands.

Submission of a paper to this journal entails the author's irrevocable and exclusive authorization of the publisher to collect any sums or consideration for copying or reproduction payable by third parties (as mentioned in article 17 paragraph 2 of the Dutch Copyright Act of 1912 and in the Royal Decree of June 20, 1974 (S.351) pursuant to article 16b of the Dutch Copyright Act of 1912) and/or to act in or out of Court in connection therewith.

Abstract

Recent evidence (oscar) indicates similar hetero-receptors, the hair cell destruction and lagena of similar to type

Keywords: Hair

1. Introduction

Early studies whereas am hair cells (hair cells) hair cell type to be a structure in the

Although cal and str anamniotes ter, 1981; evidence of and amniot two distinct teleost fish. 1990a, b; Y

* Corresponding author.
Present address:
University of

0378-5955/95/\$10.00
SSDI 0378-5955(95)00000-0

Damage and recovery of hair cells in fish canal (but not superficial) neuromasts after gentamicin exposure

Jiakun Song^{*}, Hong Young Yan¹, Arthur N. Popper

Department of Zoology, University of Maryland, College Park, MD 20742-4415, USA

Received 13 May 1995; revised 5 July 1995; accepted 24 July 1995

Abstract

Recent evidence demonstrating the presence of two types of sensory hair cell in the ear of a teleost fish (*Astronotus ocellatus*, the oscar) indicates that hair cell heterogeneity may exist not only in amniotic vertebrates but also in anamniotes. Here we report that a similar heterogeneity between hair cell types may also occur in the other mechanosensory organ of the oscar, the lateral line. We exposed oscars to the aminoglycoside (ototoxic) antibiotic gentamicin sulfate and found damaged sensory hair cells in one class of the lateral line receptors, the canal neuromasts, but not in the other class, the superficial neuromasts. This effect was not due to the canal environment. Moreover, new ciliary bundles on hair cells of the canal neuromasts were found after, and during, gentamicin exposure. The pattern of hair cell destruction and recovery in canal neuromasts is similar to that of type I-like hair cells found in the striolar region of the utricle and lagena of the oscar after gentamicin treatment. These results suggest that the hair cells in the canal and superficial neuromasts may be similar to type I-like and type II hair cells, respectively, in the fish ear.

Keywords: Hair cell; Lateral line; Canal neuromast; Superficial neuromast; Fish; Gentamicin

1. Introduction

Early studies of the vertebrate inner ear suggested that whereas amniote vestibular endorgans have two types of hair cells (types I and II), anamniotes have only type II hair cells (Wersäll, 1956). As a consequence, diversity of hair cell types in amniotes has generally been considered to be a structurally and functionally advanced characteristic in the evolution of the vertebrate ear.



Although there have been some reports of morphological and structural diversity of hair cells in the ear of anamniotes (Hama, 1969; Corwin, 1977; Popper and Hoxter, 1981; Wegner, 1982; Jensen, 1984), there was no clear evidence of a possible homology between the anamniote and amniote hair cell types until the recent discovery of two distinct types of sensory hair cells in the ear of the teleost fish, *Astronotus ocellatus*, the oscar (Saidel et al., 1990a, b; Yan et al., 1991; Chang et al., 1992). The two

types of hair cell in the oscar ear, one in the striolar region and the other in the extrastriolar region of the utricle and the lagena, may be similar to amniote type I and type II hair cells, respectively. Thus, heterogeneity of sensory hair cells may not be an advanced amniote trait, as previously proposed. In fact, hair cell diversity may extend much earlier in the evolution of the vertebrate ear than the origin of amniotes. It is still not known, however, when such diversity first occurred in the evolution of vertebrate mechanosensory systems.

The mechanosensory organs of vertebrates consist of the lateral line neuromasts and the inner ear. The neuromasts are only found in aquatic anamniotes, whereas the inner ear occurs in all the vertebrates. Ontogenetically, the sensory epithelium of both organs are closely related because the lateral line placodes differentiate from the primary otic placode (Platt, 1896; Landacre and Conger, 1913; Northcutt, 1989). The sensory epithelia in both organs contain sensory hair cells and supporting cells for detecting motion of surrounding water or fluid. Because of these similarities, the lateral line and the inner ear are collectively referred to as the octavolateralis system (Northcutt, 1980; McCormick, 1982; Popper et al., 1992). However, the inner ear has been considered to be struc-

^{*} Corresponding author. Tel.: (301) 405-6901; Fax: (301) 314-9358.

¹ Present address: Thomas H. Morgan School of Biological Sciences, University of Kentucky, Lexington, KY 40506-0225, USA.



turally and (van Berge

Despite the sensory system has been considered (Kroes, 1989; Kroes, 1989; Kroes, 1989), however, the presence of the lateral line system in the oscar is described in the literature. The evolution of the lateral line system is suggested

In the oscar, the lateral line system consists of two types of hair cells: the primary and the secondary. The primary hair cells are located in the trunk region and are characterized by their large size and the presence of a single stereocilium. The secondary hair cells are located in the head region and are characterized by their smaller size and the presence of multiple stereocilia. The lateral line system is innervated by the lateral line nerve, which carries sensory information to the brain.

2. Materials and Methods

Thirty groups of cichlids (Cichlasoma nigrofasciatum) with a total length of 46, 50, and 54 mm were used. The fish were kept in a controlled environment and fed with a commercial fish feed. The experiment was approved by the Animal Care and Use Committee of the University of California, San Diego.

Fig. 1. Peritoneal openings; the trunk is opened) from superficial layer covered by

turally and functionally more complex than the neuromasts (van Bergeijk, 1967).

Despite some morphological variation among species, the sensory hair cells of the lateral line neuromasts had been considered to be of only a single type (Jørgensen, 1989; Kroese and van Netten, 1989). It becomes important, however, to reexamine the hair cell types in the neuromasts in the light of the recent evidence showing the presence of two types of hair cells in the inner ear of the oscar to determine whether hair cell heterogeneity is present in the lateral line system. If heterogeneity is found in the lateral line, it could mean that this arose even earlier in the evolution of the vertebrate mechanosensory organs than suggested by the current evidence from the fish ear.

In the oscar ear, the aminoglycoside antibiotic gentamicin sulfate selectively damages ciliary bundles of hair cells in the striolar regions of the utricle without damaging hair cells in the extrastriolar regions (Yan et al., 1991; Lombarte et al., 1993). The pattern of damage in the striolar region appears similar to that observed for mammalian type I hair cells under various aminoglycoside antibiotic assaults (Wersäll and Hawkins, 1962; Wersäll et al., 1971). Thus, the types of hair cell in the fish striolar and extrastriolar regions of the utricle and lagena may be related to mammalian types I and II hair cells, respectively. The purpose of this study was to determine if the lateral line receptors of the oscar also have two types of hair cells as in the oscar inner ear. We hypothesized that if two types of hair cells occur in the lateral line neuromasts, one group would be more sensitive to gentamicin than the other. Therefore, we exposed the fish to a solution of gentamicin sulfate and examined the response of ciliary bundles of the hair cells in the neuromasts.

2. Materials and methods

Thirty oscars were divided into three experimental groups. One experimental group contained 6 fish (52–58 mm total length (TL)) while the other two groups had 12 fish each (42–48 mm TL). An additional 3 animals (44, 46, and 54 mm TL) were used as controls. The experimental fish were kept in aerated water containing 0.002% gentamicin sulfate. The water was completely changed each day to refresh the gentamicin. All experiments were done according to the policies of the University of Maryland Animal Care and Use Committee.

Fish were killed after being deeply anesthetized with

tricaine methane sulfonate (MS222; Sigma). Two fish each (from the group of large fish) were killed after 4, 8, and 12 days of gentamicin exposure, and 2 fish each (from a group of small fish) were killed after 1, 2, 3, 4, 8, and 12 days exposure. The third experimental group (small fish) received gentamicin treatment until day 4, followed by 4 or 8 additional days survival during which time the water was changed daily but without gentamicin. Two fish each from this group were killed on the same schedule as the other small fish group. Control animals were kept in similarly sized tanks as the experimental animals, and their water was changed daily. The fish were killed after 4 days.

Tissue from experimental and control fish was prepared for scanning electron microscopic (SEM) examination following procedures described previously (Song and Northcutt, 1991a, b). Briefly, fish were immersed in 4% glutaraldehyde in 0.1 M phosphate buffer (pH 7.4) and fixed for a period of time ranging from overnight to 2 days. After decalcification in 0.1% HCl for about 30 min, the fish were washed with distilled water and cut mid-sagittally. The lateral line canals were opened with microneedles. Each half of the fish was cut transversely to separate the head, trunk, and tail. After dehydration in a graded series of ethanol and critical-point drying with liquid CO₂, samples containing lateral line neuromasts were mounted on aluminum stubs, sputter-coated with gold-palladium, and viewed and photographed with an Amray 1820 SEM.

3. Results

3.1. Normal organization

The mechanosensory lateral line receptors in the oscar, as in most fishes, consist of canal and superficial neuromasts that are distributed over the surface of the head and the body in a species-specific pattern (Fig. 1). The morphological differences between the canal (or superficial) neuromasts on the head and the body are minimal. The trunk of the oscar has a dorsorostral and a mediocaudal lateral line. Each of the trunk scales in the lateral line has a single canal neuromast that is covered by a bony canal with an opening on either end of the scale. Only a few trunk canal neuromasts in each fish are not covered by bony canals (Fig. 1A). There are two or three superficial neuromasts found dorsally to each canal neuromast on the trunk of fish in the size range of our experimental groups (42–58 mm TL).

Fig. 1. Peripheral distribution of the lateral line canal and superficial neuromasts in an oscar (56 mm TL). Large dot, nares; small circles, lateral line canal openings; small dots, superficial neuromasts; small oval dots, uncovered canal neuromasts. A: SEM overview of the skin surface (with caudal portion of the trunk lateral lines) on the body. Arrowheads (upper right) point to a pair of superficial neuromasts dorsal to a lateral line scale (canal is surgically opened) from the dorso-rostral section of the trunk lateral line. Arrows point to the uncovered canal neuromast. Bar = 1 mm. B: high-power view of a superficial neuromast (arrowhead in upper left). An adjacent canal neuromast sits in a surgically opened canal (long arrow in center). Both neuromasts are covered by a gelatinous cupula. Bar = 100 μ m. Orientation: head to the right, dorsal to the top.

Although the canal and the superficial neuromasts in the oscar share the same basic organization, they also differ in several ways. In our experimental fish, the canal neuromasts are larger and possess over 120 hair cells, whereas the superficial neuromasts are smaller and each has about 40 hair cells (Fig. 2A,C). The density of hair cells and the length of their kinocilia are greater in superficial neuromasts than in canal neuromasts (compare Fig. 2B,D).

3.2. Gentamicin exposure

After 4 days of gentamicin exposure, superficial neuromasts in experimental animals did not show a loss of the ciliary bundles (Fig. 2C,D). However, canal neuromasts, both covered and not covered by scales, had a substantial loss of ciliary bundles with as little as 1 day of exposure (Fig. 3A,B). Control animals had no loss of ciliary bundles in either canal (Fig. 2A) or superficial neuromasts.

New small ciliary bundles on hair cells were found among the support cell microvilli in the canal neuromasts by the 2nd or 3rd day of treatment in the smaller fish and by the 4th day in the larger fish. These new bundles, however, did not attain the normal hair cell density even after 8 and 12 days of treatment (compare Fig. 4A-C and 2A).




Fig. 2. Low (A) and high (B) magnification SEMs of a canal neuromast from a trunk canal of a control specimen. The bony canal was surgically opened during SEM preparation. Low (C) and high (D) magnification SEMs of a superficial neuromast from the epidermal surface dorsal to the trunk canal neuromasts from an animal continuously exposed to gentamicin sulfate for 4 days. No damage to the ciliary bundles is found. Long arrows point to a kinocilium; arrowhead points to a set of stereocilia. Bar = 5 μ m.

In contrast, there were far more ciliary bundles present in the canal neuromasts of the animals that had 4 days (Fig. 4) or 8 days (Fig. 4E,F) recovery in plain water than in animals that had 8 days (Fig. 4A) or 12 days (Fig. 4B,C) of gentamicin exposure. The area covered by ciliary bundles in the canal neuromasts of the animals exposed to gentamicin for 4 days followed by 8 days of recovery (Fig. 4E,F) approached the pattern found in control animals (compare to Fig. 2A,B). The new hair cell bundles, however, appeared to have longer kinocilia than those in the canal neuromasts of the control animals (compare Figs. 4E,F and 2A,B), a characteristic similar to that of normal superficial neuromast hair cells (Fig. 2C,D).

4. Discussion

The results of our experiments demonstrate that the sensory hair cells in the canal and the superficial neuromasts respond differently to a gentamicin sulfate solution at the concentration used in our study. Moreover, the different responses to gentamicin parallel, respectively, the responses of the types I and II hair cells in the utricle and

lagna of These data types occ organs, the experimen generated even unde

4.1. Class

Neuron depending Coombs e ing the o open groo cial neuro (Fig. 1). the same in a num morpholo Münz, 19 neuomas frequen (Münz, 1 findings, cance of difference rudimenta Even tional, va the epide receptors (Jørgense fore, our gentamic time, the cells be added to canal an lateral li from one While neuoma gobies (canals in It will b these ne the norr though epiderm be base position

4.2. Typ

The p neuoma

ndles present
ad 4 days of
(4) or 8 days
animals that
of gentamicin
in the canal
micin for 4
(4E,F) ap-
(compare to
er, appeared
l neuromasts
and 2A,B), a
ficial neuro-

ate that the
ficial neuro-
fate solution
oreover, the
ectively, the
e utricle and



gically opened
he trunk canal
ows point to a

lagena of the oscar and in amniote vestibular endorgans. These data may suggest that the heterogeneity of hair cell types occurs in both of the vertebrate mechanosensory organs, the inner ear and the lateral line neuromasts. Our experiments also demonstrate that ciliary bundles are regenerated in the canal neuromasts after exposure to, and even under the continuing presence of, gentamicin.

4.1. Classification of lateral line neuromasts

Neuromasts have been classified as canal or superficial, depending on their epidermal position (reviewed by Coombs et al., 1988; Münz, 1989). In most fishes, including the oscar, the canal neuromasts are enclosed within open grooves, bony canals, or scales, whereas the superficial neuromasts are found on the surface of the epidermis (Fig. 1). Although superficial and canal neuromasts exhibit the same basic organization, they differ from one another in a number of ways, including innervation pattern and morphology of hair cell arrangement (Hama, 1965, 1978; Münz, 1989; Song and Northcutt, 1991a, b). The canal neuromasts also appear to be more sensitive to higher frequencies of vibration than are the superficial neuromasts (Münz, 1985; Kroese and van Netten, 1989). Despite these findings, however, our knowledge of the biological significance of the morphological, physiological, and functional differences between superficial and canal neuromasts are rudimentary.

Even though there is morphological, and some functional variation between neuromasts and their position on the epidermis, the sensory hair cells in the lateral line receptors have been considered to be only a single type (Jørgensen, 1989; Kroese and van Netten, 1989). Therefore, our results showing differences in the response to gentamicin by hair cells in the oscar indicate, for the first time, the existence of a biochemical heterogeneity of hair cells between the canal and superficial neuromasts. This, added to the known physiological differences between canal and superficial neuromasts, suggests that the two lateral line receptors may be functionally quite distinct from one another.

While the oscar, as most fishes, has two types of neuromasts, there are a number of fish species, such as the gobies (Song, 1993), that have reduced the lateral line canals in certain regions of the fish but not the neuromasts. It will be interesting and important to determine whether these neuromasts maintain distinct types of hair cells from the normal canal instead of superficial neuromasts even though both neuromasts are located on the surface of epidermis. If so, the classification of the neuromasts will be based on more meaningful criteria other than their position on the epidermis.

4.2. Types of hair cells in the lateral line

The pattern of destruction of ciliary bundles of the canal neuromast by gentamicin sulfate observed in this study is

reminiscent of the damage caused by gentamicin to striolar hair cells of the oscar (Yan et al., 1991; Lombarte et al., 1993) and goldfish ears (Platt and Yan, 1993) and to mammalian vestibular type I hair cells (Wersäll, 1960; Lindeman, 1969). In contrast, the ciliary bundles of the hair cells in the superficial neuromasts were not damaged by the same dosage of gentamicin sulfate treatment. This is similar to the findings for extrastriolar hair cells of the oscar ear (Yan et al., 1991; Lombarte et al., 1993) and mammalian type II hair cells (Wersäll, 1960; Lindeman, 1969).

One question that arose in our study was whether the different response to gentamicin damage in canal versus superficial neuromasts was associated with the environment within the canal. Accumulation of gentamicin in the canal could, theoretically, have produced a much higher concentration of the drug than that presented to superficial neuromasts. To test this, we took advantage of the fact that the oscar, like most teleost fish, has a number of trunk canal lateral line scales that fail to form canals over their neuromasts (Fig. 1). If the differential damage to the canal organs was due to the concentration of drug in the canal environment, we would expect no ciliary bundle damage in




Fig. 3. SEMs of canal neuromasts after 1 day in gentamicin sulfate showing the loss of ciliary bundles on the hair cells. A: Canal neuromast where the canal cover was surgically removed during SEM preparation. B: Canal neuromast that was not normally covered by a bony canal. This is the next neighbor to the neuromast in A. Bar = 10 μm.

the neuromasts of uncovered canals. However, we found that the canal neuromasts on the uncovered scales (Fig. 3A) had damage that was similar to the damage on the neighboring covered canal neuromasts (Fig. 3B) in the same fish.

The other question concerned whether the different response to gentamicin of the hair cells in canal versus superficial neuromasts was related to the state of development of the hair cells, since the hair cells in the superficial neuromasts and the immature hair cells in the canal neuro-

masts share a morphological feature of having a long kinocilia (Fig. 2C-D vs. Fig. 4). It has been known that immature hair cells are less susceptible to aminoglycoside toxicity than are mature hair cells (Richardson and Russell, 1991). Our new observation of the differences in ultrastructure, innervation, and postembryonic growth pattern between the hair cells in the two classes of neuromasts, however, suggests that it is not likely the case (Song et al., 1995).

Although further study of the gentamicin susceptibility




Fig. 4. SEMs of canal neuromasts after 8 (A) and 12 days (B) in gentamicin sulfate. C: Higher magnification view of (B). D: Canal neuromast after 4 days in gentamicin sulfate followed by 4 days of recovery in water with no drug (total 8 days). Low (E) and high (F) magnification view of a canal neuromast after 4 days in gentamicin sulfate and 8 days recovery (total 12 days). Long arrow points to a kinocilium. Short arrow head points to a set of stereocilia. Bar = 10 μ m.

of the sup
exposure
is requir
susceptib
that the
cial neur
chemical
as the tw
ear. Bec
receptors
mal plac
Metcalf
heteroge
organs a
studied.

4.3. Reg

Our r
neuroma
ciliary b
the repl
(Fig. 3A
examine
crosscop
dence th
eration.
romasts.
the can:
cells be
suggest:
and tha
cells. T
examine
gentami
crosscop
to Rege
not pre
or fish
in the c
that ha:
the pre:
1991)
damag:
al., 199
may no
hair ce
that in:
ble for
(Hashi:
our res
know :
by sim
were a
(intran
this ma

giving a long
known that
aminoglycoside
and Russell,
differences in ultra-
growth pattern
neuromasts,
(Song et al.,
susceptibility

of the superficial neuromast hair cells in different length of exposure time and in different dosages of the gentamicin is required in order to understand the mechanism of the susceptibility. The results from this study, at least, suggest that the hair cells of the lateral line canal and the superficial neuromasts in the oscar may share the similar biochemical heterogeneity in response to gentamicin sulfate as the two distinct types of hair cells in the vertebrate inner ear. Because the sensory epithelia of both the lateral line receptors and the inner ear are from closely related epidermal placodes embryonically (Landacre and Conger, 1913; Metcalfe, 1985; Northcutt, 1989), it is possible that the heterogeneity between the two types of hair cells in both organs are homologous. This hypothesis should be further studied.

4.3. Regeneration of lateral line hair cells

Our results not only demonstrate that intact lateral line neuromasts of the oscar are capable of replacing hair cell ciliary bundles after aminoglycoside treatment but also that the replacement can occur in the presence of gentamicin (Fig. 3A-C) that is refreshed daily. Still, since we did not examine the hair cells with transmission electron microscopy in this study, we do not have histological evidence that there was actual hair cell destruction and regeneration. Our SEM examination of gentamicin-treated neuromasts, however, indicated that the damaged hair cells in the canal neuromasts appear to be replaced by supporting cells before the new ciliary bundles arise (Fig. 4). This suggests to us that the hair cells were completely destroyed and that new ciliary bundles must arise from new hair cells. To test this hypothesis, it will be necessary to examine the hair cell bodies of the canal neuromasts after gentamicin treatment with transmission electron microscopy.

Regeneration in the presence of an ototoxic drug has not previously been encountered in lateral line neuromasts or fish ears. The only comparable data comes from studies in the chick basilar papilla where it has been demonstrated that hair cell regeneration and differentiation can occur in the presence of ototoxic levels of gentamicin (Girod et al., 1991) and that the regenerated hair cells are not easily damaged by additional kanamycin treatment (Hashino et al., 1994). Hashino et al. (1994) suggested that kanamycin may not be incorporated into the cytoplasm of regenerated hair cells after cumulative administration of the drug and that inhibition of aminoglycoside uptake may be responsible for the drug resistance in the regenerated hair cells (Hashino et al., 1995). How this hypothesis might apply to our results, however, is not clear. In particular, we do not know if hair cell uptake of kanamycin and gentamicin is by similar mechanisms. In addition, the ototoxic drugs were administered to the hair cells by different methods (intramuscular in chicks and by immersion in fish), and this may affect the way that hair cells respond to the drugs.

Even though the suggestion of Hashino et al. (1994) may be correct, there are still other possible explanations for our results that need further study. First, it is possible that gentamicin did damage the hair cell but did not destroy the cell body, as it does in the amniote vestibular system, and that the cilia were regenerated by otherwise intact cells. Second, the response to the ototoxic effect of gentamicin may only occur late in the development of sensory hair cells and before they reach full maturity. Thus, the cells we encountered may only get to a certain stage of development and then they again are 'killed' (or damaged) by the gentamicin. The third possibility is that the regenerated cells are more like the hair cells of the superficial neuromast than those found in untreated canal neuromasts.

It is difficult to differentiate the second and third possibilities without more extended survival time after gentamicin treatment. Our results show that the regenerated ciliary bundles in the canal neuromasts have longer kinocilia than those in the control animals (compare Figs. 4E,F and 2A,B), a characteristic similar to that of normal superficial neuromast hair cells (Fig. 2C,D). The presence of longer kinocilia has been reported in regenerated hair cells in the inner ear of fishes and amphibians (Corwin, 1985; Baird et al., 1993; Lombarte et al., 1993). It also has been reported that, after aminoglycoside ototoxic drug treatment, type I hair cells are replaced by hair cells that are morphologically similar to the type II hair cells in the avian vestibular system (Weisleder and Rubel, 1992) and in the mammalian inner ear (Forge et al., 1993). Even though little is known about the regeneration or developmental stages of sensory hair cells, it is possible that vestibular type I cells develop by going through a type II stage. Therefore, the ciliary bundles we observed in the regenerated canal neuromast hair cells may be in an early stage of development in which they resemble the superficial neuromast hair cells (or the 'type II'). Thus, at the early stages, the hair cells may not be affected by ototoxic drugs because they are more type II-like. As the cells start to become more like mature canal neuromast hair cells (or striolar hair cells in the utricle), however, they may become more susceptible to the gentamicin sulfate. It is possible that, after drug damage, regenerating type II vestibular hair cells are observed first and the more normal type I hair cells would show up several weeks (or more than 60 days) later as in the cristae of birds (Weisleder and Rubel, 1992).

The third possibility, that the regenerated canal hair cells take on the characteristics of superficial neuromast hair cells and stay that way, can only be tested by observing regenerated canal hair cells for extended periods of time after gentamicin exposure.

5. Conclusions

Although further ontogenetic and phylogenetic studies to test the homology of hair cell types of the lateral line



last after 4 days
canal neuromast
of stereocilia.

receptors to the mammalian inner ear are necessary, the results of this study and new observations of the ultrastructural differences between the hair cells in the canal versus superficial neuromasts (Song et al., 1995) suggest that the lateral line receptors have considerable potential for studies on the heterogeneity of mechanosensory hair cells and their susceptibility to ototoxic drugs. The simplicity, versatility, and efficiency of ototoxic drug exposure, and the greater anatomical accessibility of the lateral line system, provide an optimal system for the study of damage and regeneration of sensory hair cells. Even though further investigations are needed to determine whether ototoxic drugs cause functional damage to the lateral line receptors, we do expect this is the case because behavioral studies have indicated that increasing doses of both neomycin and streptomycin produce increasing loss of an animal's ability to localize a wave source (the lateral line reaction) (Kaus, 1987, 1992; Blaxter and Fuiman, 1990).

Acknowledgements

We thank Dr. L.R. Parenti and Dr. R.A. Code, for their comments on the manuscript, Ms. Helen Popper for editorial work, and Mr. T. Maugele for his technical guidance in electron microscopy. This study was supported by grants from NIDCD/NIH DC01986 to J.S., DC01729 to H.Y.Y., NASA NAG 2-787 and ONR N-00014-92-J-1114 to A.N.P. This work is contribution No. 68 from the Laboratory for Biological Ultrastructure, University of Maryland at College Park.

References

- Baird, R.A., Torres, M.A. and Schuff, N.R. (1993) Hair cell regeneration in the bullfrog vestibular otolith organs following aminoglycoside toxicity. *Hear. Res.* 65, 164–174.
- Blaxter, J.H.S. and Fuiman, L.A. (1990) The role of the sensory system of herring larvae in evading predatory fishes. *J. Marine Biol. Assoc. UK* 70, 413–427.
- Chang, J.S., Popper, A.N. and Saidel, W.M. (1992) Heterogeneity of sensory hair cells in a fish ear. *J. Comp. Neurol.* 324, 1–20.
- Coombs, S., Janssen, J. and Webb, J. (1988) Diversity of lateral line systems: evolutionary and functional considerations. In: J. Atema, R.R. Fay, A.N. Popper, and W.N. Tavolga (Eds.), *Sensory Biology of Aquatic Animals*, Springer, New York, pp. 553–593.
- Corwin, J.T. (1977) Morphology of the macula neglecta in sharks of the genus *Carcharhinus*. *J. Morphol.* 152, 341–362.
- Corwin, J.T. (1985) Perpetual production of hair cells and maturational changes in hair cell ultrastructure accompany postembryonic growth in an amphibian ear. *Proc. Natl. Acad. Sci. USA* 82, 3911–3915.
- Forge, A., Li, L., Corwin, J.T. and Nevill, G. (1993) Ultrastructural evidence for hair cell regeneration in the mammalian inner ear. *Science* 259, 1616–1619.
- Girod, D.A., Tucci, D.L. and Rubel, E.W. (1991) Anatomical correlates of functional recovery in the avian inner ear following aminoglycoside ototoxicity. *Laryngoscope* 101, 1139–1148.
- Hama, K. (1965) Some observations on the fine structure of the lateral line organ of the Japanese sea eel, *Lycoteuthis nystromi*. *J. Cell Biol.* 24, 193–210.
- Hama, K. (1969) A study on the fine structure of the saccular macula of the goldfish. *Z. Zellforsch. Mikrosk. Anat.* 94, 155–171.
- Hama, K. (1978) The fine structure of the pit organs of the common Japanese sea eel, *Conger myriaster*. *Cell Tissue Res.* 189, 375–388.
- Hashino, E., Salvi, R.J. and Shero, M. (1994) Are regenerated hair cells more resistant to aminoglycoside ototoxicity than pre-existing hair cells? *Abstr. Assoc. Res. Otolaryngol.* 17, 115, 459.
- Hashino, E., Shero, M. and Salvi, R.J. (1995) Possible inhibitory mechanisms for kanamycin uptake in regenerated hair cells. *Abstr. Assoc. Res. Otolaryngol.* 18, 46, 183.
- Jensen, J.C. (1984) On the polarization and innervation of the pars inferior sensory epithelia of the herring labyrinth. *Acta Zoologica* 65, 61–74.
- Jørgensen, J.M. (1989) The fine structure of the pit organs of the common Japanese sea eel, *Conger myriaster*. In: S. Coombs, P. Görner, and H. Münz (Eds.), *The Mechanosensory Lateral Line*, Springer, New York, pp. 113–145.
- Kaus, S. (1987) The effect of aminoglycoside antibiotics on the lateral line organ of *Aplocheilichthys lineatus* (Cyprinodontidae). *Acta Otolaryngol. (Stockh.)* 103, 291–298.
- Kaus, S. (1992) The influence of calcium on the ototoxicity of aminoglycosides. *Acta Otolaryngol. (Stockh.)* 112, 83–87.
- Kroese, A.B.A. and van Netten, S.M. (1989) Sensory transduction in lateral line hair cells. In: S. Coombs, P. Görner, and H. Münz (Eds.), *The Mechanosensory Lateral Line*, Springer, New York, pp. 265–284.
- Landacre, F.L. and Conger, A.C. (1913) The origin of the lateral line primordia in *Lepidosteus osseus*. *J. Comp. Neurol.* 23: 575–633.
- Lindeman, H.H. (1969) Regional differences in sensitivity of the vestibular sensory epithelia to ototoxic antibiotics. *Acta Otolaryngol.* 67, 177–189.
- Lombarte, A., Yan, H.Y., Popper, A.N., Chang, J.S. and Platt, C. (1993) Damage and regeneration of hair cell ciliary bundles in a fish ear following treatment with gentamicin. *Hear. Res.* 64, 166–174.
- McCormick, C.A. (1982) The organization of the octavolateralis area in actinopterygian fishes: A new interpretation. *J. Morphol.* 171, 159–181.
- Münz, H. (1985) Single unit activity in the peripheral lateral line system of the cichlid fish *Sarotherodon niloticus*. *J. Comp. Physiol. A* 157, 555–568.
- Münz, H. (1989) Functional organization of the lateral line periphery. In: S. Coombs, P. Görner, and H. Münz (Eds.), *The Mechanosensory Lateral Line*, Springer, New York, pp. 285–298.
- Northcutt, R.G. (1980) Central auditory pathways in anamniotic vertebrates. In: A.N. Popper and R.R. Fay (Eds.), *Comparative Studies of Hearing in Vertebrates*, Springer, New York, pp. 79–118.
- Northcutt, R.G. (1989) The phylogenetic distribution and innervation of cranial mechanoreceptive lateral lines. In: S. Coombs, P. Görner, and H. Münz (Eds.), *The Mechanosensory Lateral Line*, Springer, New York, pp. 17–78.
- Platt, C. and Yan, H.Y. (1993) Dramatic differences between fish species for gentamicin pharmacotoxicity. *Abstr. Assoc. Res. Otolaryngol.* 16, 562.
- Platt, J.B. (1896) Ontogenetic differentiations of the ectoderm in *Necturus*. Study II. On the development of the peripheral nervous system. *Quart. J. Microsc. Sci.* 38: 485–547.
- Popper, A.N. and Hoxter, B. (1981) The fine structure of the sacculus and lagena of a teleost fish. *Hear. Res.* 5, 245–263.
- Popper, A.N., Platt, C. and Edds, P.L. (1992) Evolution of the vertebrate inner ear: An overview of ideas. In: D.B. Webster, R.R. Fay and A.N. Popper (Eds.), *Evolutionary Biology of Hearing*, Springer, New York, pp. 49–57.
- Richardson, G.P. and Russell, I.J. (1991) Cochlear cultures as a model system for studying aminoglycoside induced ototoxicity. *Hear. Res.* 53, 293–311.
- Saidel, W.M., Presson, J.C. and Chang, J.S. (1990a) S-100 immunoreactivity identifies a subset of hair cells in the utricle and saccule of a fish. *Hear. Res.* 47, 139–146.

Saidel, W.M.
logical
utricle.
Song, J. (1
line rec
649.17.
Song, J., Ji
growth
superfi
gol. 18
Song, J. :
innerva
platyph
Song, J. a
lateral
Behav.
Van Berge
Neff (1
New Y
Wegner, N

- ocular macula of 171.
- of the common 189, 375–388.
- erated hair cells ore-existing hair
- hibitory mecha- s. Abstr. Assoc.
- ion of the pars ta Zoologica 65,
- s of the common . Görner, and H. nger, New York,
- s on the lateral . Acta Otolaryn-
- ity of aminogly-
- transduction in H. Münz (Eds.), k, pp. 265–284.
- the lateral line 3: 575–633.
- of the vestibulo- Otolaryngol. 67,
- Platt, C. (1993) s in a fish ear 66–174.
- lateralis area in ohol. 171, 159–
- eral line system Physiol. A 157,
- e periphery. In: echanosensory
- amniotic verte- ative Studies of 118.
- innervation of P. Görner, and Springer, New
- een fish species Otolaryngol. 16,
- oderm in *Nec-* rous system.
- ne sacculus and
- f the vertebrate . Fay and A.N. ger, New York,
- es as a model ity. Hear. Res.
- 100 immuno- and sacculle of
- Saidel, W.M., A.N. Popper, and Chang, J.S. (1990b) Spatial and morphological differentiation of trigger zones in afferent fibers to the teleost utricle. *J. Comp. Neurol.* 302, 629–642.
- Song, J. (1993) Morphology, distribution and innervation of the lateral line receptors in gobies (Teleosts). *Soc. Neurosci. Abstr.* 19, 1580, 649.17.
- Song, J., Jia, X. And Popper, A.N. (1995) Differences in postembryonic growth and ultrastructure of sensory and supporting cells in canal and superficial neuromasts of the lateral line. *Abstr. Assoc. Res. Otolaryngol.* 18, 44, 175.
- Song, J. and Northcutt, R.G. (1991a) Morphology, distribution and innervation of the lateral line receptors of the Florida gar, *Lepisosteus platyrhincus*. *Brain Behav. Evol.* 37, 10–37.
- Song, J. and Northcutt, R.G. (1991b) The primary projections of the lateral line nerves of the Florida gar, *Lepisosteus platyrhincus*. *Brain Behav. Evol.* 37, 38–63.
- Van Bergeijk, W.A. (1967) The evolution of vertebrate hearing. In: W.D. Neff (Ed.), *Contribution to Sensory Physiology*, Vol. 2, Academic, New York, pp. 1–49.
- Wegner, N. (1982) A qualitative and quantitative study of a sensory epithelium in the inner ear of a fish (*Colisa labiosa*, Anabantidae). *Acta Zoologica* 63, 133–146.
- Weisleder, P. and Rubel, E.W. (1992) Hair cell regeneration in the avian vestibular epithelium. *Exp. Neurol.* 115, 2–6.
- Wersäll, J. (1956) Study on the structure and the innervation of the sensory epithelium of the cristae ampullares in the guinea pig. A light and electron microscopic investigation. *Acta Otolaryngol. (Suppl.)* 126, 1–85.
- Wersäll, J. (1960) Vestibular receptor cells in fish and mammals. *Acta Otolaryngol. (Suppl.)* 163, 25–29.
- Wersäll, J. and Hawkins, J.E., Jr. (1962) The vestibular sensory epithelium in the cat labyrinth and their reactions in chronic streptomycin intoxication. *Acta Otolaryngol.* 54, 1–22.
- Wersäll J., Björkroth, B. Flock, Å. and Lundquist P.-G. (1971) Sensory hair fusion in vestibular sensory cells after gentamicin exposure. *Arch. Klin. Exp. Ohr. Nasu. Kehlk. Heilk.* 200, 1–14.
- Yan, H.Y., Saidel, W.M., Chang, J.S., Presson, J.C. and Popper, A.N. (1991) Sensory hair cells of the fish ear, evidence of multiple type: based on ototoxicity sensitivity. *Proc. R. Soc. Lond. B* 245, 133–138

ARTICLES should deal with original research not previously published or being considered for publication elsewhere. The act of submitting a manuscript to the journal carries with it the right to publish that paper and implies the transfer of the Copyright from the author to the publisher. This transfer will ensure widest possible dissemination of information. Papers should be submitted to:

Professor Aage R. Møller (Editor-in-Chief),
P.O. Box 99187
Pittsburgh, PA 15233-4187, USA
Cleat Szczepaniak (Editorial Secretary)
Telephone (412) 648-3901

Address for Courier Mail
Department of Neurological Surgery,
Suite B-400 Presbyterian-University Hospital,
200 Lothrop Street
Pittsburgh, PA 15213, USA

One or two reviewers from the Editorial Board may be suggested by the authors, and these will be taken into consideration by the Editor-in-Chief: Fax: (412) 648-8924.

MANUSCRIPTS in English, should be complete in all respects, i.e., originals plus three copies of all items including figures and tables: *Please note* four original sets of all micrographs must be submitted. The manuscript should be typed with double spacing and wide margins, on one side of the paper only, and full-length papers should be divided into sections (e.g. Introduction, Materials and Methods, Results, Discussion, etc.). Authors' full names and academic addresses should be given on the title page, as well as an address for correspondence along with their telephone number and fax (or telex) numbers to enable the Editor/Publisher to contact them. Papers describing experimental work on living subjects must indicate that the reported investigations have been performed in accordance with the principles of the Declaration of Helsinki. Three copies of any article by the authors cited in the manuscript and noted in the reference list as being 'in press' must accompany the submitted manuscript.

Title should be informative and preferably not exceed 85 characters, including spaces. Extraneous words such as 'study', 'investigation', etc., should be avoided. Summary not exceeding 200 words should be given at the beginning of the paper, followed by 3-6 indexing terms (key words).

If a manuscript involves experimentation on animals or humans, a statement must be made at the end of the Methods section relative to the following: The care of humans ... or the care and use of the animals reported on in this study were approved by (one of the following)

- 1) A grant application agency
 - 2) A specific university's Animal Care and Use Committee provided that the committee adheres to the guidelines of the Declaration of Helsinki.
- In both cases please provide a grant title name/number.

If the authors cannot provide approval as outlined above they must, in the case of human studies, state that the experiments were performed in accordance with the guidelines of the Declaration of Helsinki, a copy of which is available from the Editor's office upon request. Manuscripts not complying with the above will be returned to the authors.

Tables should be typed, with double spacing, each on a separate sheet, numbered consecutively, and should only contain horizontal lines. A short descriptive heading should be given above each table, and possible footnotes and explanations underneath.

Figures should be original laser prints or very sharp, well-contrasting prints on glossy paper, suitable for immediate reproduction. Half-tone figures should be in black-and-white, very sharp, well-contrasting, and on glossy paper. Any lettering in the figures should be large enough to stand photographic reduction. Contributors should prepare suitably their figures for either one column width (84 mm) or the entire page width (175 mm). The maximum height is 240 mm. The Publisher will determine the degree of any reduction or enlargement required and, in general, line drawings will be reduced to one column width if possible. Authors may, however, specifically request a larger reproduction. Particular requests should be typed on the relevant figure legend page. Micrographs will usually not be reduced unless the reduction involved is small or the height of the figure necessitates reduction. Figure legends should be typed, with double spacing, on a separate sheet.

References should be assembled in alphabetical order on a separate sheet. In the text they should be referred to by name and year (Harvard System). More than one paper from the same author in the same year must be identified by the letters a, b, c, etc., placed after the year of publication. In the text, when referring to a work by more than two authors, the name of the first author should be given followed by et al. Literature references must consist of names and initials of all authors, year, title of paper referred to, abbreviated title of periodical, volume number and first and last page numbers of the paper. Periodicals (i), books (ii) and multi-author books (iii) should accord with the following examples:

Sellick, P.M., Patuzzi, R. and Johnstone, B.M. (1982) Measurement of basilar membrane motion in guinea pig using the Mössbauer technique. *J. Acoust. Soc. Am.* 72, 131-141. (i)

Békésy, G. von (1960) Experiments in Hearing. McGraw-Hill, New York. (ii)

Wilson, J.P. and Evans, E.F. (1983) Some observations on the 'passive' mechanics of cat basilar membrane. In: W.R. Webster and L.M. Aitkin (Eds.), *Mechanisms of Hearing*, Monash University Press, Clayton, Australia, pp. 30-35. (iii)

Abbreviations of journal titles should conform to the List of Serial Title Word Abbreviations, International Serials Data System, 20, rue Bachaumont, 75002 Paris, France. ISBN 2-904938-02-8.

Typescripts should be carefully checked before submission to obviate alterations after acceptance.

Submissions on disc. The manuscript, plus three copies, a floppy disc of the word-processed manuscript and a list of any non-standard characters that were used on the disc should be submitted in the usual manner. The preferred storage medium is a 5.25 or 3.5 inch disc in MS-DOS or MS-DOS compatible format, although other systems, e.g. Macintosh, are acceptable. Please specify the type of computer and word processing package used (do not convert your text file to plain ASCII). It is essential that the disc submitted for publication (i.e. after acceptance) is the final version of the article and exactly matches the accompanying manuscript. Accurate keyboard practice is essential, for instance, one (1) and 'el' (!) must be clearly different. Non-reproducible characters should not be left as a blank space in the file but replaced by characters not used elsewhere: their use must be consistent and clearly stated. Recommended reading: *Chicago Guide to Preparing Electronic Manuscripts*, The University of Chicago Press, Chicago (1987). The Publisher is under no obligation to use the submitted floppy disc, but will make every attempt to do so.

PROOFS will be sent to the first-named author of an article, unless an alternative is requested on the title page of the manuscript. They should be checked carefully and returned by airmail within 3 days of receipt. Only printer's errors may be corrected: no changes in or additions to the edited manuscript will be allowed at this stage.

REPRINTS may be ordered by filling in and returning to the Publishers the order form sent to the authors with their proofs. 50 free reprints per contribution will be made available.

A CUMULATIVE INDEX for *Hearing Research*. Volumes 1 through to the current volume is available upon an IBM-formatted 5.25" inch floppy disc (ASCII) free of charge. Please send \$ 10.00 (to cover postage and handling) by personal check or money order payable to: Cleat Szczepaniak, Hearing Research, P.O. Box 99187, Pittsburgh, PA 15233-4187, USA. The index is also available on the Internet: gopher://spib.nice.edu:70/11/SPIB/bibliography/hr

No responsibility is assumed by the Publisher for any injury and/or damage to persons or property as a matter of products liability, negligence or otherwise, or from any use or operation of any methods, products, instructions or ideas contained in the material herein. Because of rapid advances in the medical sciences, the Publisher recommends that independent verification of diagnoses and drug dosages should be made.

Although all advertising material is expected to conform to ethical (medical) standards, inclusion in this publication does not constitute a guarantee or endorsement of the quality or value of such product or of the claims made of it by its manufacturer.

Submission of a paper to this journal entails the author's irrevocable and exclusive authorization of the publisher to collect any sums or consideration for copying or reproduction payable by third parties (as mentioned in article 17 paragraph 2 of the Dutch Copyright Act of 1912 and in the Royal Decree of June 20, 1974 (S.351) pursuant to article 16b of the Dutch Copyright Act of 1912) and/or to act in or out of Court in connection therewith.