

NASA-CR-201825

100-10-102  
16-11-

NAG 2-341

FINAL

JUL 13 1986  
CASI

THE EFFECTS OF HYPERGRAVIC FIELDS ON NEURAL SIGNALLING  
IN THE HIPPOCAMPUS

CONTENTS

- A. Summary . . . . . 1
- B. Research Plan . . . . . 2
  - 1. Introduction . . . . . 2
  - 2. Background . . . . . 2
  - 3. Specific questions . . . . . 10
  - 4. Rationale and experimental procedures . . . . . 10
  - 5. Data interpretation . . . . . 14
- C. Significance . . . . . 15
- D. References . . . . . 16
- E. Progress report (1985-1987) . . . . . 19
  - 1. Hippocampal studies . . . . . 19
  - 2. Vestibular and auditory system . . . . . 20
  - 3. Thermoregulatory system . . . . . 21
- F. Biographical sketch of PIs . . . . . 24
- G. Budget . . . . . 49
- H. Budget justification . . . . . 50
- I. Available facilities . . . . . 50
- K. Appendix A Effects of pH and temperature on hippocampal population spikes . . . . . A1
- Appendix B Temperature effects on evoked potentials of hippocampal slices from noncold-acclimated, cold-acclimated and hibernating hamsters . . . . . B1
- Appendix C Simulation of the effects of temperature on hippocampal neurons . . . . . C1

A. SUMMARY

Experiments in this grant renewal have as a general long-term objective the elucidation of the effects of altered gravitational fields on neural regulatory mechanisms. More specifically, they focus on the effects of the neurotransmitter serotonin in the hippocampus. We have selected the hippocampus for study because rats flown on Spacelab-3 showed an increased number of serotonergic receptors in this region of the central nervous system after 7 days in space. A more recent study has shown that rats exposed to a hypergravic field of 2.0 G for 7 days show a decreased number of receptors. We propose to examine population and single cell electrical activity associated with these receptors using the in vitro hippocampal slice preparation. After development of experimental techniques, we will utilize centrifuges at the Chronic Acceleration Research Laboratory to expose animals to a hypergravic environment. Receptor numbers change relatively slowly so that altered activity can be studied in rats at 1 G following prolonged exposure to an increased gravitational field. This study will provide information on the modulation of signaling in the hippocampus by serotonin before and after exposure to hypergravic fields.

## B. RESEARCH PLAN

### 1. Introduction

The goal of this grant is to study the effect of hypergravic fields on the modulation of hippocampal electrical activity by serotonin (5-HT). The proposed study represents a shift from our previous NASA grants covering three diverse areas in neurobiology (thermoregulation, vestibular and auditory brainstem evoked responses, and the hippocampus) to consideration of only one of these areas, the hippocampus. (Thus no experiments are proposed in the areas of thermoregulation or brainstem evoked responses, and the only continuing work in these areas will be to edit manuscripts covering work supported by our previous NASA grants.)

To place our proposed hippocampal experiments in context with relevant Spacelab-3 experiments and hypergravic experiments, two experiments on receptor changes in animals exposed to altered gravitational fields are first described. Our experiments build on these structural/biochemical observations and extend investigations to related electrical activity at 1 G and in hypergravic fields. The background continues with a review of past studies at 1G related to effects of serotonin on hippocampal electrical activity (i.e., population spikes, intracellular potentials).

### 2. Background

Increased number of serotonin receptors in rats aboard Spacelab-3. Miller et al. (1985) determined the number of receptors for several neurotransmitters in selected brain areas in animals after their return from space. Six male Sprague-Dawley rats were exposed to microgravity for 7 days aboard Spacelab-3. Twelve to 14 hours after return to 1 G, the rats were sacrificed and brain tissue was removed, frozen, and subsequently assayed for receptor number and affinity. Serotonin binding in the hippocampus of these rats was 49% greater compared to that from a control group maintained at 1 G ( $120 \pm 5$  in the flight group vs  $81 \pm 11$  fmoles/mg protein in the control group;  $p < 0.05$ ). The binding of serotonin in the lateral frontal cortex was unaffected by exposure to microgravity. Moreover, in the initial survey of other neural transmitters in different regions of the brain, changes in binding were not generally observed. Thus, the hippocampus was a region of the central nervous system that showed cellular changes attributable to exposure to a microgravity environment.

Decreased number of serotonin receptors in rats exposed to 2 G. The experiments reported by Miller et al. (1985), while demonstrating an effect at microgravity, left open the question of whether hypergravic fields had any effect on serotonergic receptors. Very recently, Miller et al. (1986) reported at a Tokyo conference the results of studies in which they measured serotonergic binding in rats after exposure to a 2 G field for seven days. Twelve rats were exposed to chronic acceleration on an 18 foot diameter centrifuge and then binding was determined using the ligand [ $^3$ H]-5-hydroxytryptamine. A 27% decrease in receptor number was observed in rats exposed to the 2 G field.

These studies by Miller et al. (1985, 1986) show that hypergravic and microgravity environments have opposite effects on hippocampal 5-HT receptor number. In both environments the effects appear to be terminal field specific, since changes in serotonin binding were not observed in the lateral frontal cortex. Although receptor changes have been described in both microgravic and hypergravic environments, there are no parallel electrophysiological studies of serotonergic effects on neural activity. It is therefore unknown if the changes in receptor number alter the functional characteristics of the neurons. It is this issue that this grant will address.

Neurochemistry and multiple serotonergic receptors. The population of 5-HT receptors in rat cerebral cortex and hippocampus can be divided into two types denoted 5-HT<sub>1</sub> and 5-HT<sub>2</sub> receptors (Peroutka and Snyder, 1979; Creese and Snyder, 1978), and recent studies (Pedigo et al., 1981) suggested a further subdivision of 5-HT<sub>1</sub> receptors into two subtypes, 5-HT<sub>1A</sub> and 5-HT<sub>1B</sub>, a classification based on the biphasic inhibition of spiroperidol (or spiperone) on serotonin binding. Binding inhibited by low concentrations of spiroperidol defined the 5-HT<sub>1A</sub> site, whereas the component that was unaffected by low concentrations of spiroperidol defined the 5-HT<sub>1B</sub> site. Beck et al. (1985), De Vivo and Maavani (1986), Mauk et al. (1985), and Andrade and Nicoll (1985) have identified serotonergic receptors found on hippocampal pyramidal cells as 5-HT<sub>1A</sub> receptors. This grant covers effects of binding of serotonin to 5-HT<sub>1A</sub> receptors.

The effect of serotonin on population spikes. We plan experiments on changes in hippocampal electrical activity and the effects of serotonin using both extracellular and intracellular procedures. Extracellular methods (described in this section) allow assessment of the system as a whole, while intracellular methods allow a closer look at the cellular mechanisms underlying serotonin's effects. The results of the proposed experiments would provide baseline data for future studies using rats exposed to microgravic environments.

Several studies have shown that serotonin decreases the amplitude of a population response of hippocampal neurons (Beck et al., 1985; Beck and Goldfarb, 1985; Rowan and Anwyl, 1985; Mauk et al., 1985). (The decrease in amplitude is often preceded by a small transient increase.) When Schaffer collaterals, axons forming a fiber tract in the hippocampus, are stimulated, pyramidal cells are excited and generate a synchronous burst of action potentials. The electrical fields of the action potentials add to produce an easily detected potential, a population spike. In experiments where serotonin was superfused over the tissue and then washed out with buffer, Beck and Goldfarb (1985) showed that serotonin reversibly decreased the amplitude of the population spike. The magnitude of this decrease was concentration dependent and was greater on submaximal than on maximal population spikes. The EC<sub>50</sub> for serotonin was 3.2  $\mu$ M. Repeated applications of a maximal dose of serotonin did not produce tachyphylaxis. Beck and Goldfarb (1985) concluded that the reversibility, reproducibility, and concentration dependence of the serotonergic response in the rat hippocampal slice preparation make the hippocampus useful for pharmacological experiments on the modulatory role of serotonin.

Recent pharmacological studies involving measurement of hippocampal population spike amplitudes have indeed further characterized serotonergic mechanisms. Thus Beck et al. (1985) were among the first to suggest that the decrease in population spike amplitude involved a 5-HT<sub>1A</sub> receptor because spiperone at 100 nM shifted serotonin and 5-carboxyamidotryptamine concentration-response curves to the right. Mauk et al. (1985) also observed that 5-HT<sub>1A</sub> agonists depressed the amplitude of population spikes. Slices from Wistar rats were placed in a submersion chamber, superfused with a continuous flow of modified Krebs Ringer and maintained at 33-34°C. Evoked potentials following stimulation of Schaffer collaterals were recorded in the pyramidal layer. When field potentials had been stable for a least 20 min, dose-response curves were obtained by applying increasing concentrations of serotonergic agonists every 30 min. Application of the agonists buspirone, TVX-Q 7821, and DPAT produced a dose dependent reduction of the population spike. Effective concentrations were 25-100 µM.

The method of measuring spike amplitudes allows assessment of the response of a population of pyramidal cells to a variety of compounds. Beck and Goldfarb (1985) describe in detail superfusion procedures for obtaining concentration-response curves. They then applied these procedures (Beck et al., 1985) to show shifts in the concentration-response curves for a variety of ligands (serotonin alone and in the presence of 100nM spiperone or 100nM ketanserin). This procedure should prove sufficiently sensitive to detect an altered response to serotonin in animals exposed to hypergravic environments if the decrease in amplitude of the population spike is similar in magnitude to the 27% decrease in receptor number found in binding studies.

The effect of temperature, pH, and stimulus frequency on population spikes. Bath temperature has a marked effect on population spike parameters (amplitude and width) and pH a much smaller effect (Hooper et al., 1985; Thomas et al., 1986; appendix A). In experiments designed to detect changes in the physiological response to serotonin in animals exposed to a hypergravic environment, these factors must be controlled or monitored.

Experiments on hippocampal slices have been done at various bath temperatures -- some investigators maintain bath temperature between 32 and 35°C, others between 29 and 30°C. So that results can be more readily compared with other studies, one approach is to perform critical experiments over a range of temperatures.

Stimulus frequency can also modify the height of population spikes and the activation of intracellular events. Long-term potentiation (LTP) is a widely studied model of activity-dependent change in synaptic efficacy that has attracted wide interest as the basis for information storage in the mammalian brain. LTP is induced by increasing stimulus frequency above about 3 shocks/sec, and is reliably induced by frequencies greater than 10 Hz. Its onset occurs in seconds and can last for weeks in the intact animal. During LTP, the amplitude of a population spike is greatly enhanced. The role of N-methyl-D-aspartate (NMDA) receptors in inducing hippocampal LTP has been recently established by a variety of

studies in different laboratories (Collingridge and Bliss, 1987; Collingridge et al., 1983; Harris et al., 1984; Peet et al., 1986).  $Mg^{++}$  blocks NMDA-gated channels in a voltage dependent manner. When the cell is at its resting potential, many of the channels are blocked, and when the cell is depolarized  $Mg^{++}$  blockade is alleviated. Calcium then enters the cell via these NMDA channels and triggers the sequence of events that culminate in LTP.

Stimulus frequency must be carefully controlled in studies where population spike amplitude is measured because high but not low frequency stimulation induces LTP. As an added complication, Bliss et al. (1983) found reduction of LTP in the dentate gyrus of the rat following depletion of serotonin, and no effect following depletion of norepinephrine (NE). They used an in vivo preparation and it was not possible to determine whether the effect on LTP was due to depletion in the hippocampus or depletion of other brain areas. Stanton and Sarvey (1985) used the in vitro slice preparation and found that while NE reduced the occurrence of LTP in the dentate, serotonin did not prevent LTP in the hippocampus in either CA1 or the dentate. Further experiments are required to fully clarify the involvement of serotonin in LTP. On one hand, by using high frequency stimulation to evoke LTP, the involvement of serotonin on LTP can be further studied. On the other hand, by keeping the stimulus frequency less than 1 Hz, LTP is not evoked (NMDA channels are not opened) and possible effects of serotonin on LTP are avoided.

The effect of serotonin on intracellular potentials. There are many recent electrophysiological studies of the hippocampus using intracellular rather than extracellular techniques. In many cases intracellular techniques have an advantage; namely, that intermediate steps in the overall response of the system can be monitored and effects on particular cellular mechanisms can be tested. For example, transmembrane potential and membrane resistance provide information on combined receptor and channel activity. We propose to measure both potentials and resistance in rats and hamsters exposed to 2G fields. In addition, factors such as the potential role of GTP/GDP binding proteins (G proteins), intracellular messengers, and various receptor types can be investigated using intracellular techniques.

Using intracellular techniques, Andrade and Nicoll (1985) provided further electrophysiological evidence, in agreement with previous pharmacological studies, that serotonin binding appears to be predominantly of the 5-HT<sub>1</sub> type. Recordings were obtained from pyramidal cells of the CA1 region. Serotonin administered either in the bath (300nM - 20 $\mu$ M) or by microiontophoresis elicited a dose-dependent hyperpolarization (that could be as large as 20 mV) accompanied by reduced input resistance. No depolarizing responses to serotonin were ever observed. Administration of the selective 5-HT<sub>1A</sub> agonist 8-OHDPAT (200 nM - 4 $\mu$ M) also hyperpolarized these cells. The classical serotonin antagonists cyproheptadine and cinanserin were relatively ineffective in antagonizing responses to iontophoretically applied serotonin, with IC<sub>50</sub>s of 50  $\mu$ M and 100  $\mu$ M respectively. The selective 5-HT<sub>2</sub> antagonist ketanserin did not reduce serotonin responses in concentrations up to 200  $\mu$ M. The ergot methysergide elicited a small hyperpolarization and

reduced the serotonin response with an IC<sub>50</sub> of 30  $\mu$ M. Similarly, the non-benzodiazepine anxiolytic, buspirone, which has been reported to be potent and selective at displacing serotonin from hippocampal membranes, also elicited a small hyperpolarization (1-4 mV) and reduced serotonin responses (IC<sub>50</sub> = 3 $\mu$ M). Thus, a variety of intracellular results are consistent with the proposal that 5-HT<sub>1</sub> receptors are present on pyramidal cells in the hippocampus.

Intracellular studies have also characterized the channels opened by serotonin (Segal, 1980; Jahnsen, 1980). Serotonin reduces the input resistance and hyperpolarizes the cell, results consistent with the opening of some type of potassium ion channel (Segal, 1980; Andrade et al., 1986). Intracellular measurements have also indicated that G proteins are involved in the coupling of signals from serotonergic receptors to the opening of these potassium ion channels, a major advance in understanding serotonergic mechanisms (see below).

Receptors, G proteins and potassium ion channels. Membrane receptors often signal cellular responses via a coupling system. The measurement of transmembrane potential and resistance serves to characterize this system, from the binding of serotonin through the opening of potassium ion channels. Since we plan to make these measurements in our experiments and in this way characterize plasma membrane mechanisms for the recognition and transduction of serotonergic signals, the coupling system in pyramidal cells is described in detail.

Andrade et al. (1986) proposed that two classes of receptors in the hippocampus, 5-HT and GABA<sub>B</sub>, act through a G protein to open potassium ion channels. Thus, receptors activated by a diffuse serotonergic fiber system share the same potassium channels as do receptors activated by local inhibitory interneurons that release GABA. The critical experiment pointing toward a shared population of channels was the demonstration that potassium currents evoked by the agonists are not additive. Evidence that distinct receptors for serotonin and GABA are present in the membrane includes the observation that when spiperone (a 5-HT receptor blocker) was applied in sufficient concentration to completely block the hyperpolarization evoked by serotonin, hyperpolarization could still be readily evoked by application of the GABA<sub>B</sub> agonist baclofen.

Andrade et al. (1986) tested the system to determine if G proteins were involved in the transduction of the signal from the receptor to the opening of the channels. A model involving serotonergic receptors, G proteins, and potassium channels is shown in Figure 1. This model (proposed by Andrade et al., 1986) involves the following steps in the shuttling of the G-protein between receptors (5-HT or GABA<sub>B</sub>) and the ion channel (which, when open, allows potassium ions to leave the cell and the cell to hyperpolarize). When GTP is bound to the G-protein, the GTP-G-protein complex associates with the potassium ion channel, and the channel opens. When associated with the channel, the G-protein also acts as a GTPase, hydrolyzing its bound GTP to GDP. As a result, the G-protein then dissociates from the channel, and the channel




FIGURE 1. Sketch of a pyramidal cell showing a G protein coupling 5-HT receptors to potassium ion channels. serotonin binds with a 5-HT<sub>1</sub> receptor to activate a G protein (G<sub>i</sub>) that shuttles through the membrane (arrows A and C) to bind with a potassium channel protein, thus opening the channel. The potassium ion efflux through the open potassium channel leads to hyperpolarization and cell inhibition. The neurotransmitter GABA binds with a GABA<sub>B</sub> receptor that activates the same G protein (arrow B) and hence also opens K<sup>+</sup> channels. This process, the opening of potassium ion channels by either serotonin or GABA, can be inhibited by increased activity of protein kinase C as shown at lower left. Serotonin also inhibits the membrane bound enzyme adenylate cyclase as shown at lower right.

closes. The GDP-G-protein shuttles to the receptor through the membrane to combine with 5-HT-receptor or GABA-receptor complexes. When associated with these complexes, GTP displaces GDP at the nucleotide site on the coupling protein, and this GTP-G-protein shuttles back through the plasma membrane to repeat the cycle.

The experimental evidence that G proteins are involved in the coupling between the receptors and the potassium ion channel was demonstrated by experiments showing that the response to neurotransmitters is altered by treatments that modify G proteins. After intraventricular injection of pertussis toxin, a specific toxin for some G proteins, pyramidal cells from treated rats no longer showed hyperpolarization following application of either serotonin or GABA even though in other respects cellular activity was the same as controls (e.g., action potentials could be generated by depolarizing the cell). Additionally, in slices treated with GDP $\beta$ S, a compound that binds with G proteins and maintains them in the inactive state, application of serotonin or GABA<sub>B</sub> did not induce pyramidal cell hyperpolarization.

These experiments show that a G protein is associated with the transduction of the signal from the membrane receptors to potassium ion channels but do not in themselves show a direct coupling of G proteins to potassium ion channels. A link that involves a second messenger within the cytoplasm could be involved, and additional experiments were carried out to determine if this was the case. By eliminating three intracellular messengers as being important in this coupling, Andrade *et al.* (1986) suggest that the G-proteins directly couple to potassium ion channels in the plasma membrane. Experiments excluding second messengers described below illustrate the advantage of the slice preparation in sorting out cellular mechanisms.

First, Andrade *et al.* (1986) considered whether cAMP served as an intracellular messenger. Madison and Nicoll (1986a,b) had previously shown that norepinephrine (NE) binds to beta receptors to increase intracellular levels of cAMP. The elevated levels of cAMP reduce or block the current through potassium ion channels opened by intracellular calcium ions. These calcium-activated potassium channels hyperpolarize the cell, leading to a waveform called an AHP (afterhyperpolarization). The experiments to rule out an intracellular link of cAMP to the effects of serotonin or baclofen (a GABA<sub>B</sub> agonist) involved bathing the tissue for at least 15 minutes in 8-bromo-cAMP, a membrane-soluble analog of cAMP. Loading the cells with 8-bromo-cAMP (to saturate calcium-activated potassium channel binding sites) had no effect on the responses of the cell to serotonin or to baclofen. These cells were loaded with sufficient 8-bromo-cAMP to completely block the effects of NE on the AHP. The fact that NE had no effect and the fact that the effects of serotonin and baclofen were unchanged negated the alternative that cytosolic cAMP was necessary for serotonin to exert its effect.

Second, Andrade *et al.* (1986) considered whether intracellular levels of calcium ions formed a link between serotonin or baclofen and the influx of potassium ions. Increased levels of calcium ion in hippocampal pyramidal cells affect several types of potassium ion channels. The experiments described above ruled out the possibility that serotonin



opened one type of calcium-activated potassium channel. In additional experiments, involving the use of tetraethylammonium (TEA), they ruled out the possibility that calcium opened other types of channels. Thus Andrade et al. (1986) concluded that calcium ions did not serve as a second messenger in the serotonergic response.

Third, Andrade et al. (1986) performed experiments they interpreted as ruling out a role for inositol phospholipids as a second messenger between a G protein and potassium channels. The activation of protein kinase C (with the activator phorbol-12,13-dibutyrate) showed that protein kinase C is not involved in a second messenger pathway. In addition, inositol triphosphate often acts by releasing calcium from intracellular storage sites, but changes in intracellular calcium had no effect on responses evoked by serotonin or GABA.

The lack of evidence for a second messenger involving inositol phospholipids, together with a lack of evidence for other second messengers (cAMP and calcium), led Andrade and Nicoll (1985) to propose direct coupling between G proteins and potassium proteins. The coupling can modify and can be modified by other cellular mechanisms. The schematic diagram of a pyramidal cell (Figure 1) illustrates added interactions in the process initiated by serotonin binding on hippocampal pyramidal cells. First, De Vivo and Maayani (1986) have reviewed evidence showing that 5-HT<sub>1A</sub> agonists inhibit adenylate cyclase. In addition, the serotonergic response can be inhibited by activation of protein kinase C using phorbol-12, 13-dibutyrate (Andrade et al., 1968). Figure 1 includes these added interactions to indicate mechanisms associated with events following binding in addition to the G-protein coupling between the 5-HT<sub>1A</sub> receptor and K<sup>+</sup>-channel.

The extracellular and intracellular experiments cited above provide the background for further study of serotonergic mechanisms as modified by gravitational fields. Data are consistent with the proposals that serotonin receptors in the hippocampus conform to the 5-HT<sub>1A</sub> classification, and that the hippocampal slice preparation can be used to advantage to study functional effects of decreased receptor number. Coupled with the studies on altered binding of serotonin in microgravity (Miller et al., 1985) and in hypergravity (Miller et al., 1986), a basis is formed for further electrophysiological studies on the effect of hypergravic fields on serotonergic responses. The proposed study is on cellular events associated with 5-HT receptors after exposure of animals to a hypergravic environment.

We plan experiments on changes in hippocampal electrical activity and the effects of serotonin using both extracellular and intracellular procedures. Extracellular methods allow assessment of the modulatory effects of serotonin on a major hippocampal circuit, signals over Schaffer collaterals to hippocampal pyramidal cells. Modulation can be monitored by measuring the amplitude of the population spike. Intracellular methods allow a closer look at the effect of serotonin on intermediate steps in the process. Within the plasma membrane a signal is coupled from 5-HT<sub>1A</sub> receptors to potassium ions channels via a G protein, and these membrane mechanisms are reflected by changes in transmembrane potentials and membrane resistance. When potassium ion

channels open in many pyramidal cells, the hyperpolarization decreases the amplitude of population spikes. The results of the proposed experiments would provide baseline data for future studies using rats exposed to microgravic environments.

### 3. Specific questions

- (1) Does exposure to altered gravity (2G for 7 days) alter the effect of serotonin on population spike amplitude?
- (2) Does exposure to altered gravity (2G for 7 days) alter the effect of serotonin on transmembrane potential of hippocampal neurons?
- (3) Does exposure to altered gravity (2G for 7 days) alter the effect of serotonin on the input resistance of hippocampal pyramidal cells?

### 4. Rationale and experimental procedures

Experiments are designed to further characterize the response of hippocampal neurons to serotonin under differing gravitational conditions. The techniques used build on our current methods for recording extracellular activity (the population spike) in the slice preparation (Hooper et al., 1985, Appendix items A and B).

Tissue preparation. Hippocampal slices will be prepared in our laboratory following procedures we are currently using (Hooper et al., 1985; Appendices A and B). Male Sprague-Dawley rats are decapitated, and their brains quickly dissected and placed in chilled artificial cerebrospinal fluid (ACSF). The neocortex is removed to expose the bilateral hippocampi, which are removed. Slices 400-450 microns thick are cut perpendicular to the long axis of the hippocampus with a McIlwain tissue chopper. The slices are removed from the razor blade with a fine sable brush and placed in a petri dish containing chilled ACSF. The slices are then transferred with a large bore pipette to a plexiglass holding chamber where they rest at room temperature on ACSF-saturated filter paper. The holding chamber is provided with humidified 95% O<sub>2</sub>/5% CO<sub>2</sub> gas. The ACSF consists of (in mmoles): 124 NaCl, 5 KCl, 1.25 NaH<sub>2</sub>PO<sub>4</sub>, 2 MgSO<sub>4</sub>, 26 NaHCO<sub>3</sub>, 2 CaCl<sub>2</sub>, and 10 dextrose. At 37°C, the osmolarity is 305 ±5 mOsm and the pH is 7.4.

Recording chamber. We will record from slices in a recording chamber we have recently built. The chamber is a modification of the one described by Nicoll and Alger (1981) to include thermoelectric modules for controlling bath temperature. Approximately 1-2 hours after tissue dissection, a slice is transferred from the holding chamber to the constant-perfusion slice chamber. The slice is submerged in ACSF between two nets, the top net serving to hold the slice in place. The tissue is constantly perfused with oxygen-saturated ACSF at a rate of 1.5 to 2 ml/min via a gravity-fed reservoir system. The ACSF is removed by suction from a second chamber located behind and connected to the recording chamber. The entire chamber assembly is mounted in a machined

brass plate. The temperature of the slice is controlled by heating or cooling the brass plate with thermoelectric modules situated between the plate and an aluminum heat sink. Temperature is monitored via a calibrated thermistor placed in the recording chamber next to the slice.

Extracellular recording following Schaffer collateral stimulation. The methods will follow those used in our recent studies on population spikes (Hooper *et al.*, 1985; Appendix items A and B). Extracellular field potentials evoked by bipolar stimulation of CA1 stratum radiatum (the Schaffer collaterals) are recorded with glass microelectrodes (4-8 megohm, filled with 3M NaCl) placed in the pyramidal cell layer of CA1. Stimulus frequency is set to 0.033 Hz to avoid LTP, (except for bursts at frequencies above 10 Hz in selected experiments to induce LTP). The software for averaging extracellular fields has been developed for a Cromemco A/D converter and Zenith 100 microcomputer. The software enables us to average population spikes and determine mean spike amplitude.

Intracellular recording. While we have recorded intracellular potentials in a variety of tissues in past studies (Flaim *et al.*, 1977; Horowitz, *et al.*, 1969, 1971, 1980, 1982), this will be the first major study in which we will measure intracellular potentials from hippocampal pyramidal cells. (Our previous slice studies have involved extracellular recording.) Fortunately, pyramidal cells are relatively large cells, which should make them relatively easy to impale. Neurons in the pyramidal cell layer of area CA1 will be impaled with glass micropipettes driven with a Kopf hydraulic microdrive. The pipettes will be pulled on a Brown-Flaming horizontal puller and filled with 4.0M potassium acetate. Pipettes will then be connected to an oscilloscope and penwriter through a WPI high-impedance preamplifier. The preamplifier is equipped with a bridge circuit for passing a calibrated current through the micropipette. Potentials will be measured relative to a chlorided silver wire located in the suction chamber. A steady hyperpolarizing current is passed through the micropipette as cells are impaled to assist in sealing the cell and stabilizing the resting potential. Input resistance will be determined after a steady resting transmembrane potential is established by delivering 100 msec hyperpolarizing pulses. Neurons will be considered healthy if they have: (a) a resting membrane potential of at least 50 mV that is steady for 10 minutes, (b) action potentials elicited by short depolarizing pulses delivered through the pipette, (c) an input resistance of at least 30 Mohm. Data from cells meeting these criteria will be used in resolving the questions posed in this study.

Drug Application. Agonists will be applied to hippocampal neurons by microinjection, microiontophoresis, or superfusion. Antagonists will be applied by adding them to the perfusion line of the recording chamber. Thus the agonists serotonin or 8-OHDPAT will be injected directly into the recording chamber at concentrations of 300 nM-20  $\mu$ M and 200 nM-4  $\mu$ M, respectively, via a calibrated microsyringe (Hamilton). The drugs will be introduced into the recording chamber through a small hole near the ACSF inlet to assure consistent application. For microiontophoretic application (Horowitz *et al.*, 1980), glass micropipettes will be filled with serotonin or 8-OHDPAT. In some cases, double-barreled pipettes will be used, one barrel filled with serotonin and the other with 8-

OHDPAT. Pipettes will be placed next to the intracellular recording electrode in area CA1 via a micromanipulator. A Dagan Model 6400 multichannel current generator is used to control drug microiontophoresis. The antagonist methylsergide or buspirone will be mixed with ACSF to concentrations of 30  $\mu$ M and 3  $\mu$ M, respectively. Switching between media reservoirs (with and without antagonist) will be accomplished through the use of electric solenoid-actuated valves to minimize pressure pulses to the impaled neuron. Rapid changeover time will be achieved with a small recording chamber volume and minimal dead space in the media tubing (2/3 turnover time is approximately 1.5 minutes at 1 ml/min medium flow rate).

Protocols. EXPERIMENT 1. EXTRACELLULAR RECORDING (see Figure 2) is devoted to the development of extracellular techniques for measuring the response of hippocampal pyramidal cells to serotonin at 1G and will be performed in the first year of the grant. We will focus on experiments to characterize variables (e.g., concentration of serotonin) that modify cellular activity in order to select an operating range for reproducible, stable population spikes. The pyramidal cell response is dependent on Schaffer collateral stimulus variables (pulse amplitude, duration and repetition rate) and these will be varied to determine appropriate ranges for obtaining stable recording. We have already completed several studies characterizing the effects of temperature and pH on the system (Hooper et al., 1985, Appendix items A and B). All studies will be performed in the same time period to avoid any circadian variation in sensitivity to serotonin. Slices from each animal will be prepared as described above and placed in a holding chamber. Slices will then be transferred individually to the recording chamber and experiments performed such that all cells are characterized between 1000 and 1700 hours.

EXPERIMENT 2. INTRACELLULAR RECORDING (Figure 2) is a study also scheduled for the first year of the grant. This study will develop intracellular techniques. Hippocampal pyramidal cells in area CA1 will be impaled and transmembrane potential and resistance measured as described above. After the membrane potential and input resistance have stabilized (thus establishing baseline values), agonists will be applied via microsyringe to the bath or via a micropipette situated next to the intracellular electrode. The following response variables will be recorded: (a) maximum change in potential, (b) time to maximum potential, (c) time to return to baseline resting potential, (d) maximum change in input resistance, and (e) time course of resistance change. Responses will not be used unless the membrane potential returns to within 10% of baseline resting potential. After the neuron has returned to baseline conditions, the agonist will be applied again with an antagonist present in the bathing medium. Responses in the presence of antagonist will be compared with those obtained without antagonist present.

EXPERIMENT 3. EXTRACELLULAR RECORDING IN RATS EXPOSED TO 2G will be done in years 2 and 3. Population spike amplitudes will be measured (Methods, Appendix items A and B) for control rats and rats exposed to 2G fields. This will be a major experimental series because spike amplitudes reflect the overall activity of the system (the output for a




FIGURE 2 -- Outline of experiments.

volley of action potentials over the Schaffer collaterals). There is a 27% decrease in receptor number in rats exposed to 7 days to a 2G field. We will be using rats similarly exposed on the small animal centrifuge at the Chronic Acceleration Laboratory).

Experiments will be performed on two groups of animals. . An experimental group, consisting of 10 male Sprague-Dawley rats, will be exposed to a 2G hypergravic field for 7 days and a control group of 10 rats will be housed at 1G. Experiments will be performed at 1G within 5 hours after transferring rats from 2G to 1G. All animals will be fed Purina rat chow and water ad libitum and housed at 23-25°C. Animals from both groups will be identified by a code number unknown to the investigator making the measurements.

EXPERIMENT 4. EXTRACELLULAR RECORDING IN RATS EXPOSED TO 1.5, 2, 2.5, 3, 3.5 & 4G will be carried out in year 3 provided results obtained in EXPERIMENT 3 show changes in the amplitude of population spikes. Once data are obtained for rats exposed to a gravitational field of 2G for 7 days (for direct comparison with data on receptor number), the gravitational field and the exposure time will be varied. The gravitational field will be varied in 0.5G steps from 1 to 3G for 7 day exposure periods. The intent of this experiment is to determine if the modulatory effect of serotonin on electrical activity is a function of the gravitational field amplitude.

EXPERIMENT 5. EXTRACELLULAR RECORDING IN RATS EXPOSED TO 2G FOR LESS THAN 7 DAYS will be carried out in year 3 provided results obtained in EXPERIMENT 3 show changes in electrical activity. Holding the field at 2G, the period of exposure will be shortened to determine the minimum time required to detect effects on acclimation.

EXPERIMENT 6. INTRACELLULAR RECORDING IN RATS EXPOSED TO 2G will be done in years 2 and 3. This is a major experimental series comparing resistance and transmembrane potential changes in rats exposed to 2G and 1G.

(In developing techniques for the six experiments both hamsters and rats will be used.)

## 5. Data interpretation

Changes in the amplitude of population spikes will first be compared between animals exposed to 1G and to 2G for 7 days. A significant decrease in mean spike amplitude in rats exposed to 2G vs 1G controls would demonstrate that functional neuronal changes accompany the observed reduction in receptor number. The extent of these changes would be then determined as the gravitational field is increased and as the period of exposure is decreased.

Changes in transmembrane potential in response to serotonin or a 5-HT<sub>1</sub> agonist will be compared in hippocampal slices from rats exposed to a 2 G hypergravic field and control rats kept at 1 G. A significant change in rats exposed to hypergravity would lend support to the hypothesis that there is a direct affect of the gravitational field on hippocampal serotonin responsiveness. No significant change in serotonin

responsiveness in rats exposed to hypergravity would indicate that membrane (receptor) changes induced by gravity are not reflected in changes of electrical activity.

Changes in input resistance in response to serotonin or 5-HT<sub>1</sub> agonist will be compared in hippocampal slices from rats exposed to a 2 G hypergravic field and control rats kept at 1 G. Changes in input resistance will be compared with changes in transmembrane potential to determine if resistance and transmembrane potentials are correlated with gravitational fields. A decreased resistance (or an increased conductance) would indicate an increase in the number of open channels.

### **C. SIGNIFICANCE**

This proposal outlines neurobiological experiments on rats at earth gravity and in hypergravic fields. The series of experiments deal with serotonin receptors in the hippocampus. Receptor binding has been shown to be modified by microgravity and hypergravic environments. This study will focus on electrical activity mechanisms at the membrane level as a link to functional changes.

Core experiments will use animals exposed to 2 G fields for 7 days. The period of 7 days is chosen so that data obtained at 1 G and 2 G can be compared with recent Spacelab 3 experiments and with hypergravic experiments on serotonin receptors. Moreover, there is a great deal of data available on rats exposed 2 G fields.

The results of the proposed experiments serve as baseline data for future studies in hypogravic fields, including experiments on a space station. The studies involving the neurotransmitter serotonin will provide data on the effects of hypergravic fields and determine whether the number of binding sites is accompanied by changes in physiological function. If so, further experiments involving this transmitter could be performed on neural tissue after prescribed periods of exposure to a microgravity environment, with analysis of tissue samples performed shortly upon return of the animals to earth.

#### D. REFERENCES

- Andrade, R. and R. A. Nicoll. The novel anolytic buspirone elicits a small hyperpolarization and reduces serotonin responses at putative 5-HT<sub>1</sub> receptors on hippocampal CA1 pyramidal cells. Soc. for Neurosci. Abs. 11:597,1985.
- Andrade, R., R. C. Malenka and R. A. Nicoll. A G protein couples serotonin and GABA<sub>B</sub> receptors to the same channels in hippocampus. Science 234:1261-1265, 1986.
- Bliss, T. V. P., G. V. Goddard and MAI Riives. Reduction of long-term potentiation in the dentate gyrus of the rat following selective depletion of monoamines. J. Physiol. (Lond.) 334:475-491, 1983.
- Beck, S. G. and J. Goldfarb. Serotonin produces a reversible concentration dependent decrease of population spikes in rat hippocampal slices. Life Sci. 36:557-563, 1985.
- Beck, S. G., W. P. Clarke and J. Goldfarb. Spiperone differentiates multiple 5-hydroxytryptamine responses in rat hippocampal slices in vitro. Euro. J. Pharm. 116:195-197, 1985.
- Collingridge, G. L. and T. V. P. Bliss. NMDA receptors -- their role in long-term potentiation. Trends in Neurosci. 10:288-293.
- Collingridge G. L., S. K. Kehl and H. McLenna. Excitatory amino acids in synaptic transmission in the schaffer collateral-commissural pathway of the rat hippocampus. J. Physiol. (Lond.) 334:33046, 1983.
- Creese, I. and S. H. Snyder. <sup>3</sup>H-Spriprioperiod labes serotonin receptors in rat cerebral cortex and hippocampus. Eur. J. Pharmacol. 49:201-202, 1978.
- De Vivo, M. and S. Maayani. Characterization of the 5-Hydroxytrptamine<sub>1A</sub> receptor-mediated inhibition of forskolin-stimulated adenylate cyclase activity in guinea pig and rat hippocampal membranes. J. Pharmacol. and Exp. Ther. 238:248-253, 1986.
- Flaim, K. E., B. A. Horwitz and J. M. Horowitz. Coupling of signals to brown fat: alpha- and beta-adrenergic responses in intact rats. Am. J. Physiol. 232(2):R101-109, 1977.
- Geschwind, Irving I., John M. Horowitz, Gene M. Mikuckis and Roberg Dewey. Iontophoretic release of cyclic AMP and dispersion of melanosomes within a single melanophore. J. Cell Physiol. 74:928-939, 1977.
- Harris, E. W., A. H. Ganong and C. W. Cotman. Long-term potetntiation in the hippocampus involves activation of N-methyl-D-aspartate receptors. Brain Res. 323:132-137, 1984.



- Hooper, D. C., S. M. Martin and J. M. Horowitz. Temperature effects on evoked potentials of hippocampal slices from euthermic chipmunks, hamsters and rats. *Thermal Biol.* 10:35-40, 1985.
- Horowitz, J. M., J. Hamilton and B. A. Horowitz. Catecholamine-induced changes in transmembrane potential and temperature in normal and dystrophic hamster brown adipose tissue. *Life Sciences* 32:725-731, 1985.
- Horowitz, J. M., B. A. Horowitz and R. Em. Smith. Effect in vivo of norepinephrine on the membrane resistance of brown fat cells. *Experientia* 27:1419-1421, 1971.
- Horowitz, J. M., G. M. Mikuckis and M. A. Longshore. The response of single melanophores to extracellular and intracellular iontophoretic injection of melanocyte-stimulating hormone. *Endocrinology* 106:770-777, 1980.
- Horowitz, B. A., J. M. Horowitz and R. Em. Smith. Norepinephrine-induced depolarization of brown fat cells. *Proc. Nat. Acad. Sci.* 64:113-120, 1969.
- Henrik, J. The action of 5-Hydroxytryptamine on neuronal membranes and synaptic transmission in area CA1 of the hippocampus in vitro. *Brain Res.* 197:83-94, 1980.
- Madison, D. V. and R. A. Nicoll. Actions of noradrenaline recorded intracellularly in rat hippocampal CA1 pyramidal neurones, in vitro. *J. Physiol. (Lond.)* 372:221-244, 1986.
- Madison, D. V. and R. A. Nicoll. Cyclic adenosine 3',5'-monophosphate mediates  $\alpha$ -receptor actions of noradrenaline in rat hippocampal pyramidal cells. *J. Physiol. (Lond.)* 372:245-259, 1986.
- Mauk, M. D., P. J. Peroutka and J. D. Kocsis. Modulation of synaptic transmission in the hippocampus produced by novel anxiolytics (Buspirone, TVX-Q 7821 & DPAT. *Soc. Neurosci. Abstr.* 11:135, 1985]
- Miller, J. D., B. A. McMillen, D. M. Murakami, M. M. McConnaughey, H. L. Williams and C. A. Fuller. Effects of weightlessness on neurotransmitter receptors in selected brain areas. *The Physiologist* 28:S203-S204, 1985.
- Miller, J. D., B. McMillen and C. A. Fuller. Hypergravity effects on serotonin receptor binding in the rat center nervous system. (Abstract) Eighth Annual Meeting IUPS Commission on Gravitational Physiology, Tokyo, Japan, November 4-8, 1986.
- Nicoll, R. A. and B. E. Alger. A simple chamber for recording from submerged brain slices. *J. Neurosci. Meth.* 4:153-156, 1981.
- Peet, M. J., H. Gregersen and H. McLennan. 2-amino-5-phosphonovalerate and  $\text{CO}_2^{2+}$  selectively block depolarization and burst firing of rat hippocampal CA1 pyramidal neurons by N-methyl-D-aspartate. *Neuroscience* 17:635-641, 1986.

- Pedigo, N. W., H. I. Yamamura, I and D. L. Nelson. Discrimination of multiple [<sup>3</sup>]5-Hydroxytryptamine-binding sites by the neuroleptic spiperone in rat brain. *J. Neurochem.* 36:220-226, 1981.
- Peroutka, S. J. and S. H. Snider. Multiple serotonin receptors: differential binding of [<sup>3</sup>H]5-hydroxytryptamine, [<sup>3</sup>H]lysergic acid diethylamide and [<sup>3</sup>H]spiroperidol. *Mol. Pharmacol.* 16:687-699, 1979.
- Rowan, M. J. and R. Anwyl. The effect of prolonged treatment with tricyclic antidepressants on the actions of 5-hydroxytryptamide in the hippocampal slice of the rats. *Neuropharmacology* 24:131-137, 1985.
- Segal, M. The action of serotonin in the rat hippocampal slice preparation. *J. Physiol. (Lond.)* 303:423-439, 1980.
- Stanton, P. K. and J. M. Sarvey. Depletion of norepinephrine, but not serotonin, reduces long-term potentiation in the dentate gyrus of rat hippocampal slices. *J. Neuroscience* 5:2169-2176, 1985.
- Thomas, M. P., S. M. Martin and J. M. Horowitz. Temperature effects on evoked potentials of hippocampal slices from noncold-acclimated, cold-acclimated and hibernating hamsters. *J. Thermal Biol.* 11:213-218, 1986.

## E. PROGRESS REPORT (1985-1987)

**Description of research.** Over the past year we have shifted our experimental work almost completely to the hippocampus and away from studies on thermoregulation and the vestibular and auditory system. Our hippocampal experiments are related to a striking experimental finding from Spacelab-3, namely that there is an elevation in serotonin binding in rats exposed to microgravity for 7 days. This observation led us to focus primarily on developing techniques to study the hippocampus where the changes in receptor number have been reported. The goal of the experiments was to lay the basis for determining if there are functional changes in hippocampal neurons after exposure to hypergravic fields.

In addition to hippocampal studies, we completed experiments on vestibular, auditory and thermoregulatory systems that concluded a series of NASA supported projects on sensory mechanisms and regulatory systems.

Thus we finished one series of experiments and started another one on hippocampal slices. All focused on central neural function as modified by altered acceleration fields.

**Hippocampal studies.** Our major emphasis in recent experimental studies was centered on mastering the hippocampal slice preparation. A primary goal was to gain command of the techniques for recording electrical activity in the hippocampal slice preparation. We focused on thermal and pH effects on hippocampal electrical activity. With our shift in emphasis to the effects of neurotransmitters on cellular activity, a simulation study was completed on the effects of temperature on ion channel activity in hippocampal cells (Appendix C).

Experimental methods The hippocampus was sliced and placed in chilled artificial cerebrospinal fluid (ACSF). Slices 400-450 microns thick were cut perpendicular to the long axis of the hippocampus with a McIlwain tissue chopper and placed in a holding chamber. The chamber was provided with humidified 95% O<sub>2</sub> / 5% CO<sub>2</sub> gas. The ACSF consisted of (in mmoles): 124 NaCl, 5 KCl, 1.25 NaH<sub>2</sub>PO<sub>4</sub>, 2 MgSO<sub>4</sub>, 26 NaHCO<sub>3</sub>, 2 CaCl<sub>2</sub>, and 10 dextrose. At 37°C, the osmolarity was 305 ±5 mOsm and the pH was 7.4.

Approximately 1-2 hours after tissue dissection, a slice was transferred from the holding chamber to the submerged, constant-perfusion slice chamber. The slice was submerged in ACSF between two nets, the top net serving to hold the slice in place. The tissue was constantly perfused with oxygen-saturated ACSF at a rate of 1.5 to 2 ml/min via a gravity-fed reservoir system. The ACSF was removed by suction from a second chamber located behind and connected to the recording chamber. Temperature was monitored via a calibrated thermistor placed in the recording chamber next to the slice. Action potentials from a population of hippocampal pyramidal neurons were evoked by stimulating an afferent fiber tract, the Schaffer collaterals. The temperature and the pH of the ACSF bathing the slice were varied by controlling the temperature of a water chamber jacketing the recording chamber.

Experimental results. The effects of pH and temperature on hippocampal slices were studied in detail (references 5A, 1, 3, 4 and 6). Thresholds for evoked activity were significantly different in noncold-acclimated, cold-acclimated and hibernating hamsters, and may reflect acclimation of hippocampal neurons to cold. Plots of population spike amplitude (action potentials from a group of pyramidal cells) versus temperature have bell-shaped curves. The population spikes increased in amplitude as temperature was lowered from 35°C, reached a peak amplitude between 25 and 20°C, and then decreased until a response could not be evoked when temperature was further lowered. Techniques were thus developed for studying population spikes in the slice over a range of bath temperatures. Pilot studies on the effect of serotonin on population spikes were initiated. These techniques will be used in the experiments proposed in this grant renewal.

Simulation of the effect of temperature on ion channels in nerve membranes. We have developed a model to predict the effect of temperature on the electrical activity of a hippocampal pyramidal cell (5). With the development of brain slice techniques for voltage-clamping single cells, several types of ion channels in membranes of mammalian nerve cells have been identified and characterized in sufficient detail to allow a simulation of channel currents. Four populations of membrane channels in the pyramidal cell were simulated. Equations for current through these ion channels are similar to those first developed by Hodgkin and Huxley for sodium and potassium channels in the squid axon and more recently extended by Traub to include not only these channel types but, in addition, calcium and calcium-mediated potassium channels in hippocampal cells. Voltage and/or concentration dependent rate functions were used to describe the kinetic behavior of each population of channels. A temperature dependent term was included for each rate function to simulate the effect of changing temperature on neural activity. Model simulations correspond to experimental data over a range of temperature from 40°C to 35°C.

**Vestibular and auditory system.** Pulse angular acceleration evokes short latency far-field responses that can be attributed in large measure to the activation of the vestibular periphery and brain stem nuclei. The frequency spectrum of bone conducted vibrations coupled to the skull of rats during impulse angular acceleration stimulation was estimated to have greatest power at 2-3 kHz. The intensity of these vibrations was approximately 5 dB lower than the vibrations evoked by bone-conducted auditory clicks, which had their greatest power between 9 and 11 kHz. Moreover, the amplitudes of the first two major components of the response evoked by angular acceleration were greater than the first two major components of the response evoked by bone-conducted clicks. These results indicate that in rats, one can evoke far-field responses due to activity over the vestibular system and that this activity can be distinguished from auditory evoked responses (2, 9, 10).

Our data confirms a previous report that noninvasive procedures can be used to record brainstem vestibular evoked responses in rats. Particular attention was devoted to ruling out the possibility that what had been

called a vestibular response was in reality an auditory response. Stimulus characteristics were precisely determined. Using this type of stimulation, the amplitudes of the first two major components of the response evoked by angular acceleration were greater than the first two major components of the response evoked by bone-conducted clicks.

Thus, two independent laboratories have reported brainstem vestibular responses. These far-field potentials provide a means to record overall brainstem activity using noninvasive techniques. We have concluded experiments in this area and have manuscripts in preparation. (While we have turned our attention to the hippocampus, one of our former collaborators, Dr. T. Jones, is carrying on an independent series of studies on brainstem-evoked responses based, in part, on these studies.)

**Thermoregulatory studies** Previous studies in a variety of laboratories showed that the rat, dog, and monkey have an impaired ability to regulate their body temperature when exposed to hypergravic fields. One set of experiments using the rat was directed toward clarifying mechanisms underlying this impairment. The rat was chosen as an experimental animal because of studies at earth gravity, 1 G, that provided basic background for further studies both at zero-G and in hypergravic fields from 1.5 to 4 G.

One series of completed experiments (7) showed that rats acclimated to a gravitational field of 2.1 G are able to regulate their core temperature better when cold stressed at 2.1 G than are rats acclimated at earth gravity (1 G). Rats acclimated to 2.1 G also increased tail temperature ( $T_t$ ) and decreased core temperature during exposure to 5.8 G. Thus, rats acclimated to 2.1 G were not able to regulate their temperature when exposed to higher gravitational fields. Acclimation did not result in a change in thermoregulatory ability at 1 G. It appears that rats acclimated to 1 G conditions continue to regulate their core temperature, albeit at a lower core temperature, when placed in a hypergravic field of 2.1 G.

Another series of completed experiments (2, 8) compared groups of rats acclimated to gravitational fields of 1 or 2.1 G. That is, one group was born and raised at 2.1 G and belonged to the 12th generation of rats living continuously on a centrifuge in a 2.1 G field, except for brief periods of routine care at 1 G. The finding that rats acclimated at 2.1 G could thermoregulate better than rats raised at 1 G when cold stressed at 2.1 G shows that acclimation to a hypergravic environment can modify the activity of a neural control system in mammals. However, rats acclimated to 2.1 G were not able to regulate their core temperature when first exposed to 5.8 G indicating that acclimation at one level of a hypergravic field does not improve the ability of the animal to thermoregulate at higher field levels.

(While we have concluded the experimental series on temperature regulation in rats exposed to hypergravic fields, one of our early collaborators, Dr. C. Fuller, is carrying on related hypergravic studies on temperature regulation and circadian rhythms.)

## References

### Abstracts

- 1A. Hoffman, L., Jones T. A., Fullerton, B. and Horowitz, J. Hypothermia-Induced changes in interpeak latency of brainstem auditory evoked responses recorded from hamsters and rats. APS Fall Meeting, 1985.
- 2A. Horowitz, J., S. Patterson and C. Monson. Oxygen consumption during cold exposure at 2.1 g in rats adapted to hypergravic fields. APS Fall Meeting, 1985.
- 3A. Hoffman, L. and J. Horowitz. Can impulse angular acceleration activate the vestibular and auditory systems in rats? Abstract. ASGSB Annual Meeting, Charlottesville, Va., 1986.
- 4A. Horowitz, J., M. Lindley and M. Thomas. A model of hippocampal pyramidal cells and the effects of temperature. FASEB, 1986.
- 5A. Thomas M. P., P. J. Eckerman and J. M. Horowitz. Temperature and pH effects on evoked responses from hamster hippocampal slices. Abstract. Society for Neuroscience, 1986.
- 6A. Hamill, N. J., J. M. Horowitz and M. D. McGinn. Peripheral and brainstem auditory responses in squirrels arousing from hibernation. FASEB, 1987.

### Papers

1. Hooper, D. C., S. M. Marin, and J. M. Horowitz. Temperature effects on evoked potentials of hippocampal slices from euthermic chipmunks, Hamsters and rats. J. of Therm. Biol. 10:35-40, 1985.
2. Horowitz, J., S. Patterson, and C. Monson. Oxygen consumption during cold exposure at 2.1 G in rats adapted to hypogravic fields. Physiologist 28:S313-S142, 1985.
3. Thomas, M. P., S. M. Martin, and J. M. Horowitz. Temperature effects on evoked potentials of hippocampal slices from noncold-acclimated, cold-acclimated and hibernating hamsters. J. Therm. Biol. 11:213-218, 1986. (Included in this grant proposal as Appendix B)
4. Horowitz, J. M., M. P. Thomas, and P. Eckerman. Thermal dependence of neural activity in the hamster hippocampal slice preparation. J. Therm. Biol. In press.
5. Lindley, M. A., M. P. Thomas, and J. M. Horowitz. A simulation of the effects of temperature on hippocampal neurons. Computers in Biomed. In press. (Included in this grant proposal as Appendix C).
6. Eckerman, P., K. Scharruhn, and J. M. Horowitz. Effects of

temperature and pH on hippocampal population spikes. In preparation. (Included in this grant proposal as Appendix A).

7. Monson, C. B., J. M. Horowitz and B. A. Horowitz. Core temperature is regulated, although at a lower temperature, in rats exposed to hypergravic fields.
8. Monson, C. B., S. L. Patterson, J. M. Horowitz. Thermoregulation in hypergravity-acclimated rats. In preparation.
9. Hoffman, L. and J. M. Horowitz. Peripheral and brainstem vestibular responses in rats: The effect of masking. In preparation.
10. Hoffman, L. and J. M. Horowitz. Peripheral and Brainstem vestibular responses in rats: The effect of hypothermia. In preparation.

## F. BIOGRAPHICAL SKETCH OF PIs

### CURRICULUM VITA John Horowitz

BIRTHDATE: [REDACTED]

NATIONALITY: US citizen

EDUCATION:	Institution	Degree	Year Conferred	Scientific Field
	Univ. of Calif., Berkeley	B.S.	1959	Electrical Engineering
	Univ. of Calif., Berkeley	M.S.	1961	Electrical Engineering
	Univ. of Calif., Berkeley	Ph.D.	1968	Biophysics

#### MEMBERSHIP IN PROFESSIONAL SOCIETY

Sigma Xi  
American Physiological Society  
Society for Neuroscience

#### MAJOR RESEARCH INTERESTS:

Neural regulation of temperature, neural networks

#### RESEARCH AND/OR PROFESSIONAL EXPERIENCE

1978-present	Professor of Physiology and Research in the Experiment Station (UC Davis)	Physiologist
1973-1978	Associate Professor of Physiology and Research Physiologist in the Experiment Station (UC Davis)	
1968-1973	Assistant Professor of Physiology (UC Davis)	
1963-1968	Teaching and Research Assistantships (UC Berkeley)	
1961-1963	Graduate Research Engineer (UC Berkeley)	
1959-1961	Research Assistant (UC Berkeley)	

#### PUBLICATIONS -- ABSTRACTS:

1. Horowitz, J. M. and W. J. Freeman. Evoked potentials arising from neural population elements excited at different times on a warped surface. *The Physiologist* 8:197, 1965.
2. Horowitz, B. A., P. A. Herd, J. M. Horowitz and R. E. Smith. Brown fat: Effect of norepinephrine on intracellular potentials and  $\text{Na}^+/\text{K}^+$  ATPase activity. *The Physiologist* 12:257, 1969.
3. Horowitz, J. M. and R. Em. Smith. Neural control of thermogenesis in brown adipose tissue. *Fed. Proc.* 29:659, 1970.



4. Horowitz, J. M. and R. Em. Smith. Effect of norepinephrine on membrane conductance of brown fat cells. Fed. Proc. 30:320, 1971.
5. Scobey, R. and J. M. Horowitz. The response of monkey retinal ganglion cells to the flashing and moving luminous spot. Society for Neuroscience -- First Annual Meeting, October 1971.
6. Horowitz, J. M. and R. Em. Smith. Response of a hippocampal network with a background level of excitation. Fed. Proc., 1972.
7. Mackay, R. S., G. Eilers, J. M. Horowitz and E. Marg. Ultrasonic echo imaging in eye research. Digest of the 1962 15th Annual Conference in Medicine and Biology, Paper 9, 1962.
8. Horowitz, J. M., R. Karem and M. Saleh. Response of a hippocampal network to changes in cutaneous temperature. Fed. Proc. 32:407, 1973.
9. Horowitz, J. M. and C. A. Fuller. Central neural signals and the regulation of nonshivering thermogenesis. International Symposium on Depressed Metabolism and Cold Thermogenesis, Prague, October 9-12, 1974.
10. Horowitz, J. M. and C. A. Fuller. Concurrent neural control of shivering and nonshivering thermogenesis. XXVI International Congress of Physiological Sciences - Satellite Conference, Israel, 1974.
11. Fuller, C. A., J. M. Horowitz and B. A. Horowitz. Evaluation of neural models controlling heat production in cold-exposed animals. Biomedical Engineering Symposium Abstracts, p. 10, UCD, 1974.
12. Smietana, P. J., J. M. Horowitz and T. J. Willey. CSMP programs for dispersion of signals over a fiber tract and for coupling between a fiber tract and a neural network. Biomedical Engineering Symposium Abstracts, p. 11, UCD, 1974.
13. Horowitz, J. M. and J. Mates. Correlation of hippocampal pyramidal cell activity with the theta rhythm and with dispersion over a feedback branch of a neural network. Biomedical Engineering Symposium Abstracts, p. 12, UCD, 1974.
14. Thompson, A. C., J. M. Horowitz and I. H. Wagman. Rates of radiant thermal stimulation and threshold levels of cutaneous nociception in man. Fed. Proc. 35:768, 1976.
15. Horowitz, B. A. and J. M. Horowitz. Thermoregulatory responses of unanesthetized rats exposed to 1 to 4G. Fed. Proc. 36:579, 1977.
16. Horowitz, J. M. and I. I. Geschwind. Melanosome dispersion within a single melanophore following iontophoretic release of cyclic AMP. Fed. Proc. 36:284, 1977.

17. Horowitz, J. M. and R. E. Plant. Simulation of energy conversion in brown adipose tissue using network thermodynamics: I. Interaction of cellular components. Biomedical Engineering Symposium Abstracts, p. 2, UCD, 1977.
18. Plant, R. E. and J. M. Horowitz. Simulation of energy conversion of brown adipose tissue using network thermodynamics: II. Coupling between chemical reactions and ion transport. Biomedical Engineering Symposium Abstracts, p. 3, UCD, 1977.
19. Giacchino, J., J. M. Horowitz and B. Horwitz. Simulation of the thermoregulatory system of rats exposed concurrently to cold and to gravitational fields of 1 to 4G. Biomedical Engineering Symposium Abstracts, p. 28, UCD, 1977.
20. Giacchino, J., J. M. Horowitz and B. A. Horwitz. Simulation of the thermoregulatory system of rats exposed to hypergravic fields and to low ambient temperatures. Fed. Proc. 37:426, 1978.
21. Giacchino, J., J. M. Horowitz and B. A. Horwitz. Thermoregulation of unanesthetized rats during and after exposure to hypergravitational fields. Fed. Proc. 38:1053, 1979.
22. Horowitz, J. M. and M. A. Longshore. Comparison of hippocampal evoked activity in vivo and in vitro. Fed. Proc. 38:898, 1979.
23. Longshore, M., J. M. Horowitz and G. M. Mikuckis. Extracellular and intracellular injection of melanocyte stimulating hormone and the response of single melanophores in frog skin. The Physiologist 22:79, 1979.
24. Horowitz, J. M., E. Schertel and B. A. Horwitz. Centrifuge high-g effects on temperature regulation in unanesthetized rats. The Physiologist 22:58, 1979.
25. Longshore, M. A., J. M. Horowitz and H. L. Nguyen. Localization and characterization of adrenergic plasma membrane receptors on frog skin melanophores. Fed. Proc. 39:732, 1980.
26. Giacchino, J. L., J. M. Horowitz and E. R. Schertel. Effect of p-chlorophenylalanine on thermoregulation of unrestrained rats. Fed. Proc. 39:990, 1980.
27. Horowitz, J. M., J. L. Giacchino and K. S. Kott. Signal dispersion within a hippocampal neural network in the rabbit and rat. Society for Neuroscience, Abstracts 6:564, 1980.
28. Wickler, S. J., B. A. Horwitz and J. M. Horowitz. Metabolic organization of muscle and brown fat in normal and dystrophic hamsters. Fed. Proc. 40:518, 1981.
29. van Kan, P. L. E., C. Morgan, J. Horowitz and B. A. Horwitz. Na<sup>+</sup>-dependent glucose transport in dystrophic and normal hamster ileum. Fed. Proc. 40:347, 1981.

30. Rothfeld, J., K. Kott, J. Horowitz and B. A. Horowitz. The in vitro and in vivo effects of norepinephrine on the membrane potential of hamster skeletal muscle. Fed. Proc. 40:614, 1981.
31. Giacchino, J. L. and J. M. Horowitz. Modulation of background activity in a hippocampal neural network. Society for Neuroscience, Abstracts 7:451, 1981.
32. Jones, T. A., L. Hoffman and J. M. Horowitz. Altered auditory function in rats exposed to hypergravic fields. The Physiologist 24:213, 1982.
33. Horowitz, J. M., J. Hamilton and B. A. Horowitz. Norepinephrine-induced transmembrane potential changes in normal and dystrophic hamster brown adipocytes. Fed. Proc. 41:1613, 1982.
34. Horowitz, J. M., C. B. Monson and B. A. Horowitz. Effects of restraint hypothermia on heat production in rats exposed to hypergravic fields. The Physiologist 24:212, 1982.
35. Jones, T. A. and J. M. Horowitz. The effects of interstimulus noise on short-latency auditory responses. Society for Neuroscience, Abstracts, 8:975, 1982.
36. Horowitz, J. M., S. J. Wickler and B. A. Horowitz. Blood flow in normal and dystrophic hamsters during nonshivering thermogenesis. Fed. Proc. 42:1323, 1983.
37. Monson, C. B., J. M. Horowitz and B. A. Horowitz. Evidence for parallel controllers in rats exposed to hypergravic fields. Fed. Proc. 42:1124. 1983.
38. Hamilton, J., J. M. Horowitz and B. A. Horowitz. Norepinephrine-induced depolarization of hamster skeletal muscle in vitro. Fed. Proc. 42:995, 1983.
39. Horowitz, J. M. and B. A. Horowitz. Thermoregulation in cold- and noncold-acclimated rats cold-exposed while in hypergravic fields. XXIX International Congress, International Union of Physiological Sciences, Australia, 1983.
40. Hoffman, L. and J. Horowitz. Wide band noise masks brainstem responses evoked by clicks more effectively than responses evoked by angular acceleration in rats. Soc. for Neuroscience, Abstracts, 10:112, 1984.
41. Monson, C. B., J. M. Horowitz and J. Oyama. Thermoregulation during cold exposure at 2.1 g in rats adapted to hypergravic fields. Fed. Proc. 43:907, 1984.
42. Hoffman, L. F. and J. M. Horowitz. Far-field brainstem responses evoked by vestibular and auditory stimuli exhibit increases in interpeak latency as brain temperature is decreased. Sixth Annual Meeting IUPS Commission on Gravitational Physiology, Switzerland, 1984.

43. Monson, C. B., S. Patterson, J. Oyama and J. M. Horowitz. Cold tolerance of rats adapted to hypergravity. Fed. Proc. 44(3):623, 1985.
44. Martin, S. M., M. Thomas, K. Scharruhn and J. M. Horowitz. Temperature effects on evoked potentials of hippocampal slices from euthermic, cold-acclimated and hibernating hamsters. Fed. Proc. 44:1045, 1985.
45. Horowitz, J., S. Patterson and C. Monson. Oxygen consumption during cold exposure at 2.1 g in rats adapted to hypergravic fields. American Physiological Society, New York, 1985.
46. Hoffman, L., T. A. Jones, B. Fullerton and J. Horowitz. Hypothermia-induced changes in interpeak latency of brainstem auditory evoked responses recorded from hamsters and rats. American Physiological Society, New York, 1985.
47. Horowitz, J. M., M. P. Thomas and P. J. Eckerman. Hippocampal activity in hibernating and nonhibernating animals. Living in the Cold, an international symposium, Fall Leaf Lake Conference Center, California, Oct 6-11, 1985.
48. Hoffman, L. and J. Horowitz. Can impulse angular acceleration activate both the vestibular and auditory systems in rats? ASGSB Annual Meeting, Charlottesville, Va., 1986.
49. Horowitz, J., M. Lindley and M. Thomas. A model of hippocampal pyramidal cells and the effects of temperature. Fed. Proc. 45:408, 1986.
50. Thomas, M. P., P. J. Eckerman and J. M. Horowitz. Temperature and pH effects on evoked responses from hippocampal slices. Soc. for Neuroscience, Abstracts, 16th Annual Meeting, 12:1529, 1986.
51. Hamill, N. J., J. M. Horowitz and M. D. McGinn. Peripheral and brainstem auditory responses in squirrels arousing from hibernation. Fed. Proc. 46:839, 1987.
52. Thomas, M. P. and J. M. Horowitz. Simulation of thermal effects on four membrane channels in hippocampal pyramidal cells. Soc. for Neuro., 1987.
53. Horowitz, J. M. and M. P. Thomas. Serotonergic mechanisms and the hippocampal slice preparation. ASGSB, Utah State Univ., Logan, Utah, 1987.

PUBLICATIONS -- FULL PAPERS:

1. Everhart, T. E. and J. Horowitz. Limitations of space-time harmonics for microwave amplification. Electronic Research Laboratory, University of California Series No. 60, Issue No. 440, 1962.

2. Horowitz, J. M. and W. F. Freeman. Evoked potentials arising from neural population elements excited at different times on a warped surface. *Bulletin of Math Biophysics* 28:519-535, 1966.
3. Horowitz, B. A., J. M. Horowitz and R. Em. Smith. Norepinephrine-induced depolarization of brown fat cells. *Proc. Nat. Acad. Sci.* 64:113-120, 1969.
4. Horowitz, J. M., B. A. Horowitz and R. Em. Smith. Control of brown fat thermogenesis: A systems approach. *J. de Physiologie* 63:273-276, 1971.
5. Horowitz, J. M., B. A. Horowitz and R. Em. Smith. Effect in vivo of norepinephrine on the membrane resistance of brown fat cells. *Experientia* 27:1419-1421, 1971.
6. Horowitz, J. M. Evoked activity of single units and neural populations in the hippocampus of the cat. *Electroenceph. Clin. Neurophys.* 32:227-240, 1971.
7. Walters, R. F., G. M. Nishimoto, J. M. Horowitz and R. E. Burger. Initial development of individualized instruction with computer support. *Proceedings of the 1972 Conference on Computers in Undergraduate Curricula*, pp. 199-207, 1972.
8. Horowitz, J. M. Neural control of thermogenesis in brown adipose tissue. *Fed. Proc.*, pp. 115-121, *Fed. of the Amer. Soc. of Exper. Biol.*, 1972.
9. Scobey, R. and J. M. Horowitz. The detection of small image displacements by cat retinal ganglion cells. *Vision Res.* 12:2133-2143, 1972.
10. Horowitz, J. M., W. J. Freeman and P. J. Stoll. A neural network with a background level of excitation in the cat hippocampus. *Internat. J. Neurosci.* 5:113-123, 1973.
11. Horowitz, J. M. and L. K. Erskine. Central regulation of temperature in cold environments. I. A dynamic model with two temperature inputs. *Computers and Biomed. Res.* 6:57-73, 1973.
12. Fuller, C. A., J. M. Horowitz and B. A. Horowitz. Central regulation of temperature in cold environments. II. A dynamic model with three temperature inputs. *Computers and Biomed. Res.* 7:164-182, 1974.
13. Horowitz, J. M., M. A. Saleh and R. D. Karem. Correlation of hippocampal theta rhythm with changes in cutaneous temperature. *Am. J. Physiol.* 227(3):635-642, 1974.
14. Teskey, N., B. Horowitz and J. Horowitz. Norepinephrine-induced depolarization of skeletal muscle cells. *Eur. J. Pharmacol.* 30:352-355, 1975.

15. Martin, S. M., G. P. Moberg and J. M. Horowitz. Glucocorticoids and the hippocampal theta rhythm in loosely restrained, unanesthetized rabbits. *Brain Res.* 93:535-542, 1975.
16. Horowitz, J. M., P. J. Smietana, Jr. and T. J. Willey. Dispersion along fiber tracts and in the coupling between tracts and a cortical network. *Computer Prog. Biomed.* 4:253-262, 1975.
17. Fuller, C. A., J. M. Horowitz and B. A. Horowitz. Sorting of signals from thermosensitive areas. *Computer Prog. Biomed.* 4(4):263-273, 1975.
18. Horowitz, J. M. Neural models on temperature regulation for cold-stressed animals. In: Temperature Regulation and Drug Action, (P. Lomax, E. Schonbaum and J. Jacob, Editors), Karger:New York, pp. 1-10, 1975.
19. Fuller, C. A., B. A. Horowitz and J. M. Horowitz. Shivering and nonshivering thermogenic responses of cold-exposed rats to hypothalamic warming. *Am. J. Physiol.* 228:1519-1524, 1975.
20. Horowitz, J. M. and J. W. B. Mates. Signal dispersion within a hippocampal neural network. *Computers Biol. Med.* 5:283-296, 1975.
21. Scobey, R. P. and J. M. Horowitz. Detection of image displacement by phasic cells in peripheral visual fields of the monkey. *Vis. Res.* 16:15-24, 1976.
22. Horowitz, J. M., C. A. Fuller and B. A. Horowitz. Central neural pathways in the control of nonshivering thermogenesis. In: Regulation of Depressed Metabolism and Thermogenesis, (L. Jansky and X. J. Musacchia, Editors), Chapter 1, pp. 3-25, Charles C. Thomas Publisher:Springfield, Illinois, 1976.
23. Mates, J. and J. M. Horowitz. Instability in a hippocampal neural network. *Computer Prog. Biomed.* 6:74-84, 1976.
24. Horowitz, J. M., R. G. Tearce and B. Mulloney. Comparison of electrical activity and behavioral correlates of neural networks in the cat and locust. *Computer Prog. Biomed.* 6:92-100, 1976.
25. Mates, J. W. B. and J. M. Horowitz. Interacting neural networks on a cortical surface. In: Proceeding 1975 Winter Simulation Conference, Society for Computer Simulation, La Jolla, CA, pp. 187-193, 1976.
26. Flaim, K. E., J. M. Horowitz and B. A. Horowitz. Functional and anatomical characteristics of nerve-brown adipose interaction in the rat. *Pflugers Archives* 365:9-14, 1976.
27. Horowitz, J. M. Concurrent neural control of shivering and nonshivering thermogenesis. *Israel J. Med. Sci.* 12:1082-1085, 1976.

28. Horowitz, J. M., B. A. Horowitz and J. W. B. Mates. Thermoregulatory models and their relationship to fever and neural transmitters. *Biometerology* 6(Part II):137-149, 1976.
29. Fuller, C. A., J. M. Horowitz and B. A. Horowitz. Effects of acceleration on the thermoregulatory responses of unanesthetized rats. *J. Appl. Physiol.* 42:74-79, 1977.
30. Fuller, C. A., J. M. Horowitz and B. A. Horowitz. Spinal cord thermosensitivity and sorting of neural signals in cold-exposed rats. *J. Appl. Physiol.* 42:154-158, 1977.
31. Horowitz, J. M. Norepinephrine and thermoregulatory networks. In: Drugs, Biogenic Amines and Body Temperature, (Copper, Lomax and Schonbaum, Editors), pp. 1-4, 1977.
32. Flaim, K. E., B. A. Horowitz and J. M. Horowitz. Coupling of signals to brown fat: alpha- and beta-adrenergic responses in intact rats. *Am. J. Physiol.* 232(2):R101-109, 1977.
33. Horowitz, B. A. and J. M. Horowitz. Role of the plasma membrane in brown fat thermogenesis. *Experientia* 33:1126-1128, 1977.
34. Geschwind, Irving I., John M. Horowitz, Gene M. Mikuckis and Roberg Dewey. Iontophoretic release of cyclic AMP and dispersion of melanosomes within a single melanophore. *J. Cell Physiol.* 74:928-939, 1977.
35. Horowitz, J. M. and B. A. Horowitz. Thermoregulatory responses of unanesthetized rats exposed to gravitational fields of 1 to 4g. In: COSPAR: Life Sciences and Space Research XVI, (R. Holmquist and A. C. Strickland, Editors), Pergamon Press Oxford and New York, pp. 77-82, 1978.
36. Horowitz, J. M. and B. A. Horowitz. An overview of neural models postulated for thermoregulation. In: Body Temperature: Regulation, Drug Effects and Therapeutic Indications, (P. Lomax and E. Schonbaum, Editors), Dekker:New York, pp. 25-37, 1978.
37. Horowitz, J. M. and R. E. Plant. Controlled cellular energy conversion in brown adipose tissue thermogenesis. *Am. J. Physiol.* 253:R121-R129, 1978.
38. Plant, R. E. and J. M. Horowitz. Simulation of coupling between chemical reactions and ion transport in brown adipose tissue using network thermodynamics. *Computer Prog. Biomed.* 8:171-179, 1978.
39. Giacchino, J., J. M. Horowitz and B. A. Horowitz. Simulation of the thermoregulatory system of rats exposed concurrently to cold and to acceleration fields. *Computers Biol. Med.* 9:205-211, 1979.
40. Plant, R. E. and J. M. Horowitz. Energy conversion in biological systems. I. Chemical reactions and ion transport. *Journal of the Franklin Institute* 308:269-280, 1979.

41. Horowitz, J. M., J. L. Giacchino and B. A. Horowitz. Energy conversion in biological systems. II. Heat production in brown adipose tissue. *Journal of the Franklin Institute* 308:281-295, 1979.
42. Giacchino, J., J. M. Horowitz and B. A. Horowitz. Thermoregulation in unrestrained rats during and after exposure to 1.5 to 4G. *J. Appl. Physiol.* 46(6):1049-1053, 1979.
43. Horowitz, J. M., E. R. Schertel and B. A. Horowitz. Centrifuge high-g effects on temperature regulation in unanesthetized rats. *The Physiologist* 22:S57-S58, 1979.
44. Horowitz, J. M., G. M. Mikuckis and M. A. Longshore. The response of single melanophores to extracellular and intracellular iontophoretic injection of melanocyte-stimulating hormone. *Endocrinology* 106:770-777, 1980.
45. Teskey, N. J., J. M. Horowitz and B. A. Horowitz. Norepinephrine-induced membrane responses in normal and dystrophic hamster muscle. *Proc. Soc. Exp. Biol. Med.* 164:331-335, 1980.
46. Ross, D., J. M. Horowitz and R. E. Plant. Oscillatory neural networks in the rabbit hippocampus. *Biol. Cybernetics* 37:115-124, 1980.
47. Schertel, E. R., J. M. Horowitz and B. A. Horowitz. Effects of gravitational profiles on the rat's thermoregulatory response to cold. *J. Appl. Physiol.: Respir. Environ. Exer. Physiol.* 49:663-668, 1980.
48. Horowitz, J. M., B. A. Horowitz and J. Oyama. Alterations in heat loss and heat production mechanisms in rats exposed to hypergravic fields. *The Physiologist* 23:S119-S120, 1980.
49. Longshore, M. A. and J. M. Horowitz. Localization and characterization of adrenergic receptors on frog skin melanophores. *Am. J. Physiol.* E84-E89, 1981.
50. Horowitz, J. M., K. Whitelaw, A. Lim and C. Jacobs. Real-time analysis of hippocampal neural activity in the intervals between interictal spikes. *Computer Prog. in Biomed.* 13:19-26, 1981.
51. Monson, C. B. and J. M. Horowitz. Simulation of thermoregulation in serotonin-depleted rats subjected to acceleration fields. *Computer Prog. in Biomed.* 13:9-17, 1981.
52. Jones, T. A. and J. M. Horowitz. Core temperature and brainstem auditory evoked potentials as complimentary noninvasive measures of central neural function during exposure to hypergravic fields. *The Physiologist* 24:S107-S108, 1981.
53. Monson, C. B., J. M. Horowitz and B. A. Horowitz. Restraint hypothermia in cold-exposed rats at 3G and 1G. *The Physiologist* 25:S89-S90, 1982.



54. Jones, T. A., L. Hoffman and J. M. Horowitz. Altered auditory function in rats exposed to hypergravic fields. *The Physiologist* 25:S93-S94, 1982.
55. Jones, T. A. and J. M. Horowitz. Bone-conducted auditory stimulation in unrestrained, unanesthetized animals. *J. Neuroscience Methods* 7:261-267, 1983.
56. Horowitz, J. M., J. Hamilton and B. A. Horowitz. Catecholamine-induced changes in transmembrane potential and temperature in normal and dystrophic hamster brown adipose tissue. *Life Sciences* 32:725-731, 1983.
57. Giacchino, J. L., E. R. Schertel, J. M. Horowitz and B. A. Horowitz. Effect of p-chlorophenylalanine on thermoregulation in unrestrained rats. *Am. J. Physiol.* 244:R299-R302, 1983.
58. Monson, C. B., J. M. Horowitz and B. A. Horowitz. Hypergravic fields and parallel controllers for thermoregulation. *J. Appl. Physiol.: Respirat. Environ. Exercise Physiol.* 55:990-995, 1983.
59. Willey, T. J., G. Maeda, R. L. Schultz, W. S. Seibly and J. M. Horowitz. The principal projection pathway between the olfactory bulb and the prepyriform cortex in the cat. *J. Neurosci. Res.* 9:253-277, 1983.
60. Horowitz, J., B. A. Horowitz and C. B. Monson. Thermoregulation in cold- and noncold-acclimated rats cold exposed in hypergravic fields. *The Physiologist* 26:S169-S172, 1983.
61. Monson, C. B. and J. M. Horowitz. Impairment of thermogenesis and heat conservation in rats during 3 hours of 3G exposure. *J. Aviat. Space and Environ. Med.* 55:542-545, 1984.
62. Horowitz, J. and B. Horowitz. Changes in the number and characteristics of brown adipocytes underlies the impaired thermogenic response of dystrophic hamsters. In: *Thermal Physiology* (J. R. S. Hales, editor). Raven Press: New York, pp. 179-182, 1984.
63. Hoffman, L. F. and J. M. Horowitz. Far-field brainstem responses evoked by vestibular and auditory stimuli exhibit increases in interpeak latency as brain temperature is decreased. *The Physiologist* 27:S89-S90, 1984.
64. Mlayj, W., R. E. Smith and J. M. Horowitz. New directions in scientific computing: Impact of advances in microprocessor achitecture and system design. *Comp. Prog. in Biomed.* 18:149-162, 1984.
65. Hooper, D. C., S. M. Martin and J. M. Horowitz. Temperature effects on evoked potentials of hippocampal slices from euthermic chipmunks, hamster and rats. *J. Thermal Biol.* 10:35-40, 1985.

66. Horowitz, J., S. Patterson and C. Monson. Oxygen consumption during cold exposure at 2.1G in rats adapted to hypergravic fields. *The Physiologist* 28:S141-S142, 1985.
67. Horowitz, J. M., J. Hamilton and B. Horwitz. Norepinephrine depolarizes sartorius but hyperpolarizes soleus muscle in vitro. *Eur. J. Pharmacol.* 119:85-92, 1985.
68. Thomas, M. P., S. M. Martin, and J. M. Horowitz. Temperature effects on evoked potentials of hippocampal slices from noncold-acclimated, cold-acclimated and hibernating hamsters. *J. Thermal Biol.* 11:213-218, 1986.
69. Smith A. H., E. M. Bernauer, A. L. Black, R. E. Burger, J. H. Crowe, J. M. Horowitz, G. P. Moberg and E. M. Renkin. History of physiology at University of California, Davis. *The Physiologist* 29:46-57, 1986.
70. Horowitz, J. M., M. Thomas and P. Eckerman. Thermal dependence of neural activity in the hamster hippocampal slice preparation. *J. Thermal Biol.* in press. 1987.
71. Lindley, M. A., M. P. Thomas, and J. M. Horowitz. A simulation of the effects of temperature on hippocampal neurons. *Comp. Prog. in Biomed.* in press. 1987.

CURRICULUM VITA

Barbara Horwitz

BIRTHDATE: 9/26/40

NATIONALITY: US citizen

EDUCATION:

Institution	Degree	Year Conferred	Scientific Field
University of Florida, Gainesville	B.S.	1961	Biology
University of Florida, Gainesville	M.S.	1962	Biology
Emory University, Atlanta, Georgia	Ph.D.	1966	Physiology
Marine Biol. Lab, Woods Hole, Mass.	—	—	Physiology
University of California, Los Angeles & Davis	Postdoc.	1966-1968	Physiology

THESIS WORK:

Master of Science

"Temperature Effects on Oxygen Uptake of Liver and Kidney Tissues of a Hibernating and a Non-hibernating Mammal"

Doctor of Philosophy

"The Effect of Temperature on the Oxidative Metabolism and Morphology of Liver Mitochondria from a Hibernating and a Non-hibernating Mammal"

HONORS AND MEMBERSHIP IN PROFESSIONAL SOCIETIES:

Phi Beta Kappa

Phi Kappa Phi

Sigma Xi

Phi Sigma

Alpha Epsilon Delta

Alpha Lambda Delta

Outstanding undergraduate in Biological Sciences  
(Univ. of Florida, 1960-61),

"Who's Who in American Colleges and Universities"  
(1965 - representing Emory Grad. School)

Member of: American Physiological Society  
American Institute of Nutrition  
New York Academy of Sciences  
American Society of Zoologists

MAJOR RESEARCH INTERESTS:

Regulation of Cellular Metabolism  
Environmental Physiology  
Temperature Regulation/Cellular Thermogenesis  
Catecholamine-receptor Interaction  
Cellular Adaptation  
Obesity

## RESEARCH AND/OR PROFESSIONAL EXPERIENCE

1978-present Professor of Physiology (UC Davis)

1975-1978 Associate Professor of Physiology (UC Davis)

1972-1975 Assistant Professor of Physiology (UC Davis)

1968-1972 Assistant Research Physiologist (UC Davis)

1966-1968 USPHS Postdoctoral Fellow (UCLA, UC Davis)

1963-1964 USPHS trainee in Fertilization and Gamete Physiology (Marine Biology Lab, Woods Hole, Mass.)

1963-1966 USPHS trainee in Physiology (Emory University)

1961-1962 USPHS Predoctoral Fellow (University of Florida)

1959-1961 Teaching/Research Assistant — Biology Department (University of Florida)

PUBLICATIONS — ABSTRACTS:

1. Horwitz, B. A. and L. Nelson. Myokinase activity in sea urchin fragments. Biol. Bull. 125:366, 1963.
2. Horwitz, B. A. and L. Nelson. Adenylate kinase and ATPase activity in spisula and Asterias eggs. Biol. Bull. 127:374, 1964.
3. Horwitz, B. A. and L. Nelson. Rates of oxygen consumption of Arbacia, Asterias, and Spisula eggs. Biol. Bull. 127:374, 1964.
4. Horwitz, B. A., L. Nelson and R. E. Smith. Differential temperature sensitivities of rates of respiration and phosphorylation in liver mitochondria from cold-adapted rats. Fed. Proc. 26, 1967.
5. Horwitz, B. A., R. E. Smith and E. T. Pengelley. Estimated thermal contribution of brown fat during arousal of the ground squirrel, C. lateralis. The Physiologist 10:206, 1967.
6. Horwitz, B. A., P. A. Herd and R. E. Smith. Electron transport-coupled phosphorylation in brown adipose tissue. Fed. Proc. 27:633, 1968.
7. Herd, P. A. and B. A. Horwitz. Factors controlling brown fat mitochondrial respiration. Fed. Proc. 28:721, 1969.
8. Horwitz, B. A., P. A. Herd, J. M. Horowitz, Jr. and R. E. Smith. Brown fat: Effect of norepinephrine on intracellular potentials and  $\text{Na}^+/\text{K}^+$  ATPase activity. The Physiologist 12:257, 1969.
9. Herd, P. A. and B. A. Horwitz. Brown fat mitochondria: Coupled endogenous respiration. Fed. Proc. 29:659, 1970.
10. Horwitz, B. A. and J. F. Detrick. Role of brown fat in the norepinephrine-induced thermogenic response of cold-acclimated rats. Fed. Proc. 30:319, 1971.
11. Horwitz, B. A. Role of active sodium transport in brown fat thermogenesis. Proc. International Symposium on Temperature Regulation, Jerusalem, Israel, Oct., 1974.
12. Horwitz, B. A. Effects of alpha- and beta-adrenergic agonists on responses of brown adipocytes. Fed. Proc. 34:477, 1975.
13. Jacobs, J. R. and B. A. Horwitz. Characteristics of a lipolytic extract from rabbit urine. Fed. Proc. 35:518, 1976.
14. Leung, P. M. B. and B. A. Horwitz. Feeding patterns of rats in response to changes in energy demand. Fed. Proc. 35:520, 1976.
15. Horwitz, B. A. and J. M. Horowitz. Thermoregulatory responses of unanesthetized rats exposed to 1 to 4G. Fed. Proc. 36:579, 1977.
16. Hamilton, J. and B. A. Horwitz. Brown fat lipolysis: Effects of adrenergic agonists and cyclic nucleotides. Fed. Proc. 36:579, 1977.
17. Giacchino, J., J. M. Horowitz and B. A. Horwitz. Simulation of the thermoregulatory system of rats exposed to hypergravic fields and to low ambient temperatures. Fed. Proc. 37:426, 1978.

18. Giacchino, J., J. M. Horowitz and B. A. Horwitz. Thermoregulation of anesthetized rats during and after exposure to hypergravitational fields. Fed. Proc. 38:1053, 1979.
19. O'Donnell, M. E. and B. A. Horwitz. Effect of an uncoupler on ouabain-sensitive norepinephrine-induced  $\text{VO}_2$  in brown adipocytes. Fed. Proc. 38:1055, 1979.
20. Horowitz, J. M., E. Schertel, and B. A. Horwitz. Centrifuge high-g effects on temperature regulation in unanesthetized rats. The Physiologist 22:58, 1979.
21. O'Donnell and B. A. Horwitz. Effects of butyrate and oligomycin on ouabain sensitive norepinephrine (NE)-induced  $\text{VO}_2$  in brown adipocytes. Fed. Proc. 39:1182, 1980.
22. Hettinger, D. R. and B. A. Horwitz. Catecholamine stimulation of the  $(\text{Na}^+/\text{K}^+)\text{-ATPase}$  in brown adipose tissue. Fed. Proc. 39:979, 1980.
23. van Kan, P. L. E., C. Morgan, J. Horowitz and B. A. Horwitz.  $\text{Na}^+$ -dependent glucose transport in dystrophic and normal hamster ileum. Fed. Proc. 40:374, 1981.
24. Rothfeld, J., K. Kott, J. Horowitz and B. A. Horwitz. The in vitro and in vivo effects of norepinephrine on the membrane potential of hamster skeletal muscle. Fed. Proc. 40:614, 1981.
25. Cornett, L. E., B. A. Horwitz and J. M. Roberts. Beta-adrenergic receptors in brown fat from dystrophic and normal hamsters. Fed. Proc. 40:254, 1981.
26. Wickler, S. J., B. A. Horwitz and J. M. Horowitz. Metabolic organization of muscle and brown fat in normal and dystrophic hamsters. Fed. Proc. 40:518, 1981.
27. Horwitz, B. A. Diverse approaches to teaching environmental physiology. Fed. Proc. 40:377, 1981.
28. Horwitz, B. A. and M. Mooney. Effects of a  $\text{Na}^+$  ionophore on the norepinephrine-stimulated  $(\text{Na}^+/\text{K}^+)\text{-ATPase}$  in brown adipose tissue. Fed. Proc. 40:612, 1981.
29. Wickler, S. J., B. A. Horwitz and J. S. Stern. Catecholamine-stimulated blood flow is impaired to brown adipose tissue in genetically obese rats. Fed. Proc. 41:389, 1982.
30. Horwitz, B. A., J. Finch and M. Roche. Calcium involvement in MSH-evoked melanosome dispersion. Fed. Proc. 41:1594, 1982.
31. Horowitz, J. M., J. Hamilton and B. A. Horwitz. Norepinephrine-induced transmembrane potential changes in normal and dystrophic hamster brown adipocytes. Fed. Proc. 41:1613, 1982.
32. Kott, K. S., S. J. Wickler and B. A. Horwitz. The effect of photoperiod and temperature on brown adipose tissue. The Physiologist 25:244, 1982.
33. Wickler, S., K. Kott and B. A. Horwitz. Hibernation: A natural analog to exercise inactivity? The Physiologist 25:244, 1982.
34. Horowitz, J. M., C. B. Monson and B. A. Horwitz. Effects of restraint on two modes of heat production of cold-exposed rats in hypergravic fields. The

35. Stern, J. S., S. J. Wickler and B. A. Horwitz. Blood flow is impaired to brown adipose tissue in Zucker obese rats. J. Nutr. 112:16, 1982.
36. Monson, C. B., J. M. Horowitz and B. A. Horwitz. Evidence for parallel thermocontrollers in rats exposed to hypergravic fields. Fed. Proc. 42:1124, 1983.
37. Hamilton, J., J. M. Horowitz and B. A. Horwitz. Norepinephrine-induced depolarization of hamster skeletal muscle in vitro. Fed. Proc. 42:995, 1983.
38. Horowitz, J. M., S. J. Wickler and B. A. Horwitz. Blood flow in normal and dystrophic hamsters during nonshivering thermogenesis (NST). Fed. Proc. 42:1323, 1983.
39. Arrendell, W. B. and B. A. Horwitz. Inhibition of brown fat thermogenesis and lipolysis by chlorpromazine and trifluoperazine, two calmodulin inhibitors. Fed. Proc. 42:1323, 1983.
40. Wickler, S. J., Z. Glick, J. S. Stern and B. A. Horwitz. Blood flow into brown fat and into other tissues following a single meal. Fed. Proc. 42:1190, 1983.
41. Horwitz, B. A., T. Inokuchi, S. Wickler and J. S. Stern. Removal of brown fat from obese and lean Zucker rats. Fed. Proc. 42:1188, 1983.
42. Wickler, S. J., Z. Glick, J. S. Stern and B. A. Horwitz. Blood flow into brown fat is greater after a high carbohydrate than after a high fat test meal. Fed. Proc. 43:1062, 1984.
43. Moore, B. J., B. A. Horwitz and J. S. Stern. Brown fat lipectomy decreases food intake and increases adiposity in cold-exposed rats. Fed. Proc. 43:852, 1984.
44. Kott, K. S. and B. A. Horwitz. The effect of photoperiod and castration on brown adipose tissue (BAT). Fed. Proc. 43:797, 1984.
45. Moore, B. J., J. E. Reynolds, B. A. Horwitz and J. S. Stern. Brown adipose tissue (BAT) mediates the increased energy expenditure of rat pups raised in small litters. Fed. Proc. 44:1678, 1985.
46. Gerardo, T., B. J. Moore, J. S. Stern and B. A. Horwitz. Prolactin depresses brown fat thermogenesis while increasing food intake and white fat depots. Fed. Proc. 44:1160, 1985.
47. Horwitz, B., J. Hamilton and K. Kott. Brown fat mitochondria from hibernating hamsters exhibit reduced GDP binding. Fed. Proc. 44:1161, 1985.
48. Moore, B. J., P. J. Armbruster, B. A. Horwitz and J. S. Stern. Energy expenditure is reduced in 2-day old preobese Zucker fatty rats. North Am. Assoc. for Study of Obesity meeting, Univ. Wash., 1984.
49. Stern, J. S., B. J. Moore and B. A. Horwitz. Impaired thermogenesis and its role in the development of obesity. Appetite, 6:171, 1985.
50. Moore, B. J., B. A. Horwitz, and J. S. Stern. Rates of oxygen consumption ( $VO_2$ ) at thermoneutrality and in response to cold in newborn male Zucker rats. In: Regulation of Energy Expenditure; ed. E. S. Horton and E. Danforth, Jr. London: Libby p. A-13, 1985.

51. Moore, B. J., J. S. Stern, and B. A. Horwitz. Diet alters thermogenic response to acute cold exposure in the neonatal rat. "Living in the Cold" Symposium, Fallen Leaf Lake, Ca., 1985.
52. Moore, B. J., T. Gerardo-Gettens, B. A. Horwitz, and J. S. Stern. Hyperprolactinemia: possible stimulatory effects on food intake. Presented at "Mechanisms of Appetite and Obesity", San Antonio, Tx., October 1985.
53. Moore, B. J., T. Inokuchi, J. S. Stern, and B. A. Horwitz. Brown adipose tissue lipectomy leads to decreased food intake and increased fat deposition in the Osborne-Mendel rat. Appetite 6:179, 1985.
54. Gerardo, T., B. J. Moore, J. S. Stern, and B. A. Horwitz. Hyperprolactinemia decreases GDP binding in brown fat mitochondria. "Living in the Cold" Symposium, Fallen Leaf Lake, Ca., 1985.
55. Wickler, S. J., J. S. Stern, and B. A. Horwitz. Oxygen consumption and brown adipose tissue (BAT) blood flow are normalized in adrenalectomized Zucker obese (fa/fa) rats. Fed. Proc. 45:482, 1986.
56. Gerardo-Gettens, T., B. J. Moore, J. S. Stern, and B. A. Horwitz. Ovariectomy does not alter the suppression of brown fat (BAT) GDP binding in the lactating rat. Am. J. Clin. Nutr. 43:6, A-14, 1986.
57. Moore, B. J., T. Inokuchi, B. A. Horwitz, and J. S. Stern. Brown fat metabolism is significantly reduced in nonlactating rat dams. Am. J. Clin. Nutr. 43:6, A-16, 1986.
58. Moore, B. J., T. Inokuchi, B. A. Horwitz, and J. S. Stern. Pregnancy followed by lactation or no lactation leads to increased fat cell number and increased risk of obesity in the rat. Am. J. Clin. Nutr. 43:6, A-15, 1986.
59. Moore, B. J., T. Inokuchi, B. A. Horwitz, and J. S. Stern. Recovery of brown fat metabolism is complete after one cycle of pregnancy and lactation in the rat. Fifth International Congress on Obesity, Jerusalem, 1986.
60. Moore, B. J., T. Inokuchi, B. A. Horwitz, and J. S. Stern. Factors associated with reproduction lead to increased fat cell number and predispose rat dams to obesity. Fifth International Congress on Obesity, Jerusalem, 1986.
61. Gerardo-Gettens, T., B. J. Moore, J. S. Stern, and B. A. Horwitz. Exogenous prolactin administration stimulates food intake in a dose-dependent manner. IXth International Conference on the Physiology of Food & Fluid Intake, Seattle, WA, 1986.
62. Fournier, L., N. Cortez, B. J. Moore, K.S. Kott, T. W. Castonguay, B. A. Horwitz, and J. S. Stern. Is 12:12 photoperiod appropriate for studies of food intake? IXth International Conference on the Physiology of Food & Fluid Intake, Seattle, WA, 1986.
63. McDonald, R., J. S. Stern, and B. A. Horwitz. Effects of exercise on dietary induced thermogenesis in rats. Fed. Proc. 45:973, 1986.
64. McDonald, R., B. Horwitz, and J. S. Stern. Brown adipose tissue and cold exposure in the older rat. The Gerontologist 26:154A, 1986.



65. Fournier, L.M., N. M. Cortez, B. J. Moore, K. S. Kott, T. W. Castonguay, B. A. Horwitz, and J. S. Stern. Rats are sensitive to photoperiod and pinealectomy. Fifth International Congress on Obesity, Jerusalem, 1986.
66. Gerardo-Gettens, T., B. J. Moore, J. S. Stern, and B. A. Horwitz. Is the prolactin stimulation of food intake progesterone dependent? Fifth International Congress on Obesity, Jerusalem, 1986.
67. McDonald, R. B., B. Horwitz, and J. S. Stern. Thermic effect of a meal following exercise in the rat. Am. J. Clin. Nutr. 43:6, A-13, 1986.
68. McDonald, R. B., J. S. Stern, and B. A. Horwitz. Exercise prevents the age-related decrease in core temperature of cold-exposed F344 rats. Fed. Proc. 46:1089, 1987.
69. Gong, T. W., B. A. Horwitz, and J. S. Stern. Adrenalectomy, high fat feeding and thermogenesis in obese and lean Zucker rats. Fed. Proc. 46:1487, 1987.
70. McDonald, R. B., J. S. Stern, and B. A. Horwitz. Effects of aging and exercise on fat cell number and size. Medicine and Science in Sports and Exercise 19: S75, 1987.
71. Gerardo-Gettens, T., B. J. Moore, J. S. Stern, and B. A. Horwitz. Prolactin stimulates food intake in the absence of ovarian progesterone. Fed. Proc. 46:1338, 1987.

PUBLICATIONS -- FULL PAPERS

1. Horwitz, B. A. Temperature effects on oxygen uptake of liver and kidney tissue of a hibernating and non-hibernating mammal. Physiol. Zool. 37:231-239, 1964.
2. Horwitz, B. A. Rates of oxygen consumption of fertilized and unfertilized Asterias, Arbacia and Spisula eggs. Expt. Cell. Res. 38:620-625, 1965.
3. Horwitz, B. A., L. Nelson and V. Popovic. Effect of temperature on oxidative phosphorylation in hibernators and non-hibernators. J. Appl. Physiol. 22:639-644, 1967.
4. Horwitz, B. A., R. E. Smith and E. T. Pengelley. Estimated heat contribution of brown fat in arousing ground squirrels (C. lateralis). Am. J. Physiol. 214:115-121, 1968.
5. Horwitz, B. A. and L. Nelson. Effect of temperature on mitochondrial respiration in a hibernator (Myotis austroriparius) and a non-hibernator (Rattus rattus). Comp. Biochem. Physiol. 24:385-394, 1968.
6. Imai, Y., B. A. Horwitz and R. E. Smith. Calorigenesis of brown adipose tissue in cold-exposed rats. Proc. Exp. Biol. Med. 127:717-719, 1968.
7. Smith, R. E., B. A. Horwitz and Y. Imai. Quantitative thermogenesis of brown fat in hibernation and cold adaptation. In: Quantitative Biology of Metabolism, ed. by A. Locker and F. Kruger. Springer-Verlag:New York, 1968, pp. 106-112.
8. Horwitz, B. A., P. A. Herd and R. E. Smith. Effect of norepinephrine and uncoupling agents on brown adipose tissue. Can. J. Physiol. Pharmacol. 46:897-902, 1968.
9. Smith, R. E. and B. A. Horwitz. Brown fat thermogenesis. Physiol. Rev. 49:330-425, 1969.
10. Horwitz, B. A., J. M. Horowitz, Jr. and R. E. Smith. Norepinephrine-induced depolarization of brown fat cells. Proc. Nat. Acad. Sci. 64:113-121, 1969.
11. Horwitz, B. A., P. A. Herd and R. E. Smith. Bioenergetics of brown adipose tissue. Lipids 5:30-34, 1970.
12. Herd, P. A., B. A. Horwitz and R. E. Smith. Norepinephrine-sensitive  $\text{Na}^+/\text{K}^+$  ATPase activity in brown adipose tissue. Experientia 26:825-826, 1970.
13. Horwitz, B. A. Brown fat thermogenesis: Physiological control and metabolic basis. In: Nonshivering Thermogenesis, ed. L. Jansky. Academia: Prague, 1971, pp. 221-240.
14. Horowitz, J. M., B. A. Horwitz and R. Em. Smith. Control of brown fat thermogenesis: A systems approach. J. de Physiol. (Paris) 63:273-276, 1971.
15. Horowitz, J. M., B. A. Horwitz and R. Em. Smith. Effect of norepinephrine on membrane resistance of brown fat cells. Experientia 27:1419-1421, 1971.
16. Horwitz, B. A., J. F. Detrick and R. Em. Smith. Norepinephrine-induced thermogenesis in cold-acclimated rats: Role of brown adipose tissue. Experientia 28:284-286, 1972.

17. Horwitz, B. A. and R. Em. Smith. Function and control of brown fat thermogenesis during cold exposure. In: Proceedings of International Symposium on Environmental Physiology: Bioenergetics, ed. R. E. Smith, J. P. Hannon, J. L. Shields and B. A. Horwitz. FASEB:Washington, 1972, pp. 134-140.
18. Horwitz, B. A. Ouabain-sensitive component of brown fat thermogenesis. Am. J. Physiol. 224:352-355, 1973.
19. Fuller, C. A., J. M. Horowitz and B. A. Horwitz. Central regulation of temperature in cold environments — II. A dynamic model with three temperature inputs. Computers and Biomed. Res. 8:164-182, 1974.
20. Horwitz, B. A. Physiological and biochemical characteristics of adrenergic receptors and pathways in brown adipocytes. In: Temperature Regulation and Drug Action, ed. P. Lomax, E. Schonbaum and J. Jacob. Karger:New York, 1975, pp. 150-158.
21. Horwitz, B. A. and G. E. Hanes. Isoproterenol-induced calorigenesis of dystrophic and normal hamsters. Proc. Soc. Expt. Biol. & Med. 147:392-395, 1974.
22. Fuller, C. A., B. A. Horwitz and J. M. Horowitz. Shivering and nonshivering thermogenic responses of cold-exposed rats to hypothalamic warming. Am. J. Physiol. 228:1519-1524, 1975.
23. Leung, P. M. B. and B. A. Horwitz. Free-feeding patterns of rats: Effects of pyrogen and dietary protein content. Am. J. Physiol. 228:1284-1287, 1975.
24. Fuller, C. A., J. M. Horowitz and B. A. Horwitz. Sorting of signals from thermosensitive areas. Computer Programs in Biomedicine 4:264-274, 1975.
25. Teskey, N., B. Horwitz and J. Horowitz. Norepinephrine-induced depolarization of skeletal muscle cells. Europ. J. Pharmacol. 30:352-355, 1975.
26. Horwitz, B. A. and M. Eaton. The effect of adrenergic agonists and cyclic AMP on the  $\text{Na}^+/\text{K}^+$  ATPase activity of brown adipose tissue. Europ. J. Pharmacol. 34:241-247, 1975.
27. Horwitz, B. A. The effect of cold exposure on liver mitochondrial and peroxisomal distribution in the rat, hamster and bat. Comp. Biochem. and Physiol. 54A:45-48, 1976.
28. Horwitz, B. A. and G. E. Hanes. Propranolol and pyrogen effects on shivering and nonshivering thermogenesis. Am. J. Physiol. 230:637-642, 1976.
29. Horowitz, J. M., C. A. Fuller and B. A. Horwitz. Central neural pathways and the control of nonshivering thermogenesis. In: Regulation of Depressed Metabolism and Thermogenesis, ed. L. Jansky and X. J. Mussachia. C. C. Thomas: Springfield, 1976, pp. 3-25.
30. Horwitz, B. A. Role of active sodium transport in brown fat thermogenesis. Israel J. Med. Sci. 12:1086-1089, 1976.
31. Horowitz, J. M., B. A. Horwitz and J. W. B. Mates. Thermoregulatory models and their relationship to fever and neural transmitters. Biometeorology 6 (Part II):137-149, 1976.

32. Leung, P. M. B. and B. A. Horwitz. Free-feeding patterns of rats in response to changes in environmental temperature. Am. J. Physiol. 231:1220-1224, 1976.
33. Flaim, L. K., J. M. Horowitz and B. A. Horwitz. Functional and anatomical characteristics of the nerve-brown adipose tissue interaction in the rat. Pflugers Arch. 365:9-14, 1976.
34. Fuller, C. A., J. M. Horowitz and B. A. Horwitz. Effects of acceleration on the thermoregulatory responses of unanesthetized rats. J. Appl. Physiol. 42:74-79, 1977.
35. Horwitz, B. A. and M. Eaton. Ouabain-sensitive liver and diaphragm respiration in cold-acclimated hamster. J. Appl. Physiol. 42:150-154, 1977.
36. Fuller, C. A., J. M. Horowitz and B. A. Horwitz. Spinal cord thermosensitivity and sorting of neural signals in cold-exposed rats. J. Appl. Physiol. 42:154-158, 1977.
37. Horwitz, B. A. Adrenergic-receptor involvement in brown adipose tissue activation. In: Drugs, Biogenic Amines and Body Temperature, ed. K. E. Cooper, P. Lomax and E. Schonbaum. Karger:New York, 1977, pp. 160-166.
38. Flaim, L. K., B. A. Horwitz and J. M. Horowitz. Coupling of signals to brown fat: alpha- and beta-adrenergic responses in intact rats. Am. J. Physiol. 232:R101-R109, 1977.
39. Horwitz, B. A. and J. M. Horowitz. Role of the plasma membrane in brown fat thermogenesis. Experientia 33:1126-1128, 1977.
40. Horwitz, B. A. Pathways underlying nonshivering thermogenesis in peripheral tissues. In: Depressed Metabolism and Cold Thermogenesis, ed. L. Jansky. Charles University:Prague, 1975, pp. 127-131.
41. Horowitz, J. M. and B. A. Horwitz. Thermoregulatory responses of unanesthetized rats exposed to gravitational fields of 1 to 4 G. In: COSPAR: Life Sciences and Space Research. XVI, ed. R. Holmquist and A. C. Strickland. Pergamon Press:New York, 1978, pp. 77-82.
42. Horowitz, J. M. and B. A. Horwitz. An overview of neural models postulated for thermoregulation. In: Body Temperature: Regulation, Drug Effects and Therapeutic Indications, ed. P. Lomax and E. Schonbaum. Marcel Dekker:New York, 1978, pp. 25-37.
43. Horwitz, B. A. Neurohumoral regulation of nonshivering thermogenesis in mammals. In: Strategies in Cold: Natural Torpidity and Thermogenesis, ed. L. Wang and J. Hudson. Academic Press:New York, 1978, pp. 619-653.
44. Horwitz, B. A. Plasma membrane involvement in brown fat thermogenesis. In: Effectors of Thermogenesis, ed. L. Girardier and . Seydoux. Verlag:Basel, 1978, pp. 19-23.
45. Hamilton, J. and B. A. Horwitz. Adrenergic- and cyclic nucleotide-induced glycerol release from brown adipocytes. Europ. J. Pharmacol. 56:1-5, 1979.
46. Giacchino, J., B. A. Horwitz and J. M. Horowitz. Temperature regulation in unrestrained rats during and after exposure to gravitational fields of 1.5 to 4G. J. Appl. Physiol. 46:1049-1053, 1979.

47. Horwitz, B. A. Metabolic aspects of thermogenesis: Neuronal and hormonal control. Introduction. Fed. Proc. 38:2147-2149, 1979.
48. Horwitz, B. A. Cellular events underlying catecholamine-induced thermogenesis: Cation transport in brown adipocytes. Fed. Proc. 38:2170-2176, 1979.
49. Horowitz, J. M., J. L. Giacchino and B. A. Horwitz. Energy conversion in biological systems. II. Heat production in brown adipose tissue. J. of the Franklin Institute 308:281-295, 1979.
50. Horowitz, J. M., J. Giacchino and B. A. Horwitz. Simulation of the thermoregulatory system of rats exposed concurrently to cold and to acceleration fields. Comput. Biol. Med. 9:205-211, 1979.
51. Horwitz, B. A. and D. R. Hettinger. Cold-induced enzymatic adjustments in ectotherms and homeotherms. In: Comparative Mechanisms of Cold Adaptation, ed. L. Underwood. Academic Press:New York, 1979, pp. 91-127.
52. Horowitz, J. M., E. R. Schertel and B. A. Horwitz. Centrifuge high-g effects on temperature regulation in unanesthetized rats. The Physiologist 22:S57-58, 1979.
53. Teskey, N. J., J. M. Horowitz and B. A. Horwitz. Norepinephrine-induced membrane responses in normal and dystrophic hamster skeletal muscle. Proc. Soc. Expt. Biol. Med. 164:331-335, 1980.
54. Schertel, E. R., J. M. Horowitz and B. A. Horwitz. Effects of gravitational profiles on the rat's thermoregulatory response to cold. J. Appl. Physiol.: Respirat. Environ. Exercise Physiol. 49:663-668, 1980.
55. Horowitz, J. M., B. A. Horwitz and J. Oyama. Alterations in heat loss and heat production mechanisms in rats exposed to hypergravic fields. The Physiologist 23:S119-120, 1980.
56. O'Donnell, M. E. and B. A. Horwitz. Effects of butyrate on ouabain-sensitive respiration of hamster brown adipocytes. Am. J. Physiol. 242:C46-51, 1982.
57. Milam, K. M., J. S. Stern and B. A. Horwitz. Isoproterenol alters nonshivering thermogenesis in the Zucker obese rat (fafa). Pharmacol. Biochem. and Behav. 16:627-630, 1982.
58. Wickler, S. J., B. A. Horwitz and J. S. Stern. Regional blood flow in genetically obese rats during nonshivering thermogenesis. International J. Obesity 6:481-490, 1982.
59. Monson, C. B., J. M. Horowitz and B. A. Horwitz. Restraint hypothermia in cold-exposed rats at 3 G and at 1 G. The Physiologist 25:S89-90, 1982.
60. Giacchino, J. L., E. R. Schertel, J. M. Horowitz and B. A. Horwitz. Effect of p-chlorophenylalanine on thermoregulation in unrestrained rats. Am. J. Physiol. 244:R299-302, 1983.
61. Wickler, S. J. and B. A. Horwitz. Metabolic organization of muscle and brown adipose tissue of normal and dystrophic hamsters. Am. J. Physiol. 244:R407-412, 1983.
62. Hettinger, D. R. and B. A. Horwitz. Kinetic characteristics of hamster (Mesocricetus auratus) brown fat ( $\text{Na}^+/\text{K}^+$ )-ATPase: Effects of catecholamines.

63. Kott, K. and B. A. Horwitz. Photoperiod and pinealectomy do not affect cold-induced deposition of brown adipose tissue in the Long-Evans rat. Cryobiology 20:100-105, 1983.
64. Horowitz, J. M., J. Hamilton and B. A. Horwitz. Catecholamine-induced changes in transmembrane potential and temperature in normal and dystrophic hamster brown adipose tissue. Life Sciences 32:725-731, 1983.
65. Horwitz, B. A. and S. J. Wickler. Diminished respiratory responses of brown adipocytes isolated from BIO 14.6 dystrophic hamsters. Proc. Soc. Expt. Biol. Med. 173:35-40, 1983.
66. Monson, C. B., J. M. Horowitz and B. A. Horwitz. Hypergravic fields and parallel controllers for thermoregulation. J. Appl. Physiol.:Respirat., Environ. Exercise Physiol. 55:990-995, 1983.
67. Guttas, J., A. Carter and B. A. Horwitz. Plasma membrane protection against the acute effects of inorganic lead on the respiratory rates of intact liver cells. J. Toxicol. Environ. Hlth. 12:731-736, 1983.
68. Horowitz, J. M., B. A. Horwitz and C. B. Monson. Thermoregulation in cold- and noncold-acclimated rats cold exposed in hypergravic fields. The Physiologist 26:S169-172, 1983.
69. Horwitz, B. A. Cold acclimation of hamsters leads to functional alterations in  $K^+$  transport across the plasma membrane of liver cells. In: Thermal Physiology, ed. by J. R. S. Hales. Raven Press:New York, 1984, pp. 463-466.
70. Horwitz, B. A., T. Inokuchi, S. Wickler and J. S. Stern. Lipoprotein lipase activity and cellularity in brown and white adipose tissue in Zucker obese rats. Metabolism 33:354-357, 1984.
71. Horwitz, B. A., S. D. Clark, J. Hamilton and J. Guttas. Effect of cold acclimation on  $Na^+/K^+$  transport in hamster liver cells. J. Comp. Physiol. (B) 154:159-166, 1984.
72. Horwitz, B. A. and J. Hamilton. Alpha-adrenergic induced changes in hamster (Mesocricetus) brown adipocyte respiration and membrane potential. J. Comp. Physiol. Biochem. 78C:99-104, 1984.
73. Horowitz, J. and B. Horwitz. Changes in the number and characteristics of brown adipocytes underlie the impaired thermogenic response of dystrophic hamsters. In: Thermal Physiology, ed. by J. R. S. Hales. Raven Press:New York, 1984, pp. 179-182.
74. Wickler, S. and B. A. Horwitz. Blood flow in normal and dystrophic hamsters during nonshivering thermogenesis. Am. J. Physiol. 247:R189-195, 1984.
75. Wickler, S. J., B. A. Horwitz, S. F. Flaim and K. F. LaNoue. Isoproterenol-induced blood flow in rats acclimated to room temperature and cold. Am. J. Physiol. 246:R747-752, 1984.
76. Glick, Z., S. J. Wickler, J. S. Stern and B. A. Horwitz. Regional blood flow in rats after a single low protein, high carbohydrate test meal. Am. J. Physiol. 247:R160-166, 1984.

77. Stern, J. S., T. Inokuchi, T. W. Castonguay, S. J. Wickler and B. A. Horwitz. Scapular brown fat removal enhances development of adiposity in cold-exposed obese Zucker rats. Am. J. Physiol. 247:R918-R926, 1984.
78. Glick, Z., S. J. Wickler, J. S. Stern and B. A. Horwitz. Blood flow into brown fat is greater after a high carbohydrate than after a high fat test meal. J. Nutrit. 114:1934-1939, 1984.
79. Horwitz, B. A., T. Inokuchi, B. Moore and J. S. Stern. The effect of brown fat removal on the development of obesity in Zucker and Osborne-Mendel rats. In: Regulation of Energy Expenditure, Int. J. Obesity 9, Suppl. 2:43-48, 1985.
80. Moore, B. J., T. Inokuchi, J. S. Stern and B. A. Horwitz. Brown adipose tissue lipectomy leads to increased fat deposition in the Osborne Mendel rat. Am. J. Physiol. 248:R231-235, 1985.
81. Moore, B. J., S. J. Armbruster, B. A. Horwitz and J. S. Stern. Energy expenditure is reduced in preobese 2-day old Zucker (fa/fa) fatty rats. Am. J. Physiol. 249:R262-R265, 1985.
82. Moore, B. J., B. A. Horwitz and J. S. Stern. Brown fat thermogenesis and its role in the development of obesity. Brain Res. Bull. 14:577-583, 1985.
83. Horwitz, B. A., J. S. Hamilton, and K. S. Kott. GDP binding to hamster brown fat mitochondria is reduced during hibernation. Am. J. Physiol. 249:R689-693, 1985.
84. Horowitz, J. M., J. S. Hamilton, and B. A. Horwitz. Norepinephrine depolarizes sartorius but hyperpolarizes soleus muscles in vitro. Eur. J. Pharmacol. 119:85-92, 1985.
85. Freedman, M. R., B. A. Horwitz, and J. S. Stern. Effect of adrenalectomy and glucocorticoid replacement on development of obesity in the genetically obese Zucker rat. Am. J. Physiol. 250:R595-607, 1986.
86. Moore, B. J., T. Gerardo-Gettens, B. A. Horwitz, and J. S. Stern. Hyperprolactinemia stimulates food intake in the female rat. Brain Res. Bull., 17:563-569, 1986
87. Horwitz, B. A., K. S. Kott, J. S. Hamilton, and B. J. Moore. Regulation of brown fat thermogenesis in hibernators. In: Living in the Cold, ed. by H. C. Heller, X. J. Musacchia and L. C. H. Wang, Elsevier: New York, 1986, pp. 101-108.
88. Moore, B. J., J. S. Stern, and B. A. Horwitz. Brown adipose tissue mediates the increased expenditure of cold-exposed overfed rats. Am. J. Physiol. 251:R518-R524, 1986.
89. Wickler, S. J., B. A. Horwitz, and J. S. Stern. Blood flow to brown fat in lean and obese adrenalectomized Zucker rats. Am. J. Physiol. 251:R851-858, 1986.
90. Kott, K. S., B. J. Moore, and B. A. Horwitz. Decreased testosterone levels do not mediate short photoperiod-induced brown fat changes. Am. J. Physiol. 251:R963-970, 1986.

91. Wickler, S. J., J. S. Stern, Z. Glick, and B. A. Horwitz. Thermogenic capacity and brown fat in rats exercise-trained by running. Metabolism 36:76-81, 1987.
92. Berce, P. J., B. J. Moore, B. A. Horwitz, and J. S. Stern. Metabolism at thermoneutrality and in the cold is reduced in the neonatal preobese Zucker (fa/fa) rat. J. Nutrit. 116:2478-2485, 1987.
93. Tomasi, T. and B. A. Horwitz. Thyroid function and cold acclimation in the hamster (Mesocricetus auratus). Am. J. Physiol. 252:E260-E267, 1987.
94. Tomasi, T., J. S. Hamilton, and B. A. Horwitz. Thermogenic capacity in shrews. J. Thermal Biol., in press.
95. Wickler, S. J., B. A. Horwitz, and K.S. Kott. Muscle function in hibernating hamsters: a natural analog to bed rest? J. Thermal Biol., in press.
96. McDonald, R. B., J. S. Stern, and B. A. Horwitz. Cold-induced metabolism and brown fat GDP binding in young and old rats. Expt. Gerontol., in press.



## G. BUDGET

Grant No.: NASA NAG 2-341

Period covered by budget: March 1, 1988 - February 28, 1991

	<u>1988-89</u>	1989-90	1990-91
<u>SALARIES</u>			
Staff Res. Assoc.	16,000	17,000	20,000
Employee Benefits	4,640	5,100	6,200
Total Salaries	20,640	22,100	26,200
<u>SUPPLIES</u>	3,000	4,000	5,000
Animal procurement & care, misc. lab supplies, publication costs, equipment maintenance.			
<u>EQUIPMENT</u>	4,500	3,000	1,000
Electronics & micromanipulators for intracellular recording and signal processing.			
<u>TRAVEL</u>	2,000	2,100	2,300
P.I. to present papers at FASEB and at ASGSB annual meetings.			
TOTAL DIRECT COSTS	30,140	31,200	34,500
TOTAL INDIRECT COSTS (39.2%)	10,051	11,055	13,132
TOTAL	<u>\$40,191</u>	<u>\$42,255</u>	<u>\$47,632</u>

## H. BUDGET JUSTIFICATION

We have the equipment required for the proposed experiments on hippocampal slices except for two micromanipulators to hold additional electrodes for stimulating fiber tracts and/or to hold pipettes for iontophoretic injection of compounds and for measuring electrical potentials.

## I. AVAILABLE FACILITIES

We have the standard hippocampal slice equipment required to stimulate a fiber track (in our experiments Schaffer collaterals) and record extracellular and intracellular activity in hippocampal pyramidal cells. Thus we have an air table, Brown-Flaming microelectrode puller, WPI amplifier, Zenith computer, XY plotters, WPI stimulators, and associated electronic equipment.

The PI has a laboratory of over 800 sq ft. In addition, the facilities at the Chronic Acceleration Research Unit on the Davis Campus, including animal centrifuges in the Unit, will be used to expose rats to a hypergravic environment.