

# Solutions of the Quantum Yang – Baxter Equations Associated with $(1-3/2)$ -D Representations of $SU_q(2)$

Huang Yijun, Yu Guochen and Sun Hong\*  
 Department of Foundation, the First  
 Aeronautical College of Air Force,  
 Xinyang Henan 464000 P . R . China .

## Abstract

The solutions of the spectral independent QYBE associated with  $(1-3/2)$ -D representations of  $SU_q(2)$  are derived, based on the weight conservation and extended Kauffman diagrammatic technique . It is found that there are nonstandard solutions .

## 1 Introduction

It is well known that the quantum Yang – Baxter equations ( QYBE ) play an important role in various theoretical and mathematical physics, such as completely integrable systems in  $(1+1)$  dimensions, exactly solvable models in statistical mechanics, the quantum inverse scattering method and the conformal field theories in 2-dimensions .<sup>[1]-[7]</sup> Recently, much remarkable progress has been made in construction the solutions of the QYBE associated with the representations of Lie algebras .<sup>[8]-[9]</sup> In this paper we derive the solutions of the spectral independent QYBE associated with  $(1-3/2)$ -D representations of  $SU_q(2)$ , based on the weight conservation and extended Kauffman diagrammatic technique . It is found that there are nonstandard solutions .

## 2 Braid relations of $(1-3/2)$ -D representations of $SU_q(2)$

We know that there is the relation for Universal R – matrix:

$$R_{12}^{j_1 j_2} R_{13}^{j_1 j_3} R_{23}^{j_2 j_3} = R_{23}^{j_2 j_3} R_{13}^{j_1 j_3} R_{12}^{j_1 j_2} \quad (2.1)$$

We define the new R – matrix:

$$\bar{R}^{j_1 j_2} = P R^{j_1 j_2} \quad (2.2)$$

Where P is the transposition  $(P: V^{j_1} \otimes V^{j_2} \rightarrow V^{j_2} \otimes V^{j_1})$

Then the eq . (2-1) can be rewritten as follows

$$\bar{R}_{12}^{j_1 j_2} \bar{R}_{23}^{j_1 j_3} \bar{R}_{12}^{j_2 j_3} = \bar{R}_{23}^{j_2 j_3} \bar{R}_{12}^{j_1 j_3} \bar{R}_{23}^{j_1 j_2} \quad (2.3)$$

\* Address: Jinan 250023

For the  $(1-3/2)$ -D representation of  $SU_q(2)$ ,  $(j_1, j_2, j_3) \in (1, 1, 3/2)$ , then eq. (2.3) gives the following relations

$$\bar{R}_{12}^{-11} \bar{R}_{23}^{-1\ 3/2} \bar{R}_{12}^{-1\ 3/2} = \bar{R}_{23}^{-1\ 3/2} \bar{R}_{12}^{-1\ 3/2} \bar{R}_{23}^{-11} \quad (2.4-1)$$

$$\bar{R}_{12}^{-3/2\ 1} \bar{R}_{23}^{-11} \bar{R}_{12}^{-11} = \bar{R}_{23}^{-11} \bar{R}_{12}^{-3/2\ 1} \bar{R}_{23}^{-3/2\ 1} \quad (2.4-2)$$

$$\bar{R}_{12}^{-1\ 3/2} \bar{R}_{23}^{-11} \bar{R}_{12}^{-3/2\ 1} = \bar{R}_{23}^{-3/2\ 1} \bar{R}_{12}^{-11} \bar{R}_{23}^{-1\ 3/2} \quad (2.4-3)$$

These are the braid relations associated  $(1-3/2)$ -D representations of  $SU_q(2)$ . We suppose that the  $\bar{R}$  satisfies the C-P invariance, then eq. (2.4-1) is equal to eq. (2.4-2).

### 3 The weight conservation and the solutions of QYBE

To determine the structure for the solutions, We consider the weight conservation

$$(\bar{R})_{cd}^{ab} = 0 \quad \text{unless } a+b=c+d \quad (3.1)$$

where

$$\bar{R} = \bar{R}^{-1\ 3/2}, \quad \bar{R}^{-3/2\ 1}, \quad \bar{R}^{-11}$$

$$a, b, c, d \in (\pm 3/2, \pm 1/2, \pm 1, 0)$$

It is well known that  $\bar{R}^{-11}$  which satisfied the conditions of c-p invariance and eq. (3.1) be written as

$$\begin{aligned} \bar{R}^{-11} = & \sum_a u_a E_{aa} \otimes E_{aa} + \sum_{a < b} W^{(a, b)} E_{ab} \otimes E_{ab} + \sum_{a \pm b} P^{(a, b)} E_{ab} \otimes E_{ba} \\ & + \sum_{a \leq b} q^{(a, c)} E_{ab} \otimes E_{cd} + E_{cd} \otimes E_{ab} \end{aligned} \quad (3.2)$$

Where

$$u_0 = 1, u_{\pm 1} = q^2, \quad p^{(0, 1)} = p^{(1, 0)} = 1, \quad p^{(+1, +1)} = q^{-2}$$

$$w^{(0, 1)} = w = q^2 - q^{-2}, \quad w^{(-1, 1)} = (1 - q^{-2}) w, \quad q_0^{(-1, 0)} = q_0^{(0, -1)} = q^{-1} w \quad (3.3)$$

By the weight conservation  $\bar{R}^{-1\ 3/2}$  can be constructed in the form

$$\bar{R}^{-13/2} = \sum_{a, b} P_{a+b}^{a, b} E_{ab} \otimes E_{ba} + \sum_{\substack{a < d \\ c < d}} q^{a, c} E_{ac} \otimes E_{bd} \quad (3.4)$$

Where

$$a, b \in (\pm 1, 0); \quad b, c \in (\pm 3/2, \pm 1/2) \quad P_{a+b}^{(a, b)} \quad \text{and} \quad q_{a+b}^{(a, c)}$$

are the determined parameters .

Substituting eq . (3.2), (3.4) into eq . (2.4-1.3), We obtain the unknown parameters by extended Karffiman diagrammatic techique .

$$P_{5/2}^{(1, 3/2)} = q^3, \quad P_{-5/2}^{(-1, -3/2)} = P_{5/2}^{(1, 3/2)} \quad Q = q^3 Q \quad (3.5-1)$$

$$P_{-3/2}^{(-1, -1/2)} = P_{3/2}^{(1, 1/2)} \quad Q = qQ, \quad P_b^{(0, b)} = Q^{1/2}$$

$$q_{3/2}^{(0, 1/2)} = Q^{-1/2} q_{3/2}^{(-1, -3/2)} = (1 - q^{-2}) q^{3/2} ([3]!)^{1/2} Q^{1/4}$$

$$q_{1/2}^{(-1, 1/2)} = Q^{1/2} q_{-1/2}^{(0, -3/2)} = (1 - q^{-2}) q^{-1/2} ([3]!)^{1/2} Q^{1/4}$$

$$q_{1/2}^{(0, -1/2)} = Q^{-1/2} q_{1/2}^{(-1, -1/2)} = (1 - q^{-2}) q^{1/2} ([2]!)^{3/2} Q^{1/4}$$

$$q_{1/2}^{(-1, -1/2)} = q_{-1/2}^{(-1, -3/2)} = (1 - q^{-2}) q ([2]![3]!)^{1/2} Q^{3/4} \quad (3.5-2)$$

Where

$$[u] \equiv \frac{q^u - q^{-u}}{q - q^{-1}}, \quad [u]! \equiv [u][u-1] \dots [1], \quad [0]! \equiv 1 \quad (3.5-3)$$

Substituting eq.(3.5) into eq.(3.4), we obtain the solutions  $\bar{R}^{-13/2}$  . And we obtain the solutions  $\bar{R}^{3/2}$  by employing the c-p invariance .

We have derived the solutions of the spectral independent QYBE associated with  $(1-3/2)$ -D representations . It is easy to see that there is a new arbitrary parameter,  $Q$ , then there are new solutions . In fact when  $Q=1$ , the solutions is Universal R-matrix of  $SU_q$  (2).

$$(\bar{R}^{j_2 j_1})_{m_1 m_2}^{m_2 m_1} = \delta_{m_1+m_2}^{m_1'+m_2'} \frac{[(1 - q^{-2})^{m_1' - m_1}]}{[m_1' - m_1]!} q^{m_1 m_2' + m_2 m_1' - 1/2 (m_1' - m_1)(m_1' - m - 1)}$$

$$\left\{ \frac{(\hat{j}_1 + m_1')! (\hat{j}_1 - m_1)! (\hat{j}_1 - m_2)! (\hat{j}_2 + m_2)!}{(\hat{j}_1 - m_1)! (\hat{j}_1 + m_1)! (\hat{j}_2 + m_2)! (\hat{j}_2 - m_2)!} \right\}^{1/2} \quad (3.6)$$

Standard solutions . When  $Q \neq 1$ , there are new solutions .

## Referene

- [1] C . N . Yang, *phys . Rev . Lett .* 19 (1967) 1312; *Phys . Rev .* 168 (1968) 1920.
- [2] R . J . Baxter, Exactly Solved Models in Statistical Mechanics, Academic London (1982) .
- [3] A . B . Zamolodchikov and Al . B . Zamolochikov, *Am . of Phys .* 120 (1979) 253 .
- [4] L . D . Faddeev, Integrable Models in (1+1) -D Quantum Field Theory . Les Houches Sesio XXXIX . (1982) 536 .
- [5] E . X . Skyanin, L . A . Takhtajan and L . D . Faddeev, *Math . Phys .* 40 (1979) 194 (in Russian)
- [6] L . A . Takhtajan and L . D . Faddeev, *Uspekhi Mat . Nauk .* 24 (1979) 13 .
- [7] G . Segal, *Comformal field theory*, Oxford pereprint 1987 .
- [8] M . L . G . and K . Xue, *phys . Lett .* A146 (1990) 245 .
- [9] A . N . Kirillov and N . Yu, *Reshetikhin . Lomt . Preprint* (1988) .