

## LUMINOSITIES OF H $\alpha$ EMITTING REGIONS IN A PAIR OF INTERACTING GALAXIES IN THE BOOTES VOID

D. Weistrop, P. Hintzen, University of Nevada, Las Vegas, NV

R. Kennicutt, C. Liu, J. Lowenthal, University of Arizona, Tucson, AZ

K.-P. Cheng, R. Oliverson, B. Woodgate, NASA/Goddard Space Flight Center, Greenbelt, MD

**Abstract:** Luminosities of H $\alpha$  emission from a pair of interacting galaxies in the low density environment of the Bootes void are presented. CG 692 (IRAS 1519+5050) has an H $\alpha$  luminosity of  $2 \times 10^{42}$  ergs s $^{-1}$ , indicating a star formation rate of 18.4 M $_{\odot}$  yr $^{-1}$ . Individual extranuclear H $\alpha$  regions have luminosities of approximately  $10^{40}$  ergs s $^{-1}$ . These luminosities are similar to those found for H II regions in bright, late-type galaxies in more densely populated parts of the Universe.

Based on spectroscopy and imaging data, we have recently identified a pair of interacting galaxies within the Bootes void (Weistrop *et al.* 1991, Weistrop *et al.* 1992). The brighter galaxy, CG 692 (IRAS 1519+5050) is a spiral undergoing large amounts of star formation, while its companion, CG 693, is a previously unidentified Seyfert 1 galaxy (Figure 1). The galaxies are at the same redshift within the uncertainties of the measurements,  $z = 0.0574$ , and have a projected separation of 34 kpc, ( $H_0=50$  km s $^{-1}$ Mpc $^{-1}$  assumed throughout). The galaxies are within the boundaries of the Bootes void as defined by Kirshner *et al.* (1987). We investigate the luminosity of the H $\alpha$  emission in CG 692 and between the galaxies, to compare with galaxies in denser environments. Almost all the H $\alpha$  emission from CG 693 arises in the nucleus, and is not considered further here.

Images were obtained at the redshifted H $\alpha$  wavelength of the galaxies and a nearby continuum band, using the Goddard Fabry-Perot Imager on the University of Arizona's 90-inch telescope located at Kitt Peak (Weistrop *et al.* 1992). Flux calibration was obtained from observations of HZ 44. The total H $\alpha$  flux for CG 692 is  $1.38 \times 10^{-13}$  ergs cm $^{-2}$  s $^{-1}$ , with an estimated  $\pm 25\%$  error due to the uncertainty in the conversion from count rate to flux. Since the galaxies are located at  $b = 53^\circ$ , absorption within our galaxy is about 0.02 mag and has been ignored (Kennicutt & Kent 1983). For  $q_0 = 0$ , the total H $\alpha$  luminosity of CG 692 is  $L = 2.06 \times 10^{42}$  ergs s $^{-1}$ , similar to the H $\alpha$  + [NII] luminosities of the brightest interacting galaxies (Kennicutt *et al.* 1987). The star formation rate in CG 692 is 18.4 M $_{\odot}$  yr $^{-1}$  using the conversion given by Kennicutt (1983).

We have determined the luminosity of several extranuclear H $\alpha$  features in this system (Table 1). These regions are not spatially resolved. B and C, which are between the galaxies and may be the result of the interaction, have H $\alpha$  luminosities similar to the luminosities of the brightest H II regions in field spiral and luminous irregular galaxies (Kennicutt 1988). (Values must be converted to  $H_0=75$  km s $^{-1}$ Mpc $^{-1}$  for comparison with Kennicutt's results.) J and K, the features beyond the end of CG 692's spiral arm, have luminosities similar to those of 'giant' or 'supergiant' H II regions found in late-type normal or peculiar galaxies. The knots within the

Table 1. H $\alpha$  Luminosities<sup>1</sup>

Feature	$\log[L(H\alpha)]$
B	39.78
C	39.86
J	40.47
K	40.55
$\log[L(H\alpha) \text{ arcsec}^{-2}]$	
D	39.74
F	39.69
G	39.57
H	39.75

<sup>1</sup> $H_0=50 \text{ km s}^{-1}\text{Mpc}^{-1}$




Fig. 1. Contour plot of the H $\alpha$  image. The outermost contour represents a flux of  $8.3 \times 10^{-17} \text{ ergs cm}^{-2} \text{ s}^{-1} \text{ arcsec}^{-2}$ . Successive contours increase in intervals of  $2.76 \times 10^{-16} \text{ ergs cm}^{-2} \text{ s}^{-1} \text{ arcsec}^{-2}$ .

spiral arm are concentrations of emission embedded in an area of strong, extended H $\alpha$  emission. In Table 1 we give the luminosities per square arcsec for several locations along the spiral arm. The total H $\alpha$  emission from the strongest knot, H, is about  $5.3 \times 10^{40} \text{ ergs s}^{-1}$ .

Kennicutt (1988) finds a relationship between the average luminosity of the brightest H II regions in a galaxy and that galaxy's absolute magnitude and Hubble type.  $M_B$  for CG 692 can be calculated from the apparent magnitude estimate given by Sanduleak & Pesch (1987). For CG 692,  $B = 15$ , giving  $M_B = -22.8$ . A discussion of the accuracy of the Case magnitudes is given by Weistrop & Downes (1991). Kennicutt's relationship predicts the mean luminosity of the brightest H II regions to be  $\geq 10^{40} \text{ ergs s}^{-1}$  for late type field or Virgo cluster spirals with  $M_B = -22$  ( $H_0=75 \text{ km s}^{-1}\text{Mpc}^{-1}$ ). This is similar to the luminosity we obtain for the brightest extranuclear H II regions in CG 692.

This work was supported in part by NASA grant NAS5-31231.

#### References

- Kennicutt, Jr., R.C. 1983, ApJ, 272, 54.  
 \_\_\_\_\_ 1988, ApJ, 344, 144.  
 Kennicutt, Jr., R.C., & Kent, S.M. 1983, AJ, 88, 1094.  
 Kennicutt, Jr., R.C., Keel, W.C., van der Hulst, J.M., Hummel, E., & Roettiger, K.A. 1987, AJ, 93, 1011.  
 Kirshner, R.P., Oemler, A., Schechter, P.L., and Shectman, S.A. 1987, ApJ, 314, 493.  
 Sanduleak, N., & Pesch, P. 1987, ApJSS, 63, 809.  
 Weistrop, D., & Downes, R.A. 1991, AJ, 102, 1680.  
 Weistrop, D., et al. 1991, BAAS, 23, 1428.  
 Weistrop, D., et al. 1992, ApJL (submitted).