

30-01
145884
N93-19143

OBSERVATIONS OF COMETARY PARENT MOLECULES WITH THE IRAM RADIO TELESCOPE.

P. Colom¹⁾, D. Despois²⁾, G. Paubert³⁾, D. Bockelée-Morvan¹⁾, J. Crovisier¹⁾

- 1) Observatoire de Paris, Section de Meudon, F-92195 Meudon, France
2) Observatoire de Bordeaux, BP 89, Avenue Pierre Sémirot, F-33270 Bordeaux, France
3) IRAM, Avenida Divina Pastora, 7, N.C., E-18012 Granada, France

Abstract

Several rotational transitions of HCN, H₂S, H₂CO and CH₃OH were detected in comets P/Brorsen-Metcalf 1989 X, Austin (1989c1) and Levy (1990c) with the IRAM 30-m radio telescope. This allows us to determine the production rates of these molecules and to probe the physical conditions of the coma.

OBSERVATIONS

Comets P/Brorsen-Metcalf 1989 X, Austin (1989c1) and Levy (1990c) were observed on September 2-7 1989, May 21-25 and August 26-31 1990, respectively, with the IRAM (Institut de Radio Astronomie Millimétrique) 30-m radio telescope at Pico Veleta (Spain). Three SIS mixer receivers were used simultaneously (85-115, 130-170 and 209-270 GHz). The spectrometers consisted in two banks of 128x100 kHz channels, two banks of 512x1 MHz channels, and an AOS (Acousto Optical Spectrometer) of 864 channels with a 505 MHz bandwidth.

RESULTS

The results concerning the detected species are summarized in the table. More details were (or will be) published by Bockelée-Morvan *et al.* (1990, 1991), Colom *et al.* (1990, 1992) and Crovisier *et al.* (1990, 1991).

HCN, H₂CO and H₂S production rates were derived from the observed line intensities using models treating the evolution of the excitation conditions from the collision dominated region (inner coma, collisions with H₂O, $s = 10^{-14}$ cm², $T_{\text{kin}} = 50$ K) to the radiation dominated region (outer coma, IR excitation of the vibrational bands by the Sun). For CH₃OH, we assume LTE and used a rotational temperature of 30 K, in agreement with the observed relative line intensities. For the density distribution we assumed isotropic outflow from the nucleus at constant velocity (0.8 km s⁻¹) and took into account the molecular lifetime against photodissociation.

Hydrogen cyanide

The J(1-0) 89 GHz and J(3-2) 266 GHz rotational transitions of HCN were marginally detected in comet P/Brorsen-Metcalf, whereas clear detections were obtained in comet Austin (1989c1) and Levy (1990c). HCN seems to be more abundant by at least a factor of two in periodic comets (P/Halley, P/Brorsen-Metcalf) than in non periodic comets (Wilson, Austin, Levy). This suggests a chemical difference between periodic and new comets. The very low upper limit obtained on the relative abundance of HC₃N (5×10^{-5}) shows that it is not the major lacking source of CN radicals.

Formaldehyde

The observations of the H₂CO 3₁₂-2₁₁ transition at 226 GHz in comet P/Brorsen-Metcalf gave only a marginal detection (S/N = 4). The 226 GHz line was easily detected in comet Austin (S/N = 10) and in comet Levy (S/N = 8). Observations of the 5₁₅-4₁₄, 2₁₂-1₁₁, 3₀₃-2₀₂, 3₂₂-2₂₁ and 3₂₁-2₂₀

lines were negative, in agreement with excitation models (Bockelée-Morvan and Crovisier 1992). Production rates inferred in the assumption of release from the nucleus show that formaldehyde is a minor component of the nucleus with an abundance relative to water which ranges from 4×10^{-4} in Levy to 3×10^{-3} in P/Brosen-Metcalf (Colom *et al.* 1992). These abundances are at least an order of magnitude less than the Vega IKS value for P/Halley (4%; Combes *et al.* 1988).

Hydrogen sulfide


The observation of H_2S $1_{10}-1_{01}$ at 169 GHz in comet Austin led to the first detection of hydrogen sulfide in a comet. In addition to the 169 GHz ortho line, the $2_{20}-2_{11}$ para line of H_2S at 217 GHz was detected in comet Levy. H_2S is a minor component, with a relative abundance of 2×10^{-3} . The other sulfur-bearing molecules observed (SO_2 , OCS, H_2CS) are less abundant than hydrogen sulfide. (Crovisier *et al.* 1991.)

Production rates and abundances.

Comet and molecule	Date	Q a) [s ⁻¹]	Q/Q[H ₂ O] b)
<i>P/Brosen-Metcalf (1989 X)</i>			
HCN J(1-0)	89/09/04-07	4.5×10^{26}	1.8×10^{-3}
H_2CO $3_{12}-2_{11}$	89/09/04-07	7.6×10^{26}	3.0×10^{-3}
<i>Austin (1989c1)</i>			
HCN J(1-0)	90/05/23	2.0×10^{25}	5.0×10^{-4}
H_2CO $3_{12}-2_{11}$	90/05/21-25	4.6×10^{25}	1.1×10^{-3}
H_2S $1_{10}-1_{01}$	90/05/24-25	1.1×10^{26}	2.7×10^{-3}
CH_3OH (3,0)-(2,0)A	90/05/25	2.0×10^{26}	5.0×10^{-2}
<i>Levy (1990c)</i>			
HCN J(1-0)	90/08/29	6.6×10^{25}	2.6×10^{-4}
H_2CO $3_{12}-2_{11}$	90/08/26-30	1.0×10^{26}	4.0×10^{-4}
H_2S $1_{10}-1_{01}$	90/08/30-31	5.0×10^{26}	2.0×10^{-3}
CH_3OH (3,0)-(2,0)A	90/08/27	1.8×10^{27}	7.2×10^{-3}
HC_3N J(24-23)	90/08/27	$< 1.2 \times 10^{25}$	$< 5.0 \times 10^{-5}$
SO_2 $7_{17}-6_{06}$	90/08/29	$< 6.0 \times 10^{26}$	$< 2.5 \times 10^{-3}$
OCS J(18-17)	90/08/28	$< 5.0 \times 10^{26}$	$< 2.0 \times 10^{-3}$
H_2CS $4_{14}-3_{13}$	90/08/28	$< 2.5 \times 10^{26}$	$< 1.0 \times 10^{-3}$

a Assuming a parent distribution.

b Q/[H₂O] from OH 18-cm observations: 2.5×10^{29} s⁻¹ for P/Brosen-Metcalf and Levy, 4.0×10^{28} s⁻¹ for Austin.



The spectra of methanol (CH_3OH) observed with the IRAM 30-m radio telescope in comet Levy (1990c). The upper panels show spectra observed with a 1-MHz spectral resolution. The lower panel shows parts of the spectra around 97 GHz and 145 GHz observed with a 100-kHz resolution.

Methanol

CH₃OH was detected in comet Austin through its J(2-1) $\Delta K = 0$ transitions at 97 GHz and its J(3-2) $\Delta K = 0$ transitions at 145 GHz. It was the first detection of methanol in a solar system body. A dozen of CH₃OH lines were detected in comet Levy, as is shown in the Figure. Methanol is a substantial component of the nucleus, with a relative abundance of the order of 1% in comets Austin and Levy.

Other molecules.

Limits on many other interesting lines of potential parent molecules were also obtained, either during dedicated searches or serendipitously. Here is a preliminary list (some of the corresponding limits on the production rates are given in the table; the other ones are presently under evaluation):

Hydrocarbons: CH₃CCH (propyne; several lines at 85.3 GHz); c-C₃H₂ (cyclopropenylidene: many lines).

OH species: HDO (deuterated water: 3₁₂-2₂₁ line at 225.897 GHz).

CHO species: many lines of HCOOH (formic acid), CH₃CHO (acetaldehyde), C₂H₅OH (ethanol).

Nitrogen compounds: HC₃N (cyanoacetylene: 24-23 at 218.325 GHz); CH₃NH₂ (methylamine: lines around 85.4 and 88.6 GHz); CH₂NH (methanimine: 1₁₀-0₀₀ at 225.5 GHz); HNCO (isocyanic acid); NH₂CHO (formamide).

Sulfur compounds: SO₂, OCS, H₂CS (see Crovisier et al. 1991).

References

- Bockelée-Morvan D., Colom P., Crovisier J., Despois D., and Paubert G. (1991) Microwave detection of hydrogen sulphide and methanol in comet Austin (1989c1). *Nature* **350**, 318-320.
- Bockelée-Morvan D., and Crovisier J. (1992) Formaldehyde in comets: II. Excitation of rotational lines. *Astron. Astrophys.* (in press).
- Bockelée-Morvan D., Crovisier J., Colom P., Despois D., and Paubert G. (1990). Observations of parent molecules in comets P/Brorsen-Metcalf (1989o), Austin (1989c1) and Levy (1990c) at millimetre wavelengths: HCN, H₂S, H₂CO and CH₃OH. In *Formation of Stars and Planets and the Evolution of the Solar System*. ESA SP-315, 243-248.
- Colom P., Crovisier J., Bockelée-Morvan D., Despois D., and Paubert G. (1992). Formaldehyde in comets: I. Microwave observations of P/Brorsen-Metcalf 1989 X, Austin (1989c1) and Levy (1990c). *Astron. Astrophys.* (in press).
- Colom P., Despois D., Bockelée-Morvan D., Crovisier J., and Paubert G. (1990). Millimeter observations of comets P/Brorsen-Metcalf (1989o) and Austin (1989c1) with the IRAM 30-m radio telescope. In *Workshop on Observations of Recent Comets*. Eds W.F. Huebner et al., pp. 80-85.
- Combes et al. (1988) The 2.5 to 12 microns spectrum of comet Halley from the IKS-VEGA experiment. *Icarus* **76**, 404-436.
- Crovisier J., Bockelée-Morvan D., Colom P., Despois D., and Paubert G. (1990). Observations of parent molecules in comet Austin (1989c1) at millimeter wavelengths: HCN, H₂S, H₂CO, CH₃OH. *Bull. Amer. Astron. Soc.* **22**, 1089.
- Crovisier J., Despois D., Bockelée-Morvan D., Colom P., and Paubert G. (1991) Microwave observations of hydrogen sulfide and searches for other sulfur compounds in comets Austin (1989c1) and Levy (1990c). *Icarus* **93**, 246-258.