

An Interim Report for NASA Grant NAG-5-1843 for January to November, 1992

"Experimental Studies of the Far-Infrared Spectra of Cosmic-Type Ices"

GODDARD
GRANT
1N-89-CR
131944
p-27

Submitted to:

National Aeronautics and Space Administration
Goddard Space Flight Center
Greenbelt, MD 20771


Attention: Ms. Gloria R. Blanchard
Grants Officer, Code 286.1

Submitted by:

Dr. Reggie L. Hudson
Department of Chemistry
Eckerd College
St. Petersburg, FL 33733

November 24, 1992

Respectfully Submitted,



Reggie L. Hudson

(NASA-CR-191300) EXPERIMENTAL
STUDIES OF THE FAR-INFRARED SPECTRA
OF COSMIC-TYPE ICES Interim Report,
Jan. - Nov. 1992 (Eckerd Coll.)
27 p

N93-13660

Unclass

G3/89 0131944

The purpose of the grant was two-fold. First, travel money was budgeted for the principal investigator to attend the Second International Workshop on the Nature of Cometary Organic Matter (December 9-13, 1991) at Asilomar, California. Second, funding was requested for a study of the far-infrared spectra of astronomical ices.

Travel to Meeting

The meeting attended led to several significant events. The PI was able to interact with NASA and non-NASA scientists investigating cosmic-type ices. Experiments were compared and laboratory methods contrasted. New results were discussed with both the invited speakers and the authors of poster papers.

The PI presented a poster of his work with NASA scientist Dr. Marla H. Moore (NASA/GSFC Code 691). An abstract is reproduced as Appendix I.

Research

The grant also was used for research expenses associated with a study of clathrate hydrates. A manuscript describing the research was prepared and submitted to *Astrophysical Journal Letters*. A copy of the letter accepting the manuscript for publication and a copy of the manuscript itself are included as Appendix II.

Remaining Funds

The remaining funds, about \$ 1000, will be used to continue research on cosmic-type ices in 1993. A final report will be submitted at a later date.

APPENDIX I

FAR INFRARED SPECTRA OF AMORPHOUS AND CRYSTALLINE WATER ICE AND

CHANGES IN THESE PHASES AS THE RESULT OF PROTON IRRADIATION

M. H. Moore, and R. L. Hudson*

NASA/Goddard Space Flight Center,

Astrochemistry Branch,

Greenbelt, Maryland 20771

Far infrared spectra from 20 μm (500 cm^{-1}) to 100 μm (100 cm^{-1}) of water ice have been measured. Amorphous ice deposited at 13 K has one absorption band at 45 μm (220 cm^{-1}). Amorphous ice evolves into a crystalline form with absorptions at 44 μm (229 cm^{-1}) and 62 μm (162 cm^{-1}) as the temperature is increased to 155 K. Spectra documenting this phase change will be presented as well as spectra of crystalline ice at temperatures between 13 K and 155 K.

We also will present far infrared spectra of amorphous and crystalline water ice before and after proton irradiation. Changes in these two forms will be discussed in relation to ices in comets, grains, and planetary satellites in various radiation environments.

*Permanent address: Eckerd College, St. Petersburg, FL 33733

THE ASTROPHYSICAL JOURNAL LETTERS

A. DALGARNO, *Editor*
EUGENE H. AVRETT, *Deputy Editor*
Center for Astrophysics
60 Garden St., MS 14
Cambridge, MA 02138

telephone: 617-495-4479
fax: 617-495-8316
bitnet: apjlref@cfa
internet: apjlref@cfa.
Harvard.edu

November 20, 1992 : span: cfa7::apjlref

Dear Dr. Hudson:

Your paper FAR-INFRARED INVESTIGATIONS OF A METHANOL CLATHRATE HYDRATE:
IMPLICATIONS FOR ASTRONOMICAL OBSERVATIONS

has been accepted for publication in THE ASTROPHYSICAL JOURNAL LETTERS and has been sent by overnight express mail to the University of Chicago Press, where it will receive immediate copyediting and be sent to the typesetter. If you have any changes to make, please FAX your changes to Chicago at the number listed below:

THE ASTROPHYSICAL JOURNAL
The University of Chicago Press
5608 S. Stony Island Avenue
Chicago, IL 60637

FAX: 312-753-0811
Telephone: 312-753-8021
Telex: 6503511226

Only minor changes can be made in the proof stage without delaying the publication of your paper. The cost of substantial resetting of type will be billed to the author.


Proofs are sent to authors by overnight express mail to domestic and foreign addresses. You should receive proofs within two weeks and we ask that you respond immediately to the Chicago office with your corrections and approval.

The assignment of your paper to a particular issue of the Journal is made after the Press has received your corrections. The issue date is an average of 6 weeks after proofs are received in Chicago; however, if you urgently need an exact date, call Tulie O'Connor at 312-753-3372 after Chicago receives your proofs.

It is imperative that the enclosed offprint order and page authorization cards be completed and sent to the Chicago office as soon as possible. Since offprints are an extension of the issue print run, we must have your offprint order as soon as possible, or no offprints can be supplied.

With thanks for your cooperation, I am,

Yours sincerely,


A. Dalgarno
Letters Editor

FAR-INFRARED INVESTIGATIONS OF A METHANOL
CLATHRATE HYDRATE:
IMPLICATIONS FOR ASTRONOMICAL OBSERVATIONS

by

Reggie L. Hudson

Department of Chemistry, Eckerd College

St. Petersburg, FL 33733

and

Marla H. Moore

Astrochemistry Branch, NASA/Goddard Space Flight Center

Greenbelt, MD 20771

ABSTRACT

Observations of non-terrestrial clathrate hydrates are still lacking despite the fact that clathrates first were suggested to exist in cometary and interstellar ices over forty years ago. Spectroscopy, the most direct method of astronomical detection, has been hampered by the similarity of clathrate hydrate spectra to those of unenclathrated guest molecules and solid H_2O . We have prepared a methanol (CH_3OH) clathrate hydrate, using a recently published procedure, and have investigated its far-IR spectrum. The spectrum is quite different from that of either unenclathrated CH_3OH or solid H_2O and so should be of value in astronomical searches for this clathrate.

I. INTRODUCTION

Clathrate hydrates are crystalline solids consisting of cages of hydrogen-bonded H_2O molecules enclosing a "guest" species, usually a stable molecule. The cages' structure is distinctive and is different from the arrangement of H_2O molecules in either pure amorphous or crystalline H_2O (Davidson 1973; Jeffrey 1984). Although it was suggested more than four decades ago that these hydrates exist in cometary and interstellar ices (Delsemme & Swings 1952), there is as yet no direct evidence for clathrate hydrates in any non-terrestrial object. Astronomical observations of infrared (IR) sources might be expected to detect clathrate hydrates, but near- (Smythe 1975), mid- (Bertie & Othen 1973a; Bertie & Solinas 1975), and far-IR (Bertie & Jacobs 1977, 1978) spectra of clathrates often are dominated by broad features of the H_2O cages, which usually resemble H_2O ice spectra, and by guest molecule features, which typically differ little from spectra of the pure unenclathrated guest. In this letter we present the far-infrared spectrum of a methanol (CH_3OH) clathrate hydrate. The spectrum is very different from that of either pure H_2O or pure CH_3OH , and should be valuable in searches for the CH_3OH clathrate hydrate in astronomical objects. The fact that the guest molecule is

CH₃OH is of special importance since this molecule has been observed in several recent comets (Bocklee-Morvan *et al.* 1991) and is thought to be important in some interstellar clouds (Tielens & Allamandola 1987).

Clathrates are most commonly identified in the laboratory by x-ray or electron diffraction. Infrared spectroscopy can also be used for clathrate hydrate identification, most reliably in systems where clathrate formation is known to occur from diffraction data (Bertie & Othen 1972, 1973b; Bertie & Devlin 1983). Blake *et al.* (1991) recently have published an elegant set of experiments along these lines in which electron diffraction and mid-IR spectroscopy were combined to investigate the CH₃OH clathrate hydrate. When an amorphous H₂O:CH₃OH = 2:1 mixture was deposited at 85 K and warmed to 130 K, electron microscopy and electron diffraction showed that a single polycrystalline phase formed, with some remaining amorphous material. The diffraction pattern of the crystalline phase was indexed as the type II (Davidson 1973; Jeffrey 1984) CH₃OH clathrate hydrate. However, the authors reported that only "a small change was noticed in the profiles of some of the CH₃OH bands" in the mid-IR spectrum. Spectroscopic evidence for enclathration was

obtained by using a small amount of CO₂ as a tracer gas in a H₂O:CH₃OH mixture. (Earlier work had established the IR spectra of CO₂ clathrates (Fleyfel & Devlin 1991).)

We have independently verified the results of Blake *et al.* (1991) for the mid-IR spectrum (4000 - 400 cm⁻¹ or 2.5 - 25 μm) of the CH₃OH clathrate hydrate: spectra from 13 K to 140 K indeed are dominated by absorptions of CH₃OH and H₂O. However, by working in the far-infrared region from 400 to 100 cm⁻¹ (25 - 100 μm) we have found some rather pronounced spectral features, different from those of either CH₃OH or H₂O alone. We also have irradiated the CH₃OH clathrate with a proton beam to simulate cosmic ray bombardment. To our knowledge these are the only *in situ* radiation chemical experiments that have been done on a cometary or interstellar clathrate hydrate analog.

II. EXPERIMENTAL PROCEDURE

The experimental procedure followed was largely the same as that used for the study of other ices (Hudson & Donn 1991; Hudson & Moore 1992; Moore & Hudson 1992). Most of our experiments used slightly greater warming rates and annealing times at the higher

temperatures than those of Blake *et al.* (1991), but otherwise our procedures were identical to theirs. Briefly, a gas-phase mixture was prepared in a high-vacuum line and slowly deposited at 13 K on a pre-cooled, polished aluminum substrate in a vacuum chamber. An IR beam was passed through the resulting ice mixture (thickness ~ 4 μm) where it reflected off the underlying substrate, back through the ice mixture, and on to a detector. A spectrum of the blank substrate at 13 K served as the reference for spectral ratioing. Spectra were recorded as 60-scan accumulations with 4 cm^{-1} resolution. No corrections have been applied to the IR data for reflection and so the relative intensities of some of the features we report may be slightly different in transmission spectra, although the line positions should be the same. Most experiments were done with $\text{H}_2\text{O}:\text{CH}_3\text{OH} = 2:1$ mixtures or 2:1 mixtures with a few percent of other molecules added. Supporting experiments involved H_2O or CH_3OH alone, to examine the spectra of crystalline and amorphous phases of each compound. Radiation experiments were performed with 0.7 MeV protons from a Van de Graaff accelerator. Samples deposited at 77 K gave essentially the same results as those deposited at 13 K.

Our temperatures in the 100 K - 140 K region appear to be ~ 10

K higher than those of Blake *et al.* (1991) for the same phenomena, such as sublimation of H₂O ice. This is probably because our ices were formed, and the temperatures were measured, on a metal substrate whereas their experiments used non-metallic substrates (carbon, CsI, polyethylene).

III. RESULTS

Experiments usually began with deposition of a H₂O:CH₃OH = 2:1 mixture at 13 K over 30 min, the spectrum of a typical amorphous sample being shown in Figure 1a. Warming the sample to 130 K at 3 - 6 K min⁻¹ gave the spectrum of Figure 1b. Either holding the sample at 130 K for several hours or warming to 140 K for about 15 min, gave the spectrum shown in Figure 1c. When the sample then was cooled to 14 K, the lines marked by asterisks in Figure 1c sharpened and shifted slightly, and one line split into two components. The resulting spectrum is seen in Figure 2a. Spectra of amorphous (dashed line) and crystalline (solid line) CH₃OH and H₂O are given in the lower part of Figure 2 for 14 K. Careful comparison shows that the small, sharp feature near 230 cm⁻¹ (43 μm) in Figure 2a is probably due to crystalline H₂O (Figure 2c).

In all experiments, the lines (marked in Figure 1c) in the $\text{H}_2\text{O}:\text{CH}_3\text{OH} = 2:1$ mixtures grew in the spectrum simultaneously as the sample was warmed to 140 K, indicating a common origin. When the sample was warmed further to 150 K they disappeared simultaneously and the spectrum was replaced by that of polycrystalline H_2O (Figure 2c), indicating loss of CH_3OH .

Separate experiments investigated the influence of a third compound on the spectral changes in the Figures. Three sets of gas-phase $\text{H}_2\text{O}:\text{CH}_3\text{OH} = 2:1$ mixtures were made, with CO added in the first set, NH_3 in the second, and H_2CO in the third. Little changes in the spectra, from those in Figures 1 and 2, were seen when the added molecule's concentration was under ~5 percent (mole or molecule percent). However, above this value the lines indicated by asterisks in Figure 1 were substantially weaker if not completely absent. There appeared to be an inverse relation between the concentration of the third component (CO, NH_3 , or H_2CO) and the strength of the lines marked in Figure 1c.

Another set of experiments involved proton irradiation of ice samples. When the original amorphous $\text{H}_2\text{O}:\text{CH}_3\text{OH} = 2:1$ mixture was irradiated (incident fluence = 10^{14} p⁺ cm⁻², absorbed dose ≈ 1

eV/molecule) at 13 K, none of the lines indicated in Figure 1c appeared on subsequent warming. Other samples were warmed from 13 K to produce the lines in Figure 1c and then re-cooled to either 13 K or 77 K. Subsequent irradiation removed each of the indicated lines and a second stage of warming failed to bring them back.

IV. DISCUSSION

Our spectrum in Figure 2a shows six lines superimposed on a much broader underlying feature resembling amorphous CH_3OH and H_2O bands. The abrupt, irreversible production of the sharper lines on warming the originally amorphous $\text{H}_2\text{O}:\text{CH}_3\text{OH} = 2:1$ mixture is reminiscent of the changes seen on forming a crystalline phase in other amorphous materials, such as CH_3OH (Figure 2b), H_2O (Figure 2c), and CO_2 (Falk & Seto 1986). This suggests that the lines in Figure 2a, and those marked in Figure 1c for a higher temperature, also belong to a crystalline phase.

A comparison of Figure 2a for the $\text{H}_2\text{O}:\text{CH}_3\text{OH}$ mixture with the spectrum of crystalline CH_3OH in Figure 2b shows distinct differences. Attributing just some of the sharper lines from the mixture to crystalline CH_3OH would contradict the conclusion that

all of the features are evidently due to one chemical species since they increase together on warming the original condensate from 13 K to 130 K and decrease together on additional warming to 150 K. Furthermore, while it is conceivable that the 175 and 148 cm^{-1} bands may arise from crystalline CH_3OH formed when the clathrate hydrate structure develops, this explanation would contradict the electron diffraction result that only one crystalline phase was present under the same conditions, namely a clathrate hydrate (Blake *et al.* 1991). Since we have used the same experimental procedure as Blake *et al.*, we assign all of the sharp lines marked in Figure 1c and shown in Figure 2a to the type II CH_3OH clathrate hydrate. The positions of the clathrate hydrate's absorbances, and of others shown in the Figures, are summarized in Table 1.

Several lines of evidence support our assignment. First, adding a third molecule (CO , NH_3 , or H_2CO) to the $\text{H}_2\text{O}:\text{CH}_3\text{OH}$ mixture substantially weakened all the lines assigned to the clathrate hydrate. This can be attributed to the disruption of clathrate hydrate cages by the added (impurity) species. Second, when the original amorphous $\text{H}_2\text{O}:\text{CH}_3\text{OH} = 2:1$ mixture was irradiated before warming, none of the CH_3OH clathrate hydrate absorbances

appeared at 130 - 140 K, presumably because of the presence of radiation products such as H_2CO and CO . Third, when the original amorphous mixture was warmed to 140 K to form the clathrate, then recooled to either 13 K or 77 K and irradiated, all clathrate lines were lost, leaving a far-IR spectrum dominated by features of amorphous CH_3OH and amorphous H_2O , similar to Figure 1a. This is consistent with the known radiation-induced amorphization of crystalline materials, such as crystalline H_2O ice (Hudson & Moore 1992, Moore & Hudson 1992). Re-warming the irradiated sample failed to give the CH_3OH clathrate hydrate spectrum, again probably because of the radiation products.

Since the spectrum of the CH_3OH clathrate hydrate in Figure 2a has not been subjected to a complete theoretical analysis, and since deuterium substitution experiments have not been done, little can be said about each line's origin. However, it is likely that the spectrum is due to both inter- and intramolecular vibrations. The features at 274 and 214 cm^{-1} are probably due to intramolecular vibrations since similar lines are seen in the spectra of CH_3OH isolated in argon matrices (Barnes & Hallam 1970), but not in pure CH_3OH (Figure 2b). Their presence in our spectra is additional

evidence for enclathrated molecules. The feature at 326 cm^{-1} is probably the counterpart of the 353 cm^{-1} band in crystalline CH_3OH , which is due either to intermolecular $\text{O}\cdots\text{O}$ vibrations or $\text{O-H}\cdots\text{O}$ hydrogen-bond stretching (Wong & Whalley 1971; Durig *et al.* 1971). Assignments for the remaining features are less certain but probably they too are intermolecular in origin.

The overall appearance of our CH_3OH clathrate hydrate far-IR spectrum is quite different from clathrate hydrate spectra in which the guest molecule undergoes little or no hydrogen bonding with the H_2O cages, such spectra being strongly dominated by a broad absorption from the H_2O cages (Bertie & Solinas 1975). The presence of relatively sharp features from our annealed $\text{H}_2\text{O}:\text{CH}_3\text{OH} = 2:1$ mixture suggests that interaction of the trapped CH_3OH with the surrounding H_2O molecules occurs. This is supported by the electron diffraction data (Blake *et al.* 1991) and even earlier x-ray diffraction studies of ethanol ($\text{C}_2\text{H}_5\text{OH}$) clathrate hydrates (Boutron & Kaufman 1978). Although the diffraction data reveal formation of type II clathrate hydrates for CH_3OH and $\text{C}_2\text{H}_5\text{OH}$, it has been noted that some variations from the usual type II crystal structure probably exist (Blake *et al.* 1991; Boutron & Kaufman 1978; Potts & Davidson

1965). The nature and extent of these variations remain to be determined. Hydrogen bonding as well as hindered rotation of the methyl group or the entire molecule may be important.

Since there are almost certainly H₂O-dominated amorphous ice mixtures in space (Smoluchowski 1978, 1981, 1983; Leger *et al.* 1979; Hagen, Tielens, and Greenberg 1981), which conceivably could form clathrate hydrates on warming, it is worth summarizing CH₃OH clathrate formation, stability, and detection based on laboratory experience. Our results show that CH₃OH clathrate formation can be inhibited by the presence of "impurities," both those present initially and those formed by irradiation. In addition, ionizing radiation destroys the clathrate. Finally, there appears to be only a narrow range of temperature which produces the clathrate without forming polycrystalline H₂O ice.

These conditions aside, even if CH₃OH clathrate hydrate does form and exist in astronomical objects, it could be difficult to detect and identify. For example, on the surface of a cometary nucleus with a temperature below 150 K, certainly beyond the orbit of Mars, most of the observed energy in the 25 - 100 μ m (far-infrared) region should be from thermal emission. Locating such a nucleus, and acquiring

good quality spectra, would be challenging. Dense interstellar molecular clouds also could be searched in the far-IR for the CH₃OH clathrate hydrate. However interpretation of interstellar emissions or absorptions can involve tedious modeling. Nevertheless, the laboratory spectrum observed for the type II CH₃OH clathrate hydrate is sufficiently unique that its presence in the spectrum of an astronomical object may help determine the object's thermal and radiation history.

ACKNOWLEDGEMENTS

We acknowledge D. Blake, D. M. Hudgins, S. A. Sandford, A. G. G. M. Tielens, and L. J. Allamandola at NASA/Ames Research Center for preprints, conversations, and correspondence. S. A. Sandford is also thanked for copies of original spectra and discussions concerning this work. B. Donn is acknowledged for criticisms and conversations related to this research and R. Ferrante is thanked for assistance with mid-IR experiments. RLH acknowledges the Administration of Eckerd College for a leave during which this work was done under an Intergovernmental Personnel Act Agreement at the NASA/Goddard Space Flight Center. Both authors acknowledge NASA funding support through grant NSG 5172.

TABLE 1
PEAK POSITIONS IN FAR-IR SPECTRA

Material	Temperature (K)	Peak Positions (cm ⁻¹)
CH ₃ OH clathrate hydrate	14	326, 274, 214, 175, 154, 129
CH ₃ OH clathrate hydrate	140	317, 268, 209, 167, 125
Amorphous H ₂ O:CH ₃ OH 2:1	13	306 (broad shoulder), 206
Amorphous H ₂ O:CH ₃ OH 2:1	130	306 (broad shoulder), 216
Amorphous CH ₃ OH	14	318, 146
Crystalline CH ₃ OH	14	353, 175, 148, 113
Crystalline CH ₃ OH*	140	345, 163, 107
Amorphous H ₂ O	14	220
Amorphous H ₂ O*	130	217
Crystalline H ₂ O	14	230, 163
Crystalline H ₂ O*	140	223, 162

*Included for comparison purposes. Amorphous CH₃OH and amorphous H₂O are unstable at 140 K and convert to crystalline phases.

REFERENCES

- Barnes, A. J. & Hallam, H. E. 1970, *Trans. Farad. Soc.*, **66**, 1920

- Bertie, J. E. & Devlin, J. P. 1983, *J. Chem. Phys.*, **78**, 6340

- Bertie, J. E. & Jacobs, S. M. 1977, *Can. J. Chem.*, **55**, 1777

- Bertie, J. E. & Jacobs, S. M. 1978, *J. Chem. Phys.*, **69**, 4105

- Bertie, J. E. & Othen, A. D. 1972, *Can. J. Chem.*, **51**, 1159

- Bertie, J. E. & Othen, D. A. 1973a, *Can. J. Chem.*, **51**, 160

- Bertie, J. E. & Othen, A. D. 1973b, *Can. J. Chem.*, **51**, 3641

- Bertie, J. E. & Solinas, M. 1975, *Can. J. Chem.*, **53**, 2624

- Blake, D., Allamandola, L., Sandford, S., Hudgins, D., & Freund, F.
1991, *Science*, **254**, 548

- Bocklee-Morvan, D., Colom, P., Crovisier, J., Despois, D., Paubert, &
G. 1991, *Nature*, **350**, 318

- Boutron, P. & Kaufman, A. 1978, *J. Chem. Phys.*, **68**, 5032

- Davidson, D. W. 1973 in *Water: A Comprehensive Treatise*, ed. F. Franks (New York: Plenum, New York), vol. 2, chapter 3
- Delsemme, A. H. & Swings, P. 1952, *Ann. Astrophys.*, **15**, 1
- During, J. R., Pate, C. B., Li, Y. S., & Antion, D. J. 1971, *J. Chem. Phys.*, **54**, 4863
- Falk, M. & Seto, P. F. 1986, *Can. J. Spectros.*, **31**, 134
- Fleyfel, F. & Devlin, J. P. 1991, *J. Phys. Chem.*, **95**, 3811
- Grim, R. J. A., Baas, F., Geballe, T. R., Greenberg, J. M., & Schutte, W. 1991, *A&A*, **243**, 473
- Hagen, W., Tielens, A. G. G. M., & Greenberg, J. M. 1981, *Chem. Phys.*, **56**, 367
- Hoban, S., Mumma, M., Reuter, D. C., DiSanti, M., Joyce, R. R., & Storrs, A. 1991, *Icarus*, **93**, 122
- Hudson, R. L. & Donn, B. 1991, *Icarus*, **94**, 326
- Hudson, R. L. & Moore, M. H. 1992, *J. Phys. Chem.*, **96**, 6500

- Jeffrey, G. A. 1984 in Inclusion Compounds, ed. J. L. Atwood, J. E. D. Davies, and D. D. MacNicol (New York: Academic), vol. 1, chap. 5
- Leger, A., Klein, J., de Cheveigne, S., Guinet, C., DeFourneau, D., Belin, M. 1979, A&A, **79**, 256

- Moore, M. H. & Hudson, R. L. 1992, ApJ, in press
- Potts, A. D. & Davidson, D. W. 1965, J. Phys. Chem., **69**, 996

- Smoluchowski, R. 1978, Science, **201**, 809

- Smoluchowski, R. 1981, ApJ, **244**, L31

- Smoluchowski, R. 1983, Science, **222**, 161

- Smyth, W. D. 1975, Icarus, **24**, 421

- Tielens, A. G. G. M. & Allamandola, L. J. 1987 in Physical Processes in Interstellar Clouds, ed. G. Morfill and M. Schaler (Dordrecht: Reidel), pp. 333 - 338
- Wong, P. T. T. & Whalley, E. 1971, J. Chem. Phys., **53**, 1830

FIGURE CAPTIONS

Figure 1: Evolution of the far-IR spectrum of the $\text{H}_2\text{O}:\text{CH}_3\text{OH}$ mixture. (a) 13 K, (b) 130 K, (c) 140 K. Spectra have been displaced vertically for clarity. The initial ice had a thickness of $\sim 4 \mu\text{m}$. Features marked with asterisks in "c" sharpen and shift slightly upon recooling as shown in Figure 2a.

Figure 2: Far-IR spectra at 14 K. (a) $\text{H}_2\text{O}:\text{CH}_3\text{OH}$ mixture after warming from 13 K to 140 K and then recooling, (b) pure amorphous (dashed line) and pure crystalline (solid line) CH_3OH , (c) pure amorphous (dashed line) and pure crystalline (solid line) H_2O . The crystalline samples in (b) and (c) were made by briefly warming the amorphous solid to either 130 K (CH_3OH) or 155 K (H_2O) and then recooling. Spectra have been displaced vertically for clarity and scaled for a common sample thickness of $\sim 4 \mu\text{m}$.

POSTAL ADDRESSES

Dr. Reggie L. Hudson
Department of Chemistry
Eckerd College
St. Petersburg, FL 33733
Telephone: (813) 864-8457

Dr. Marla H. Moore
Astrochemistry Branch
NASA/Goddard Space Flight Center
Greenbelt, MD 20771
Telephone: (301) 286-9031

