

1  
17-10-02  
33016  
P-38

**Progress Report**

**December 1, 1990 - August 31, 1991**

**NASA Grant NAG 5-890**

by

**Shashi B. Verma, Principal Investigator**

**Department of Agricultural Meteorology  
Institute of Agriculture and Natural Resources  
University of Nebraska-Lincoln  
Lincoln, Nebraska 68583-0728**

**Phone: (402) 472-6702 or 472-3679**

(NASA-CR-182781) ANALYSIS OF FIELD MEASUREMENTS OF CARBON DIOXIDE AND WATER VAPOR FLUXES Progress Report, 1 Dec. 1990 - 31 Aug. 1991 (Nebraska Univ.) 28 p. (USCL 15)

NP1-30610

Unclas  
0034226

63/45

## TABLE OF CONTENTS

|   | <b>Page</b> |
|---|-------------|
| <b>1. SUMMARY OF ACTIVITIES</b> .....   | <b>2</b>    |
| <b>2. DETAILS OF WORK PERFORMED</b> .....   | <b>2</b>    |
| 2.1 Photosynthesis and Stomatal Conductance Related to Reflectance on the<br>Canopy Scale ..... | <b>2</b>    |
| 2.2 Further Data Processing and Analyses .....  | <b>13</b>   |
| <b>3. JOINT STUDIES WITH OTHER FIFE INVESTIGATORS</b> .....                                     | <b>14</b>   |
| <b>4. SCIENTIFIC PUBLICATIONS AND PRESENTATIONS STEMMING FROM<br/>    GRANT NAG5-890</b> .....  | <b>14</b>   |
| <b>5. RESEARCH PLANS</b> .....  | <b>16</b>   |
| <b>6. LIST OF REFERENCES</b> .....  | <b>17</b>   |
| <b>7. APPENDICES</b> .....  | <b>21</b>   |

## **1. SUMMARY OF ACTIVITIES**

During the past several months we have concentrated on further analysis of our field measurements of carbon dioxide and water vapor fluxes. These data were examined in conjunction with reflectances obtained from a helicopter mounted Modular Multiband Radiometer (MMR) at our grassland study site (#16) during 1987. These measurements are representative of the canopy scale (10-100 m)<sup>2</sup> and provide a good basis for investigating the hypotheses/relationships potentially useful in remote sensing applications.

All of the micrometeorological data collected at site 916 during FIFE-89 have been processed and fluxes of CO<sub>2</sub>, water vapor and sensible heat have been calculated. Soil CO<sub>2</sub> fluxes were estimated using the relationships developed at our site by Norman *et al.* (1991). Employing these soil CO<sub>2</sub> flux values, in conjunction with our micrometeorological measurements, canopy photosynthesis is being estimated.

A biochemical model of leaf photosynthesis was adapted to the prairie vegetation. The modeled leaf photosynthetic rates were scaled up to the canopy level. Further tests are underway employing our measurements made during FIFE-89. This model and a multiplicative stomatal conductance model are also used to calculate canopy conductance. Further evaluation of these calculations, using the FIFE-89 data, is in progress.

Our future plans include completion of these tests of our canopy photosynthesis and conductance models against our FIFE-89 measurements. Our FIFE-87 and FIFE-89 data will also be used to examine the contribution of this temperate grassland in northeastern Kansas as a source and sink for atmospheric CO<sub>2</sub> under a wide range of meteorological and soil moisture conditions.

## **2. DETAILS OF WORK PERFORMED**

### **2.1 Photosynthesis and Stomatal Conductance Related to Reflectance on the Canopy Scale**

#### **2.1.1 Introduction**

In recent years many researchers (*e.g.*, Monteith, 1977; Tucker *et al.*, 1981; Asrar *et al.*, 1984; Sellers, 1985, 1987) have investigated the relationships between linear combinations of reflected

visible and near infrared radiation known as vegetation indices, and biophysical quantities, such as the leaf area index, biomass, and the amount of photosynthetically active radiation (PAR) absorbed by the canopy. Two vegetation indices commonly used are:

$$\text{Simple Ratio Vegetation Index (SR)} = R_N/R_V \quad (1)$$

and

$$\text{Normalized Difference Vegetation Index (NDVI)} = (R_N - R_V)/(R_N + R_V) \quad (2)$$

where  $R_N$  and  $R_V$  are upwelling radiances or surface reflectances in the near-infrared and visible-red spectral regions, respectively.

Canopy conductance ( $g_c$ ) and canopy photosynthesis ( $P_c$ ) can be described using two simple multiplicative relationships (see Jarvis, 1976; Sellers, 1987):

$$g_c = g_c^* (PAR) \cdot [f(T) \cdot f(D) \cdot f(\theta_s)] \quad (3)$$

$$P_c = P_c^* (PAR) \cdot [f(T) \cdot f(D) \cdot f(\theta_s)] \quad (4)$$

where  $g_c^*(PAR)$  and  $P_c^*(PAR)$  are the unstressed values of  $g_c$  and  $P_c$ , regulated only by vegetation type, health and density and the amount of incoming PAR. The stress factors  $f(T)$ ,  $f(D)$ , and  $f(\theta_s)$  account for the effects of temperature (T), ambient vapor pressure deficit (D) and extractable soil water ( $\theta_s$ ). The "stress factors"  $f(x)$  vary between unity (optimal conditions) and zero (unfavorable conditions shutting down photosynthesis and transpiration). The stress factors operating on  $g_c$  and  $P_c$  are assumed to be identical and invariant within the canopy for a given set of environmental conditions. In this study, canopy level flux data and the leaf stomatal conductance of the dominant grass species measured at the site were used to obtain the functional forms of the expressions for the stress factors

(see Appendix A). The derived coefficients for these stress factors are independent of the time-varying leaf area index.

The terms  $g_c^*$  and  $P_c^*$  are dependent on the vegetation canopy and the incident PAR flux. Prior analyses (*e.g.*, Sellers, 1987) have indicated that  $g_c^*$  and  $P_c^*$  should increase almost linearly with PAR and, therefore,  $\partial g_c^*/\partial PAR$  and  $\partial P_c^*/\partial PAR$  can be treated as conservative surface properties. Put simply, a bare soil surface should show no change in conductance or "photosynthesis" with increasing PAR and so its associated partial derivatives of  $g_c^*$  and  $P_c^*$  would be zero. A thickly vegetated surface, on the other hand, would exhibit a strong sensitivity to PAR (non-zero partial derivatives of  $g_c^*$  and  $P_c^*$ ).

The theoretical study of Sellers (1987) proposed that for ideal conditions SR should be near-linearly related to the absorbed PAR (APAR) and to the derivatives of  $g_c^*$  and  $P_c^*$  with respect to PAR:

$$SR \propto APAR, \frac{\partial g_c^*}{\partial PAR}, \frac{\partial P_c^*}{\partial PAR} \quad (5)$$

The relationship described in Eq. (5) should hold where (a) the canopy consists of spatially uniform cover of green leaves, (b) the reflectance of the background and understory in the chlorophyll absorption band is approximately equal to that of the overlying vegetation and (c) the wavebands of the sensing instrument are configured so that the perceived scattering coefficients of the leaves conform to:

$$\omega_N = 1 - \left[ \frac{G(\mu)}{2\mu} \right]^2 (1 - \omega_\pi), \quad (6)$$

where  $\omega_N$  and  $\omega_\pi$  are the leaf scattering coefficients in the near infrared and PAR wavelength intervals, respectively,  $G(\mu)$  is the relative projected area of leaf elements in the direction of  $\cos^{-1} \mu$  and  $\mu$  is the cosine of the zenith angle of the incident radiation flux (see Sellers, 1987).

During 1987, the First ISLSCP Field Experiment (FIFE) was conducted in a 15 km x 15 km area near Manhattan, Kansas. In this paper we report on an analysis of the surface flux and surface reflectance measurements made at a prairie site in the FIFE study area. The fluxes of water vapor and carbon dioxide were measured using the micrometeorological eddy correlation technique (see *e.g.*, Kaimal, 1975; Kanemasu *et al.*, 1979; Verma, 1990), and the reflectance measurements were made with a helicopter-mounted Barnes Modular Multiband Radiometer (MMR). The effective narrow-band visible and near-infrared scattering coefficients for Landsat TM bands 3 and 4, which correspond to the bands on the helicopter MMR, were calculated to be 0.12-0.18 and 0.95-0.97 respectively for individual grass leaves from the data of Walter-Shea (personal communication) collected during FIFE. For calculations involving the penetration of broad-band PAR (0.4 - 0.7  $\mu\text{m}$ ) into the canopy we have assumed<sup>1</sup>  $\omega_x = 0.2$ , following the calculation of Sellers (1987). Equation (6) is, therefore, approximately satisfied for a spherical leaf angle distribution ( $G(\mu)=0.5$ ) and so we might expect that Eq. (5) should hold reasonably well for the prairie site being studied here. The purpose of this paper is to test the hypothesis represented in this equation with the field measurements collected in this study. The data presented here are representative of canopy scale and, therefore, can be considered as a test case for investigating the use of Eq. (5) for remote sensing applications.

## 2.1.2 Materials and Methods

### 2.1.2.1 Site

The study was conducted in 1987 at a tallgrass prairie site (39° 03' N, 96° 32' W, 445 m above m.s.l., FIFE site #16) near Manhattan, Kansas. The soil is predominantly Dwight silty clay loam (Typic Natrustolls). The vegetation is dominated by three C<sub>4</sub> grass species: big bluestem (*Andropogon gerardii*), indiagrass (*Sorghastrum nutans*) and switchgrass (*Panicum virgatum*). The prairie was burned in the spring of 1987. The experimental area had been lightly grazed for several years, but was not grazed in 1986 and 1987.

---

<sup>1</sup>Values of  $\omega_x$  derived from the observations of E. A. Walter-Shea (personal communication) vary from 0.17 to 0.19.

### 2.1.2.2 Micrometeorological Measurements

Fluxes of carbon dioxide, water vapor, sensible heat and momentum were measured with eddy correlation sensors mounted 2.25 m above the ground. The sensors included: a three-dimensional sonic anemometer (Kaijo Denki Co., Tokyo, Japan, Model DAT-310), one-dimensional sonic anemometers (Kaijo Denki Co., Tokyo, Japan, Model DAT-110; Campbell Scientific, Logan, Utah, Model CA27), fine wire (0.025 mm) thermocouples (Campbell Scientific, Logan, Utah, Model 127), a Lyman-alpha hygrometer (A.I.R., Inc., Boulder, CO, Model LA-1) with a 5 mm path length and a rapid response CO<sub>2</sub> sensor with a 200 mm path length (Bingham *et al.*, 1978; Anderson *et al.*, 1984). Net radiation, PAR, soil heat flux, air temperature and humidity were also measured. Details on the instrumentation and procedure can be found in Kim and Verma (1990a & b). The configuration of the flux instrumentation indicates that an area extending up to (100 m)<sup>2</sup> upwind of the instrumentation would have the strongest (~70%) influence on the measured flux (see *e.g.* Schuepp *et al.*, 1990).

### 2.1.2.3 Plant and Soil Measurements

Stomatal conductance was measured on selected leaf blades of tillers of big bluestem, indiagrass and switchgrass, employing a steady-state porometer. The surface soil water content (0-0.1 m) was monitored gravimetrically almost every day and the subsurface water content (0.1-1.4 m) was measured with a neutron probe on a weekly basis. Leaf area index was measured generally every two weeks.

### 2.1.2.4 Reflectance Measurements

Measurements of surface reflectance were made with a helicopter-mounted MMR from 230 m above the ground. The MMR has eight spectral channels, however only two channels were used in the analysis: 0.63 - 0.68  $\mu\text{m}$  (MMR3 or visible-red) and 0.75 - 0.88  $\mu\text{m}$  (MMR4 or near infrared). A second MMR was positioned on the surface to measure reflected radiance from a Barium Sulfate reference panel as a measure of irradiance. These reference panel measurements were made at one

minute intervals starting thirty minutes before the helicopter missions and ending 20 to 30 minutes after the helicopter left the FIFE study area. The reference panel was calibrated using procedures outlined in Jackson *et al.* (1987), and the MMR was calibrated using procedures described in Markham *et al.* (1988), thus allowing determination of absolute radiance from the MMR voltage responses.

A minimum of 25 and a maximum of 150 measurements were taken over the study site within 100 m of the flux station during each mission. Two to five minutes were required to obtain these measurements. Data were collected under clear sky conditions one or two times a day. The sampled areas within the site were usually concentrated upwind of the micrometeorological flux sensors. The vegetation across the site was fairly uniform, with little variation in topography.

The raw digital count values from the MMR were converted to sensor radiances using the calibration data taken prior to the experiment. The sensor radiances were then corrected for atmospheric effects using an atmospheric correction algorithm developed for FIFE by Fraser *et al.* (1989) based on the Dave (1978) algorithm that includes the effects of multiple scattering in the atmosphere and between the atmosphere and the surface. Aerosol optical depths were determined from a Sun-tracking photometer, operating in discrete bands in the 380 - 1020 nm region (Wrigley *et al.*, 1990 and Halthore *et al.* 1990) during the helicopter data acquisition. Gaseous absorption terms ( $\text{CO}_2$ ,  $\text{O}_3$ ,  $\text{H}_2\text{O}$ , etc.) were calculated using Lowtran-7 (Kniezys, 1988) from radiosonde atmospheric water-vapor profiles taken simultaneously with the helicopter data acquisition. The SR was then calculated for the site using the average site reflectance in MMR3 and MMR4.

#### 2.1.2.5 Estimating Canopy Conductance

Canopy stomatal conductance ( $g_c$ ) can be estimated from the surface conductance,  $g_c(\text{PM})$  [canopy conductance and surface conductance are roughly equivalent as the soil evaporation tends to zero - see *e.g.*, Monteith, 1965; Thom, 1975; Baldocchi *et al.*, 1991, Kim and Verma, 1991]. Values of  $g_c(\text{PM}) = 1/r_c(\text{PM})$  were calculated from the measured fluxes employing the Penman-Monteith equation (Monteith, 1965; Jarvis *et al.*, 1976):



$$r_c(PM) = [(s/\gamma)\beta - 1] r_a + (\beta + 1) (\rho C_p/\gamma) (D/A), \quad (7)$$

where  $s$  is the slope of the saturated vapor pressure-temperature curve,  $\gamma$  is the psychrometric constant,  $\beta$  is the Bowen ratio ( $=H/\lambda E$ ),  $H$  is the sensible heat flux,  $\lambda E$  is the latent heat flux,  $\rho$  is the density of air,  $C_p$  is the specific heat of air,  $D$  is the vapor pressure deficit,  $A (=R_n + G)$  is the available energy,  $G$  is the soil heat flux and  $r_a$  is the aerodynamic resistance to water vapor/heat transfer. The resistance,  $r_a$  can be given by:

$$r_a = r_{am} + r_b, \quad (8)$$

where  $r_{am}$  is the aerodynamic resistance to momentum transfer (see Appendix B for details). The term  $r_{am}$  can be expressed as:

$$r_{am} = \bar{U}/u_*^2, \quad (9)$$

where  $\bar{U}$  is the mean horizontal wind speed at a reference height  $z$  and  $u_*$  is the friction velocity. The excess resistance term,  $r_b$  (see *e.g.*, Thom, 1972; Wesely and Hicks, 1977), is approximated by:

$$r_b = (2/ku_*) (\kappa/D_v)^{2/3}, \quad (10)$$

where  $k$  is von Karman's constant,  $\kappa$  is the thermal diffusivity and  $D_v$  is the molecular diffusivity of water vapor. Friction velocity data from the three-dimensional sonic anemometer signals were used to compute  $r_a$ . Values of  $g_c(PM)$  were calculated using  $r_a$  and measured fluxes of  $R_n$ ,  $G$ ,  $H$  and  $\lambda E$  in Eq. (7).

#### 2.1.2.6 Calculation of Canopy Photosynthesis ( $P_c$ )

The eddy correlation measurement of atmospheric  $\text{CO}_2$  flux ( $F_c$ ) provides the sum of canopy and soil  $\text{CO}_2$  fluxes. An estimation of soil  $\text{CO}_2$  flux is necessary for computation of canopy photosynthesis ( $P_c$ ). Soil  $\text{CO}_2$  flux was not directly measured in this study. However, our nocturnal  $F_c$  data provided information on the respiratory release of  $\text{CO}_2$  from plant and soil. Plant respiration ( $R_d$ ) was estimated by the following relationship (Polley *et al.*, 1991):

$$R_d = R_d (@ 25^\circ\text{C}) [\exp \{E_1 (T_l - 25)/(298 R T_l)\}] L_T \quad (11)$$

where  $E_1$  is the activation energy = 45,000 J mol<sup>-1</sup>,  $T_l$  and  $T_k$  are the leaf temperature in °C and °K, respectively,  $R$  is the gas constant, and  $L_T$  is the leaf area index. In our computations, the air temperature near the top of the canopy was used to approximate the leaf temperature. Following Polley *et al.* (1991), a value of 1.55  $\mu\text{ mol m}^{-2}$  (leaf area) s<sup>-1</sup> was used for  $R_d$  (@ 25°C). Plant respiration, estimated by Eq. (11), was subtracted from the nocturnal  $F_c$  to obtain (nocturnal) soil  $\text{CO}_2$  flux. The soil  $\text{CO}_2$  flux values were then adjusted to the daytime soil temperature (0-50 mm) using a  $Q_{10}$  factor of 2 (Kucera and Kirkham, 1971) to calculate the daytime soil  $\text{CO}_2$  flux. The daytime soil  $\text{CO}_2$  flux was used in conjunction with the  $F_c$  to estimate canopy photosynthesis ( $P_c$ ). To minimize the effect of changing soil moisture and photosynthate on soil  $\text{CO}_2$  flux, the nocturnal  $F_c$  data used here were generally limited to those from the preceding or the following nights.

### 2.1.3. Results and Discussion

#### 2.1.3.1 Soil Moisture and Plant Growth

A seasonal picture of soil moisture and plant growth at the study site is presented in Fig. 1. Except for a three-week period in late July-early August, precipitation was generally ample during the study period. During the dry period, the extractable soil water,  $\theta_e$  (0-1.4 m), reached as low as 25% and moisture stress conditions prevailed. Values of green leaf area index (LAI) and stages in the annual life cycle of the prairie vegetation are also included in Fig. 1.

### 2.1.3.2 Seasonal Variations in Canopy Surface Conductance, Canopy Photosynthesis and Simple Ratio Vegetation Index

In Fig. 2, an overall seasonal pattern<sup>2</sup> of canopy conductance, computed from our measured surface fluxes using Eqs. (7-10), is presented. During the most vigorous stage of plant growth (June and July), the midday  $g_c$  averaged about  $15.0 \text{ mm s}^{-1}$ . The dry period during late July-early August produced much lower value of  $g_c$  with an average of  $3 \text{ mm s}^{-1}$ . With ample rainfall in mid-August, the vegetation began to recover from moisture stress, resulting in an increase in  $g_c$ . In mid-late August, the midday  $g_c$  averaged about  $9.0 \text{ mm s}^{-1}$ . Later in the season  $g_c$  decreased as the prairie vegetation senesced. In October the midday  $g_c$  averaged about  $1.3 \text{ mm s}^{-1}$ .

The seasonal variation<sup>3</sup> of canopy photosynthesis ( $P_c$ ) is given in Fig. 3. The midday  $P_c$  reached a seasonal peak of about  $1.5 \text{ mg m}^{-2} (\text{ground area}) \text{ s}^{-1}$  during June-early July, the most vigorous stage of plant growth. There was a drastic reduction in  $P_c$  during the dry period. The value of  $P_c$  reached a minimum of near zero around 30 July, a day with severe moisture stress. The soil moisture conditions improved during mid-August, and the midday  $P_c$  increased to about  $0.75 \text{ mg m}^{-2} \text{ s}^{-1}$ . The photosynthetic rates diminished rapidly as the vegetation senesced. The midday value of  $P_c$  in early to mid-October was about  $0.1 \text{ mg m}^{-2} \text{ s}^{-1}$ .

The days on which SR was measured are indicated on Figs. 2 and 3. The seasonal progression of SR for the study site is shown in Fig. 4. Values of SR were high, ranging from 8.5 to 9.4, in June-early July when the plants were growing vigorously. During the early senescence stage which followed the dry period in late July-early August (see Fig. 1), the SR values reduced to 5.1-6.2. The lowest values ( $\approx 2$ ) of SR were observed during mid-October toward the end of the senescence period.

### 2.1.3.3 Simple Ratio Vegetation Index, Related to Unstressed Canopy Conductance and Unstressed Canopy Photosynthesis

The procedure described in Appendix A was employed to define the form and coefficients for the stress factors,  $f(x)$ , used in Eqs. 3 and 4. Measured values of  $D$  and  $\theta_s$  were used to calculate

---

<sup>2</sup>Details of these results have been reported in Kim and Verma, 1990a.

<sup>3</sup>Details of these results have been reported in Verma *et al.*, 1991.

the stress factors  $f(D)$  and  $f(\theta_s)$ . The values of  $g_c$  and  $P_c$  were then divided by the stress factors to estimate  $g'_c$  and  $P'_c$  (see Eqs. 3 and 4). Plots of  $g'_c$  and  $P'_c$  against PAR are shown in Figs. 5 and 6. Estimates of  $\partial g'_c/\partial \text{PAR}$  and  $\partial P'_c/\partial \text{PAR}$  were obtained by applying simple linear fits to these data (only afternoon data were used in estimating  $\partial g'_c/\partial \text{PAR}$  to minimize confounding effects of rapid time rate of change in soil evaporation). The values of  $\partial g'_c/\partial \text{PAR}$  and  $\partial P'_c/\partial \text{PAR}$  are shown in Table 1. The correlation coefficient ( $r^2$ ) was generally quite high, except on 4 June, 11 August and 8 October. Because of rain on 2 and 3 June, our analysis of data on 4 June is subject to problems arising from a rapid rate of change in soil evaporation. 11 August was a day with the driest soil moisture conditions of all the days during June-August when  $g'_c$  and  $P'_c$  were computed. Our empirical approach to calculate the stress factors under such severe moisture stress conditions is not adequate. The low  $r^2$  on 8 October is likely because the canopy conductance and photosynthetic rates diminished rapidly as vegetation senesced in October.

On five days (16 and 17 August and 7, 10 and 13 October), the canopy photosynthesis and conductance measurements were not available. The  $\partial g'_c/\partial \text{PAR}$  and  $\partial P'_c/\partial \text{PAR}$  values on 16 and 17 August were interpolated from the values on 15 and 20 August. On 7, 10 and 13 October, the values of  $\partial g'_c/\partial \text{PAR}$  and  $\partial P'_c/\partial \text{PAR}$  were assumed to be the same as those on 8 October.

In Figs. 7 and 8, SR is plotted against  $\partial g'_c/\partial \text{PAR}$  and  $\partial P'_c/\partial \text{PAR}$ , respectively. As indicated above, the data included in these two figures are from a range of soil moisture and meteorological conditions and cover different stages in the annual life cycle of the prairie vegetation (Table 1). The data from 4 June-11 July are from a period when the soil water was not limiting and the plants were growing vigorously. The data point from 11 August is during the dry period, when severe moisture stress conditions prevailed. With about 67 mm of rainfall on 13 August, the soil water conditions improved significantly. The data collected during 15-20 August represent the improved soil water conditions. The data during 7-13 October are from the senescence period.

Whereas the SR and canopy biophysical variables ( $\partial g'_c/\partial \text{PAR}$ ,  $\partial P'_c/\partial \text{PAR}$ ) track each other fairly well, the same cannot be said of the reported green LAI values and these three quantities. One

reason<sup>4</sup> may be that the canopy was much less lush in August than in June-July, but the bulk of the foliage was still classified as "green" rather than brown in August. This would explain the lower SR values in August (in spite of larger values reported for green LAI) and also the lower values of  $\partial g_c^* / \partial \text{PAR}$  and  $\partial P_c^* / \partial \text{PAR}$ .

A linear regression of all of the SR- $\partial P_c^* / \partial \text{PAR}$  data yielded a  $r^2$  of 0.41 (Fig. 8). Similarly, a linear regression of all the data in the SR- $\partial g_c^* / \partial \text{PAR}$  relationship (Fig. 7) resulted in  $r^2 = 0.56$ . Our representation of the effect of moisture stress on  $g_c^*$  and  $P_c^*$  is inadequate on 11 August when extreme dry soil moisture conditions prevailed. Also, as mentioned above, the value of  $\partial g_c^* / \partial \text{PAR}$  on 4 June was probably affected by problems associated with a rapid change in soil evaporation. Excluding the data points on 4 June and 11 August a linear regression of the SR- $\partial g_c^* / \partial \text{PAR}$  data (solid line shown in Fig. 7) resulted in  $r^2 = 0.69$ . Similarly, excluding the data point on 11 August, a linear regression of the SR- $\partial P_c^* / \partial \text{PAR}$  data (solid line shown in Fig. 8) yielded  $r^2 = 0.57$ .

Even though there is some scatter in the data, Figs. 7 and 8 seem to support the hypothesis suggested by Sellers (1987) that there should be near-linear relationships between the simple ratio vegetation index (SR) and the response of the unstressed canopy conductance and the photosynthetic capacity to the incident PAR (Eq. 5). The data scatter in Figs. 7 and 8 could have resulted from several simplifying assumptions used in the analysis presented above. First, there was no allowance for soil evaporation in the calculation of  $g_c(\text{PM})$ ; it was assumed that all of the latent heat flux was due to canopy transpiration. Second, the soil respiration was not measured directly, but was estimated using empirical relationships. Third, the expressions used to calculate the environmental stress factors are empirical and need further evaluation.

The scatter in Figs. 7 and 8 is consistent with expectations given the experimental methodologies and simple analytical tools we have applied. Further work is needed to address the problems outlined above to bring us closer to the goal of using satellite data to provide information on energy and mass exchanges in various terrestrial ecosystems.

---

<sup>4</sup>To estimate LAI, leaves of eight 0.18m<sup>2</sup> area were harvested. The harvested foliage was divided into live and dead components, and then fed through an area meter. However, this procedure took no account of changes in the greenness/lushness of live vegetation and in August the leaves were less green than in June/July.

#### **2.1.4. Summary and Concluding Remarks**

The surface fluxes of carbon dioxide and water vapor were measured employing the micrometeorological eddy correlation technique during FIFE in 1987. Concurrent measurements were also made of reflectance using a helicopter-mounted Barnes MMR, from which the simple ratio vegetation index (SR) was calculated. Both of these data sets provide information on the grassland canopy at a comparable scale, (10-100 m)<sup>2</sup>.

Using our data, both on the leaf level (*e.g.*, stomatal conductance of the three dominant species) and the canopy level (*e.g.*, water vapor flux), empirical relationships for the environmental stress factors,  $f(D)$  and  $f(\theta_s)$ , were developed to account for the effect of ambient vapor pressure deficit ( $D$ ) and extractable soil water ( $\theta_s$ ). These factors were then used in conjunction with the measured surface fluxes to compute unstressed canopy conductance ( $g_c'$ ) and unstressed canopy photosynthesis ( $P_c'$ ).

The derivatives ( $\partial g_c' / \partial \text{PAR}$  and  $\partial P_c' / \partial \text{PAR}$ ) of unstressed canopy conductance and unstressed canopy photosynthesis are dependent on the amount and type of green vegetation, and therefore, should be amenable to remote sensing. The analysis of Sellers (1987) indicated that  $\partial g_c' / \partial \text{PAR}$  and  $\partial P_c' / \partial \text{PAR}$  should be near-linearly related to SR and the first order, preliminary analysis of data presented in this paper seems to support this hypothesis. However, before such relationships can be employed in satellite remote sensing applications, more work is needed. Future studies should include: (a) adequate allowance of soil evaporation in calculating canopy conductance, (b) direct measurement of soil respiration, (c) further development and evaluation of environmental stress functions and (d) more frequent measurements of canopy reflectance under similar illumination conditions.

#### **2.2 Further Data Processing and Analyses**

All of the eddy correlation data collected at site 916 during 1989 have been processed and fluxes of energy and mass have been calculated. Supporting meteorological, soil water and plant growth data have also been computed.

Soil CO<sub>2</sub> flux has been estimated using the relationships developed at our site by Norman *et al.* (1991). Employing these estimates in conjunction with our eddy correlation measurements during FIFE-89, canopy photosynthesis of the prairie vegetation is being estimated. A biochemical model of leaf photosynthesis was applied to the tallgrass prairie. The modeled photosynthetic rates of individual leaves were scaled up to the canopy level using a radiative transfer model. The FIFE-89 data are now being used to further test the canopy photosynthesis model. This model as well as a "multiplicative" stomatal conductance model are employed to evaluate canopy conductance. Further tests of these models using the FIFE-89 data are underway.

### 3. JOINT STUDIES WITH OTHER FIFE INVESTIGATORS

The following includes a list of major joint studies presently underway:

- Relating surface fluxes to reflectance values measured by the helicopter-mounted multiband radiometer (MMR)—with Dr. C. Walthall, University of Maryland, College Park, MD, Drs. P. Sellers and F. Hall, NASA-Goddard Space Flight Center, Greenbelt, MD.
- Comparison of canopy surface conductance at two contrasting sites within the FIFE study area—with Dr. J. B. Stewart, Institute of Hydrology, Wallingford, Oxon, United Kingdom.
- Intercomparison of rapid response CO<sub>2</sub> sensors—with Dr. M. Wesely, Argonne National Laboratory, Argonne, IL and Dr. J. Moncrieff, University of Edinburgh, Edinburgh, U.K.

### 4. SCIENTIFIC PUBLICATIONS AND PRESENTATIONS STEMMING FROM GRANT NAG5-890

The following scientific publications and presentations stem from our work on Grant No. NAG5-890.

#### Journal Articles

Hall, F. G., P. J. Sellers, I. McPherson, R. D. Kelly, S. B. Verma, R. Markham, B. L. Blad, J. Wang and D. E. Strebel. 1989. FIFE: Analysis and Results—A Review. *Advances in Space Research*. Cospar Publication 9:275-293.

- Kim, J. and S. B. Verma. 1990. Components of surface energy balance in a temperate grassland ecosystem. *Boundary-Layer Meteorology* 51:401-417.
- Kim, J. and S. B. Verma. 1990. Carbon dioxide exchange in a temperate grassland ecosystem. *Boundary-Layer Meteorology* 52:135-149.
- Verma, S. B. 1990. Micrometeorological methods for measuring surface fluxes of mass and energy. In: Instrumentation for Studying Vegetation Canopies for Remote Sensing in Optical and Thermal Infrared Regions, N. S. Goel and J. M. Norman (co-editors), Special Issue of *Remote Sensing Reviews* 5(1):99-115.
- Kim, J. and S. B. Verma. 1991. Modeling canopy stomatal conductance in a temperate grassland ecosystem. *Agric. and Forest Meteorol.* 55:149-166.
- Kim, J. and S. B. Verma. 1991. Modeling canopy photosynthesis: Scaling up from a leaf to canopy in a temperate grassland ecosystem. *Agric. and Forest Meteorol.* (accepted for publication).
- Verma, S. B., J. Kim and R. J. Clement. 1991. Momentum, water vapor and carbon dioxide exchange at a centrally located site in FIFE. *J. Geophys. Res.* (submitted).
- Stewart, J. B. and S. B. Verma. 1991. Comparison of surface fluxes and conductances at two contrasting sites within FIFE area. *J. Geophys. Res.* (submitted).
- Moncrieff, J. B., S. B. Verma and D. R. Cook. 1991. Intercomparison of eddy correlation carbon dioxide sensors during FIFE-89. *J. Geophys. Res.* (submitted).
- Norman, J. M., R. L. Garcia and S. B. Verma. 1991. Soil CO<sub>2</sub> fluxes and the carbon budget of a grassland. *J. Geophys. Res.* (submitted).
- Smith, E. A., A. Y. Hsu, W. L. Crosson, R. T. Field, L. J. Fritschen, R. J. Gurney, E. T. Kanemasu, W. P. Kustas, D. Nie, W. J. Shuttleworth, J. B. Stewart, S. B. Verma, H. L. Weaver and M. L. Wesely. 1991. Area averaged surface fluxes and their time-space variability over the FIFE experimental domain. *J. Geophys. Res.* (submitted).

### Conference Presentations

- Verma, S. B. 1988. Carbon dioxide and water vapor fluxes over a tallgrass prairie during FIFE. Invited Paper. American Geophysical Union Spring Meeting, Baltimore, MD, May 18-20, 1988.
- Sellers, P. J., F. G. Hall, D. E. Strebel, R. D. Kelly, S. B. Verma, B. L. Markham, B. L. Blad, D. S. Schimel, J. R. Wang, W. Brutsaert, R. Desjardins, L. Fritschen, R. Frouin, R. Grossman, E. Kanemasu, I. McPherson, J. L. Dorman and C. Walthall. 1988. First ISLSCP Field Experiment: Experiment Execution and Preliminary Analyses. European Center for Medium Range Weather Forecasts (ECMWF) Symposium on land surface parameterizations, Proceedings, October 1988. Shinfield, U.K.
- Kim, J. and S. B. Verma. 1989. Energy and CO<sub>2</sub> Fluxes in a Grassland Ecosystem. Presented at the 19th Conference on Agricultural and Forest Meteorology of the American Meteorological Society, Charleston, SC, March 7-10, 1989.
- Norman, J. M., W. Polley, E. A. Walter-Shea, S. B. Verma, T. Arkebauer and D. Greigor. 1989. Comparison of measured and modeled fluxes of water and carbon dioxide at Konza Prairie.



Presented at the 19th Conference on Agricultural and Forest Meteorology of the American Meteorological Society, Charleston, SC, March 7-10, 1989.

- Stewart, J. B. and S. B. Verma. 1989. Comparison of aerodynamic and surface conductance at two contrasting sites within FIFE area. Presented at the Global Energy and Water Fluxes Symposium, Fifth Scientific Assembly of the International Association of Meteorology and Atmospheric Physics (IAMAP), Reading, U.K., July 31-August 12, 1989.
- Kanemasu, E. T., S. B. Verma, L. J. Fritschen and R. J. Gurney. 1990. Design and operation: flux measurements in FIFE. Presented at the American Meteorological Society Symposium on the First ISLSCP Field Experiment (FIFE), 70th AMS Annual Meeting, February 7-9, 1990, Anaheim, CA.
- Norman, J. M., S. B. Verma and R. Garcia. 1990. Soil surface CO<sub>2</sub> fluxes in the Konza Prairie. Presented at the American Meteorological Society Symposium on the First ISLSCP Field Experiment (FIFE), 70th AMS Annual Meeting, February 7-9, 1990, Anaheim, CA.
- Stewart, J. B. and S. B. Verma. 1990. Comparison of surface conductance at two contrasting sites within FIFE study area. Presented at the American Meteorological Society Symposium on the First ISLSCP Field Experiment (FIFE), 70th AMS Annual Meeting, February 7-9, 1990, Anaheim, CA.
- Verma, S. B., J. Kim and R. J. Clement. 1990. Fluxes of momentum, sensible heat, water vapor and CO<sub>2</sub> over a tallgrass prairie. Presented at the American Meteorological Society Symposium on the First ISLSCP Field Experiment (FIFE), 70th AMS Annual Meeting, February 7-9, 1990, Anaheim, CA.
- Stewart, J. B. and S. B. Verma. 1991. Can the simple ratio be used to estimate evaporation from the FIFE area? To be presented at the Special Session on Hydrometeorology of the American Meteorological Society, September 10-13, 1991, Salt Lake City, UT.
- Kim, J. and S. B. Verma. 1991. Modeling canopy photosynthesis in a temperate grassland ecosystem. To be presented at the 20th Conference on Agricultural and Forest Meteorology of the American Meteorological Society, September 10-13, 1991, Salt Lake City, UT.

## **5. RESEARCH PLANS**

Estimation of canopy photosynthesis during the measurement period in FIFE-89 will be completed. Further testing of canopy photosynthesis model, presently underway, will be completed in the next few months. Results from the 1989 stomatal conductance measurements will be incorporated into a multiplicative canopy conductance model. This model will also be further tested with FIFE-89 data.

Our micrometeorological measurements during FIFE-87 and FIFE-89 will be used to calculate daily net ecosystem CO<sub>2</sub> exchange. These results will be used to examine the contribution of this temperate grassland as a source and sink for atmospheric CO<sub>2</sub> under a wide range of meteorological and soil moisture conditions.

## 6. LIST OF REFERENCES

- Anderson, D. E., Verma, S. B., and Rosenberg, N. J. (1984), Eddy correlation measurements of CO<sub>2</sub>, latent heat and sensible heat fluxes over a crop surface, *Boundary-Layer Meteorol.* 29:263-272.
- Asrar, G., Fuchs, M., Kanemasu, E. T., and Hatfield, J. L. (1984), Estimating absorbed photosynthetic radiation and leaf area index from spectral reflectance in wheat, *Agron. J.* 76:300-306.
- Baldocchi, D. D., Luxmoore, R. J., and Hatfield, J. L. (1991), Discerning the forest from the trees: an essay on scaling canopy stomatal conductance, *Agric. and Forest Meteorol.* 54:197-226.
- Bingham, G. E., Gillespie, C. H., and McQuaid, J. H. (1978), Development of a miniature, rapid response CO<sub>2</sub> sensor. Report UCRL-52440, Lawrence Livermore National Laboratory.
- Brutsaert, W. (1984), *Evaporation in to the Atmosphere*, D. Reidel Publ. Co., Dordrecht.
- Chamberlain, A. C. (1966), Transport of gases to and from grass and grass-like surfaces, *Proc. Roy. Soc. A.* 290:236-265.
- Chamberlain, A. C. (1968), Transport of gases to and from surfaces with bluff and wave-like roughness elements, *Quart. J. Roy. Meteorol. Soc.* 94:318-332.
- Dave, J. V., (1978), "Extensive data sets of diffuse radiation in realistic atmospheric models with aerosols and common absorbing gases", *Solar Energy* 21:361-369.
- Fraser, R. S., Ferrare, R. A., Kaufman, Y. J., and Mattoo, S., (1989), "Algorithm for atmospheric corrections of aircraft and satellite imagery", NASA Technical Memorandum 100751, 106 pp.
- Garratt, J. R., and Hicks, B. B. (1973), Momentum, heat and water vapour transfer to and from natural and artificial surfaces, *Quart. J. Roy. Meteorol. Soc.* 99:680-687.
- Halthore, R., Brugge, C., and Markham, B., (1990), "Aerosol Optical thickness measurements during FIFE '89", pp 121-125, Preprint volume of the symposium on the First ISLSCP Field Experiment (FIFE), Feb 7-9, 1990, Anaheim, Calif., AMS, Boston, Mass.
- Jackson, R. D., Moran, M. S., Slater, P. N., and Biggar, S. F. (1987), Field calibration of reference reflectance panel, *Remote Sens. Environ.* 25:145-158.

- Jarvis, P. G. (1976), The interpretation of the variations in leaf water potential and stomatal conductance found in canopies in the field, *Phil. Trans. Roy. Soc. London*, Ser. B. 273:593-610.
- Jarvis, P. G., James, G. B., and Landsberg, J. J. (1976), Coniferous forest, *In Vegetation and the Atmosphere, Vol. 2, Case Studies* (J. L. Monteith, Ed.), Academic Press, New York, pp. 171-240.
- Kaimal, J. C. (1975), Sensors and techniques for direct measurement of turbulent fluxes and profiles in the atmospheric surface layer, *Atmospheric Technology* 7:7-14.
- Kanemasu, E. T., Wesely, M. L., Hicks, B. B., and Heilman, J. L. (1979), Techniques for calculating energy and mass fluxes, *In Modification of the Aerial Environment of Crops* (B. L. Barfield and J. F. Gerber, Eds.), Amer. Soc. of Agric. Eng., St. Joseph, MI.
- Kim, J. and Verma, S. B. (1990a), Components of surface energy balance in a temperate grassland ecosystem, *Boundary-Layer Meteorol.* 51:401-417.
- Kim, J. and Verma, S. B. (1990b), Carbon dioxide exchanges in a temperate grassland ecosystem, *Boundary-Layer Meteorol.* 52:135-149.
- Kim, J. and Verma, S. B. (1991), Modeling canopy stomatal conductance in a temperate grassland ecosystem, *Agric. and Forest Meteorol.* 55:149-166.
- Knapp, A. K. (1985), Effect of fire and drought on the ecophysiology of *Andropogon gerardii* and *Panicum virgatum* in a tallgrass prairie, *Ecology* 66:1309-1320.
- Kniezys, F. X., Shettle, E. P., Gallery, W. O., Chetwynd, J. H., Abreu, L. W., Selby, J.E.A., Clough, S. A., Fenn, R. W., (1988), *Atmospheric transmittance/radiance: computer code LOWTRAN-7*, AFGL-TR88-0177, Air Force Geophysics Lab, Hanscom AFB, Massachusetts.
- Kucera, C. L. and Kirkham, D. R. (1971), Soil respiration studies in tallgrass prairie in Missouri, *Ecology* 52:912-915.
- Lohammar, T., Larsson, S., Lindner, S., and Falk, S. (1980), FAST-Simulation models of gaseous exchange in Scots pine. *In Structure and Function of Northern Coniferous Forests - An Ecosystem Study*, *Ecological Bull.* 32:505-523.

- Markham, B. L., Wood, Jr., F. M., and Ahmad, S. P. (1988), Radiometric calibration of the reflective bands of NS001-thematic mapper simulator (TMS) and modular multispectral radiometers (MMR). Proc. of SPIE Tech. Symp., April 4-8, 1988, Orlando, Florida.
- Monteith, J. L. (1965), Evaporation and environment. *In* *The State and Movement of Water in Living Organisms* (G. E. Fogg, Ed.), Academic Press, New York, pp. 205-234.
- Monteith, J. L. (1977), Climate and the efficiency of crop production in Britain, *Phil. Trans. Roy. Soc. London* B281:277-294.
- Norman, J. M., R. L. Garcia and S. B. Verma. 1991. Soil CO<sub>2</sub> fluxes and the carbon budget of a grassland. *J. Geophys. Res.* (submitted).
- Owen, P. R., and Thomson, W. R. (1963), Heat transfer across rough surfaces, *J. Fluid Mech.* 15:321-334.
- Polley, H. W., Norman, J. M., Arkebauer, T. J., Walter-Shea, E. A., Greigor, Jr., D. H., and Bramer, C. (1991), Gas exchange of *Andropogon gerardii* Vitman, *Panicum virgatum* L., and *Sorghastrum nutans* (L.) Nash in a tallgrass prairie, *J. Geophys. Res.* (submitted).
- Schuepp, P. H., Leclerc, M. Y., MacPherson, J. I., and Desjardins, R. L. (1990), Footprint prediction of scalar fluxes from analytical solutions of the diffusion equation, *Boundary-Layer Meteorol.* 55:355-373.
- Sellers, P. J. (1985), Canopy reflectance, photosynthesis and transpiration, *Int. J. Remote Sens.* 6(8):1335-1372.
- Sellers, P. J. (1987), Canopy reflectance, photosynthesis, and transpiration. II. The role of biophysics in the linearity of their interdependence, *Remote Sens. Environ.* 21:143-183.
- Sellers, P. J., Hall, F. G., Strebel, D. E., Kelly, R. D., Verma, S. B., Markham, B. L., Blad, B. L., Schimel, D. S., Wang, J. R., and Kanemasu, E. (1988), FIFE: April 1988 Workshop Report. Code 923, NASA/Goddard Space Flight Center, Greenbelt, MD. 257 pp.
- Stewart, J. B. (1988), Modelling surface conductance of Pine forest. *Agric. For. Meteorol.* 43:19-35.
- Thom, A. S. (1972), Momentum, mass, and heat exchange of vegetation, *Quart. J. Roy. Meteorol. Soc.* 98:124-134.

- Thom, A. S. (1975), Momentum, mass, and heat exchange of plant communities. In *Vegetation and the Atmosphere, Vol. 1* (J. L. Monteith, Ed.), Academic Press, New York, pp. 57-109.
- Tucker, C. J., Holben, B. N., Elgin, J. H., and McMurtrey, E. (1981), Remote sensing of total dry matter accumulation in winter wheat, *Remote Sens. Environ.* 11:171-190.
- Verma, S. B. (1989), Aerodynamic resistances to transfers of heat, mass and momentum. In *Estimation of Areal Evapotranspiration*. International Association of Hydrological Sciences Publication No. 177, pp. 13-20.
- Verma, S. B. (1990), Micrometeorological methods for measuring surface fluxes of mass and energy, *Remote Sens. Reviews* 5:99-115.
- Verma, S. B., Kim, J., and Clement, R. J. (1991), Momentum, water vapor and carbon dioxide exchange at a centrally located prairie site in FIFE. *J. Geophys. Res.* (submitted).
- Wesely, M. L. and Hicks, B. B. (1977), Some factors that affect the deposition rates of sulfur dioxide and similar gases on vegetation, *J. Air Pollution Control Assoc.* 27:1110-1115.
- Wrigley, R. C., Spanner, M. A., Slye, R. E., Poeschel, R. F., and Livingston, J. M., (1990), "Optical depth measurements and atmospheric correction of remotely sensed data for FIFE", pp 127-134, Preprint volume of the symposium on the First ISLSCP Field Experiment (FIFE), Feb 7-9, 1990, Anaheim, Calif., AMS, Boston, Mass.

## 7. APPENDICES

### APPENDIX A

The procedure employed to derive the stress factors  $f(T)$ ,  $f(D)$  and  $f(\theta_s)$  (see Eqs. 3 and 4) is described here.

Leaf stomatal conductance ( $g_l$ ) can be described in terms of a simple multiplicative model (see *e.g.*, Jarvis, 1976; Kim and Verma, 1991):

$$g_l = g_l^*(PAR) \cdot [f(T) \cdot f(D) \cdot f(\theta_s)], \quad (A-1)$$

where  $g_l^*(PAR)$  is the unstressed value of leaf stomatal conductance. Similarly, leaf photosynthesis can be estimated as:

$$P_l = P_l^*(PAR) \cdot [f(T) \cdot f(D) \cdot f(\theta_s)], \quad (A-2)$$

where  $P_l^*(PAR)$  is the unstressed value of leaf photosynthesis. The stress factors  $f(T)$ ,  $f(D)$  and  $f(\theta_s)$  account for the effect of temperature ( $T$  in C), ambient vapor pressure deficit ( $D$  in kPa) and extractable soil water ( $\theta_s$  in %). Values of  $\theta_s$  were computed as the ratio of actual to total soil moisture held with a water potential between -1/30 and -1.5 MPa.

Equations (A-1) and (A-2) can be integrated ("scaled up") to provide estimates of canopy stomatal conductance ( $g_c$ ) and canopy photosynthesis ( $P_c$ );

$$g_c = F(\Sigma) \int_0^{L_T} g_l^*(PAR) dL = g_c^*(PAR) \cdot F(\Sigma) \quad (A-3)$$

and

$$P_c = F(\Sigma) \int_0^{L_T} P_i^*(PAR) dL = P_c^*(PAR) \cdot F(\Sigma) , \quad (A-4)$$

where  $F(\Sigma) = f(T) \cdot f(D) \cdot f(\theta_s)$  and  $L_T$  = total leaf area index. We assume that the stress factors are independent of leaf area index.

There appears to be a wide range of temperature optima for the grass species in the experimental area (Knapp, 1985) and, therefore, the effect of changes in T was assumed to be negligible (*i.e.*,  $f(T) \approx 1$ ). The response of the unstressed leaf stomatal conductance to PAR was estimated using a hyperbolic relationship (*e.g.*, Monteith, 1965):

$$g_i^*(PAR) = (a_1 PAR \downarrow) / (a_2 + PAR \downarrow) , \quad (A-5)$$

where  $PAR \downarrow$  is the normal (relative to the leaf) flux density of incident PAR, and  $a_1$  and  $a_2$  are empirically derived constants<sup>5</sup>. The leaf stomatal conductance data of the three dominant species (big bluestem, indianguass and switchgrass) measured at the site were used to evaluate parameters  $a_1$  and  $a_2$  for each of the three species (see Kim and Verma, 1991 for details). Averaged values of these parameters ( $\bar{a}_1 = 19.7 \text{ mm s}^{-1}$  and  $\bar{a}_2 = 441 \text{ } \mu\text{Ei m}^{-2} \text{ s}^{-1}$ ) were used here to yield a canopy level relationship for the mixed vegetation.

Using  $f(T) = 1$  and Eq. (A-5), Eq. (A-3) can be rewritten as:

$$g_c = \left[ \int_0^{L_T} \left( \frac{\bar{a}_1 PAR \downarrow}{\bar{a}_2 + PAR \downarrow} \right) dL \right] f(D) f(\theta_s) . \quad (A-6)$$

Substituting the Ross-Goudriaan function (see *e.g.*, Sellers, 1985, Table 3, p. 1351), Eq. (A-6) can be simplified to:

---

<sup>5</sup>The parameter,  $a_1$ , represents the asymptotic value of  $g_i^*$  when  $PAR \rightarrow \infty$ , and can be considered as the maximum stomatal conductance under optimal conditions; and  $a_2$  defines the curvature of the response to PAR.

$$g_c = \frac{\bar{a}_1}{k_x} \left[ \ln \frac{\mu \bar{a}_2 + G(\mu) PAR_o}{\mu \bar{a}_2 + G(\mu) PAR_o e^{-k_x L_T}} \right] f(D) f(\theta_s), \quad (A-7)$$

where  $k_x$  = extinction coefficient for PAR =  $(1 - \omega_x)^{1/2} G(\mu)/\mu$ ,  $\omega_x \approx 0.2$ ,  $G(\mu) \approx 0.5$  (for spherical leaf distribution) and  $PAR_o$  = PAR flux above the canopy.

A curvilinear reduction in canopy conductance with increasing vapor pressure deficit (D) was assumed (e.g., Lohammar *et al.*, 1980):

$$f(D) = 1/(1 + a_3 D), \quad (A-8)$$

where  $a_3$  is a constant. The response of canopy conductance to  $\theta_s$  (extractable soil water) was estimated using a negative exponential relationship (e.g., Stewart, 1988; Kim and Verma, 1991):

$$f(\theta_s) = 1 - \exp(-a_4 \theta_s), \quad (A-9)$$

where  $a_4$  is a constant.

As discussed in Sec. 2.5,  $g_c$  was computed from our half-hourly measurements of fluxes of net radiation, soil heat, sensible heat and latent heat. Values of  $\mu$  and  $k_x$  were calculated every half hour. The stress functions  $f(D)$  and  $f(\theta_s)$  were then evaluated by fitting (nonlinear least squares) these data ( $g_c$ ,  $\mu$ ,  $k_x$ ,  $PAR_o$ ,  $L_T$ ) to Eq. A-7:

$$\begin{aligned} a_3 &= 0.2031 \pm 0.0536 \\ a_4 &= 0.0144 \pm 0.0017 \\ (r^2 &= 0.788) \end{aligned} \quad (A-10)$$



(These values were derived from 450 half-hour  $g_c$  data points collected during 20 days distributed throughout the season.)

Thus, the following stress factors were used in this study:

$$f(D) = 1/(1 + 0.2031D) \quad (\text{A-11})$$

and

$$f(\theta_p) = 1 - \exp(-0.0144\theta_p) \quad (\text{A-12})$$

where  $D$  is the air vapor pressure deficit measured at 2.25 m above ground and  $\theta_p$  is the daily value of extractable soil water over the primary root zone (0 to 1.4 m depth).

These derived stress factors were applied to the estimates of  $g_c$  and  $P_c$  computed from flux observations, to calculate values of  $g_c^*$  and  $P_c^*$  (see Sec. 3.3). Note that while values of  $L_T$  were used to derive the coefficients given in Eqs. (A-11) and (A-12), the coefficients and the forms of stress factors are independent of the time-varying leaf area index.

## APPENDIX B

Heat (or water vapor) transfer in the immediate vicinity of a vegetated surface is primarily controlled by molecular diffusion. Momentum, however, is transferred to a large extent by pressure forces, which have no analog in heat (or water vapor) transfer. Transfer of momentum in aerodynamically rough flows (*e.g.*, over a vegetated surface) is primarily independent of molecular viscosity. Therefore, the transfer of heat (or water vapor) to or from a vegetated surface encounters greater aerodynamic resistance than does the transfer of momentum (Owen and Thomson, 1963; Chamberlain, 1966, 1968; Thom, 1972; Verma, 1989). Accordingly, an "excess resistance" term,  $r_e$  is defined:

$$r_a = r_{am} + r_b \quad (\text{B-1})$$

where  $r_{am}$  and  $r_a$  are the aerodynamic resistance to transfer of momentum and sensible heat/water vapor.

The terms  $r_{am}$  and  $r_a$  can be obtained from:

$$\tau = \rho \frac{\bar{U}}{r_{am}} \quad (\text{B-2})$$

and

$$H = \rho C_p \frac{T_m - \bar{T}}{r_a}, \quad (\text{B-3})$$

where  $\tau (= \rho u_*^2)$  is the momentum flux,  $u_*$  is the friction velocity,  $\bar{U}$  and  $\bar{T}$  are the mean horizontal wind speed and air temperature at a reference height. The term  $T_m$  represents an effective surface temperature. Equation (B-2) can be rewritten as:

$$r_{am} = \bar{U}/u_*^2. \quad (\text{B-4})$$

Owen and Thomson (1963) proposed a dimensionless parameter  $B^{-1}$  (sublayer Stanton number) to express the difference between  $r_a$  and  $r_{am}$ :

$$B^{-1} = r_a^* - r_{am}^*, \quad (\text{B-5})$$

where  $r_a^*$  and  $r_{am}^*$  are nondimensionless resistances given by:

$$r_a^* = r_a u_*, \quad (\text{B-6})$$

and

$$r_{am}^+ = r_{am} u_* \quad (B-7)$$

Several investigations (e.g., Garratt and Hicks, 1973; Brutsaert, 1984) suggest that for surfaces with densely spaced permeable roughness elements, as a first approximation  $k B^{-1}$  can be assumed to be of the order of 2 for scalars whose Schmidt ( $S_c$ ) or Prandtl ( $P_r$ ) number is of the order of 0.6 to 0.8 ( $k$  is von Karman's constant). Therefore, the excess resistance term,  $r_b$ , can be expressed as:

$$r_b = (2/ku_*) (S/P_r)^{2/3} = (2/ku_*) (\kappa/D_v)^{2/3} \quad (B-8)$$

The factor  $(S/P_r)^{2/3}$  is incorporated to account for the fact that the basic information was derived from heat transfer observations primarily (Wesely and Hicks, 1977). The term  $\kappa$  represents the thermal diffusivity and  $D_v$  represents the molecular diffusivity of water vapor.

## APPENDIX C

Table 1. Values of  $\partial g_c^*/\partial \text{PAR}$  and  $\partial P_c^*/\partial \text{PAR}$ .

| Date (1987) | $g_c^*$ vs PAR   |                | $P_c^*$ vs PAR   |                |
|-------------|--|----------------|--|----------------|
|             | $(\partial g_c^*/\partial \text{PAR}) \times 1000$<br>(mm s <sup>-1</sup> /μEI m <sup>-2</sup> s <sup>-1</sup> ) | r <sup>2</sup> | $(\partial P_c^*/\partial \text{PAR}) \times 1000$<br>(mg m <sup>-2</sup> s <sup>-1</sup> /μEI m <sup>-2</sup> s <sup>-1</sup> ) | r <sup>2</sup> |
| June 4      | 14.0   | 0.590          | 1.01   | 0.845          |
| June 5      | 9.4  | 0.976          | 1.23   | 0.951          |
| June 6      | 8.6  | 0.964          | 1.19   | 0.927          |
| July 11     | 10.8   | 0.887          | 1.24   | 0.966          |
| August 11   | 0.2  | 0.002          | 0.14   | 0.029          |
| August 15   | 8.2  | 0.961          | 1.16   | 0.961          |
| August 16*  | 8.6  | ---            | 1.18   | ---            |
| August 17*  | 9.1  | ---            | 1.19   | ---            |
| August 20   | 10.3   | 0.955          | 1.24   | 0.923          |
| October 7*  | 2.5  | ---            | 0.44   | ---            |
| October 8   | 2.5  | 0.438          | 0.44   | 0.683          |
| October 11* | 2.5  | ---            | 0.44   | ---            |
| October 13* | 2.5  | ---            | 0.44   | ---            |

\*Estimated data (see text for details).

Table 2. Information of annual life cycle stage, green leaf area index, extractable soil water ( $\theta_s$ ; 0-1.4 m depth), midday (averaged during 1230-1430 hrs, central daylight savings time) mean air temperature (T) and vapor pressure deficit (D) and photosynthetically active radiation (PAR) on selected days when reflectance and flux measurements were available. T and D were measured at 2.25 m above ground.

| Date    | Day Number | Annual Life Cycle Stage | $\theta_s$ (%) | Green LAI | T (C) | D (KPa) | PAR ( $\mu\text{Ei m}^{-2} \text{s}^{-1}$ ) |
|---------|------------|-------------------------|----------------|-----------|-------|---------|---|
| June 04 | 155        | Early Growth            | 79.3           | 1.86      | 25.4  | 1.88    | 2052  |
| June 05 | 156        | Early Growth            | 77.7           | 1.90      | 27.1  | 1.91    | 2048  |
| June 06 | 157        | Early Growth            | 76.2           | 1.94      | 27.8  | 2.11    | 2050  |
| July 11 | 192        | Peak Growth             | 68.4           | 2.79      | 31.0  | 1.85    | 1971  |
| Aug. 11 | 223        | Early Senescence        | 24.8           | 2.44      | 32.6  | 2.87    | 1844  |
| Aug. 15 | 227        | Early Senescence        | 54.4           | 2.39      | 33.7  | 2.35    | 1888  |
| Aug. 16 | 228        | Early Senescence        | 54.2           | 2.38      | 30.8  | 1.57    | 1887  |
| Aug. 17 | 229        | Early Senescence        | 54.0           | 2.37      | 31.4  | 2.71    | 1885  |
| Aug. 20 | 232        | Early Senescence        | 53.4           | 2.28      | 32.8  | 2.35    | 1860  |
| Oct. 07 | 280        | Senescence              | 30.7           | 0.30      | 15.9  | 1.71    | 1351  |
| Oct. 11 | 284        | Senescence              | 29.8           | 0.20      | 11.6  | 0.97    | 1450  |
| Oct. 13 | 286        | Senescence              | 32.5           | <0.20     | 22.7  | 2.13    | 1165  |

## APPENDIX D

### LIST OF FIGURES

- Fig. 1. Seasonal precipitation, extractable soil water (0-1.4 m depth) and green leaf area index (after Kim and Verma, 1991).
- Fig. 2. Seasonal variation in midday (1230-1430 h, central daylight savings time) values of canopy conductance, (after Kim and Verma, 1990a). The days on which SR was measured are indicated with arrows.
- Fig. 3. Seasonal variation in midday (1230-1430 h, central daylight savings time) values of canopy photosynthesis (after Verma *et al.*, 1991). The days on which SR was measured are indicated with arrows.
- Fig. 4. Seasonal variation in the simple ratio vegetation index (SR).
- Fig. 5. Unstressed canopy conductance ( $g_c^*$ ) as a function of the photosynthetically active radiation (PAR). Only afternoon data were used to minimize confounding effects. Estimates of  $\partial g_c^* / \partial \text{PAR}$  were obtained by applying simple linear fits to these data.
- Fig. 6. Unstressed canopy photosynthesis ( $P_c^*$ ) as a function of the photosynthetically active radiation (PAR). Estimates of  $\partial P_c^* / \partial \text{PAR}$  were obtained by applying simple linear fits to the data.
- Fig. 7. Simple ratio vegetative index (SR) as a function of the derivative ( $\partial g_c^* / \partial \text{PAR}$ ) of the unstressed canopy conductance with respect to the photosynthetically active radiation (PAR). The correlation coefficient ( $r^2$ ) for the linear regression is 0.56 for all data and 0.69 for the data set excluding 4 June and 11 August. The solid line represents the linear regression of the data set excluding 4 June and 11 August.
- Fig. 8. Simple ratio vegetative index (SR) as a function of the derivative ( $\partial P_c^* / \partial \text{PAR}$ ) of the unstressed canopy photosynthesis with respect to the photosynthetically active radiation (PAR). The correlation coefficient ( $r^2$ ) for the linear regression is 0.41 for all data and 0.57 for the data set excluding 11 August.




Figure 1




Figure 2






Figure 3




Figure 4




Figure 5




Figure 6




Figure 7




Figure 8