

N89-26347

75-51  
1125 11-1  
198680  
18.

MICROBIAL TRACE FOSSILS IN ANTARCTICA  
AND THE SEARCH FOR EVIDENCE OF EARLY LIFE ON MARS

E. Imre Friedmann<sup>1</sup> and Roseli O. Friedmann<sup>2</sup>

<sup>1</sup>Polar Desert Research Center and

Department of Biological Science

Florida State University FT 988847  
Tallahassee, FL 32306-2043

<sup>2</sup>Department of Biology

Florida A&M University FT 402173  
Tallahassee, FL 32307

It is possible to hypothesize that, if microbial life evolved on early Mars, fossil remnants of these organisms may be preserved on the surface. However, the cooling and drying Mars probably resembled a cold desert and such an environment is not suitable for the process of fossilization.

The frigid Ross Desert of Antarctica is probably the closest terrestrial analog to conditions that may have prevailed on the surface of the cooling and drying Mars. In this desert, cryptoendolithic microbial communities live in the airspaces of porous rocks, the last habitable niche in a hostile outside environment. The organisms produce characteristic chemical and physical changes in the rock substrate. Environmental changes (deterioration of conditions) may result in death of the community. Although no cellular structures are fossilized, the conspicuous changes in the rock substrate are preserved as trace fossils. Likewise, microbial trace fossils (without cellular structures) may also have been preserved on Mars: Discontinuities in structure or chemistry of the rock that are independent of physical or chemical gradients may be of biological origin. Ross Desert trace fossils can be used as a model for planning search strategies and for instrument design to find evidence of past Martian life.