

NEUTRON CAPTURE PRODUCTION RATES OF COSMOGENIC ^{60}Co , ^{59}Ni AND ^{36}Cl IN STONY METEORITES; M. S. Spergel, York College of CUNY, Jamaica, New York, 11451, USA; R. C. Reedy, Los Alamos National Laboratory, Los Alamos, New Mexico, 87545, USA; O. W. Lazareth and P. W. Levy, Brookhaven National Laboratory, Upton, New York 11973 USA

To unfold the cosmic-ray exposure history of a meteorite, it is best to use a variety of cosmogenic products (tracks and nuclides) with different production profiles. Neutron-capture reactions have production rates which vary considerably with sample depth and meteorite size¹. Their production profiles differ appreciably from those for tracks or nuclides created by energetic cosmic ray particle spallation reactions. In large meteorites neutron-capture reactions are the main sources of cosmic-ray produced nuclides such as ^{59}Ni and ^{60}Co .¹ The cosmogenic radionuclide ^{60}Co , from the $^{59}\text{Co}(n,\gamma)^{60}\text{Co}$ reaction, has been a very useful tool for unfolding the cosmic-ray exposure record of the large Jilin (Kirin) chondrite^{2,3}.

The neutron-capture reaction rates producing ^{36}Cl , ^{59}Ni , and ^{60}Co in meteorites were calculated by Eberhardt, et al.¹ using neutron slowing-down theory. Lingenfelter et al.⁴ used neutron-transport theory to calculate the low-energy neutron flux and neutron-capture induced isotopic anomalies in the moon. Previously we reported neutron-transport theory calculation of the low-energy neutron, as a function of depth, in spherical meteoroids⁵ and preliminary results for ^{59}Ni and ^{60}Co production rates.⁶⁻⁷ Reported here are complete results for neutron flux calculations in stony meteoroids, of various radii and compositions, and production rates for ^{36}Cl , ^{59}Ni , and ^{60}Co .

New neutron source strengths have been calculated that increase our calculated production rates by about 30% in larger meteorites.¹¹ The $^{59}\text{Ni}/^{60}\text{Co}$ production ratio in spherical L-chondrites with radii $>150\text{ g/cm}^2$ is usually within agreement with measurements on various large meteorites; but higher than the ratio as calculated by Eberhardt, et al.¹ Neutron-capture calculations for a C3-chondrite with 100-ppm hydrogen and for an aubrite ($\approx 1\%$ Fe) provide neutron-capture systematics that differ considerably from those obtained with L-chondrites. Measured neutron-capture radionuclides in the Allende meteorite agree better with calculation for a dry chondrite than for one with 100-ppm H; indicating that Allende had a low H content. Our lunar calculations agree with the calculation of Lingenfelter et al.⁴, Kornblum et al.⁹, and with the lunar neutron measurements.⁸ Both the absolute values and the activity-versus-depth profiles calculated for ^{60}Co formation in the Moon agree with the measurements of Wahlen et al.¹⁰ For large spheres the calculated results converge to those obtained for the Moon, but there are significant differences between the lunar results and those predicted for a meteorite with a radius of 1000 g/cm^2 . The calculated neutron fluxes and nuclide production rates for small spheres are quite different from those for large meteorites.

The $^{59}\text{Ni}/^{60}\text{Co}$ ratio is nearly constant with depth in most meteorites: this effect is consistent with the neutron flux and capture cross

NEUTRON CAPTURE PRODUCTION RATES

Spergel, M. S., Reedy, R. C., Lazareth, O. W. and Levy, P. W.

section properties. The shape of the neutron flux energy spectrum, varies little with depth in a meteorite. The size of the parent meteorite can be determined from one of its fragments, using the $^{59}\text{Ni}/^{60}\text{Co}$ ratios, if the parent meteorite was less than 75 g/cm^2 in radius. If the parent meteorite was larger, a lower limit on the size of the parent meteorite can be determined from a fragment. In C3 chondrites this is not possible.

In stony meteorites with $R < 50\text{ g/cm}^2$ the calculated ^{60}Co production rates (mass $< 4\text{ kg}$), are below 1 atom/min g-Co. The highest ^{60}Co production rates occur in stony meteorites with radius about 250 g/cm^2 (1.4 m across). In meteorites with radii greater than 400 g/cm^2 the maximum ^{60}Co production rate occurs at a depth of about 175 g/cm^2 in L-chondrite, 125 g/cm^2 in C3 chondrite, and 190 g/cm^2 in aubrites. Production results for ^{60}Co and ^{59}Ni in meteorites of radius 300 g/cm^2 ($\approx 86\text{ cm}$) are shown in Fig. 1 and Fig. 2 respectively. The figures contain results for L Chondrite and C3 type meteorites.




FIG. 1. ^{60}Co production (DPM/Kg) curves in a L Chondrite (A) and a C3 (B). Peak activity occurs closer to the surface in the C3 Chondrite. Cobalt activity levels are effected by the different hydrogen abundances in the meteorites.

NEUTRON CAPTURE PRODUCTION RATES

Spergel, M. S., Reedy, R. C., Lazareth, O. W. and Levy, P. W.




FIG. 2. ^{59}Ni production (DPM/Kg) curves in L Chondrite, C3 Chondrite and Aubrite type meteorites. Peak activity occurs closest to the surface with the C3 Chondrite.

1. Eberhardt, P., Geiss, T., and Lutz, M. (1963) Neutrons in meteorites, Earth Sci. and Meteoritics, North Holland, Amsterdam, p. 143-168.
2. Honda, M., Nishiizumi, K., Imamura, T., Takaoka, N., Nitoh, O., Horie, K., and Komura, K. (1982) Earth Planet. Sci. Lett. 57, p. 101-109.
3. Heusser, G. and Ouyang, Z. (1981) Meteoritics 16, p. 326-327.
4. Lingenfelter, R. E., Canfield, E. H., and Hempel, V. E. (1972) Earth Planet. Sci. Lett. 16, p. 355-369.
5. Spergel, M. S., Reedy, R. C., Lazareth, O. W., and Levy, P. W. (1980) Meteoritics, 15, p. 370.
6. Spergel, M. S., Reedy, R. C., Lazareth, O. W., and Levy, P. W. (1981) Meteoritics 16, p. 387-388.
7. Spergel, M. S., Reedy, R. C., Lazareth, O. W., and Levy, P. W. (1982) Lunar and Planetary Science XIII, p. 756-757.
8. Woolum, D. S., Burnett, D. S., Furst, M., and Weiss, J. R. (1975) The Moon 12, p. 231-250.
9. Kornblum, J. J., Fireman, E. L., Levin, M., and Aronson, A. (1973) Proc. Lunar Sci. Conf. 4th, p. 2171-2182.
10. Wahlen, M., Finkel, R. C., Imamura, M., Kohl, C. P., and Arnold, J. R. (1973) Earth Planet. Sci. Lett. 19, p. 315-320.