

1 page

NASA TECHNICAL MEMORANDUM

TM-88423

IN-18389

PRELIMINARY RESULTS OF DETERMINATION OF CHEMICAL ELEMENT
CONCENTRATIONS IN THE AEROSOL OF VENUS CLOUDS

B.M. Andreychikov, L.M. Mukhin, B.N. Korchuganov, I.K.
Akhetshin, Ye.N. Tokarev, A.V. Medvedev, M.N. Gol'dfel'd,
V.M. Faynboym, A.V. Kalyuzhnyy, I.V. Petryanov, B.I. Ogorodnikov
and V.I. Skitovich

Translation of "Predvaritel'nyye rezul'taty opredeleniye soder-
zhaniya khimicheskikh elementov v aerozole oblakov venery",
Pis'ma v AZh, vol. 12, No. 2, 1986, pp. 120-122.

(NASA-TM-88423)	PRELIMINARY RESULTS OF	N86-30617
	DETERMINATION OF CHEMICAL ELEMENT	
	CONCENTRATIONS IN THE AEROSOL OF VENUS	
	CLOUDS (National Aeronautics and Space	Unclas
Administration)	7 p	CSSL 03B G3/91 43510

ORIGINAL PAGE IS
OF POOR QUALITY

STANDARD TITLE PAGE

1. Report No. TM-88423		2. Government Accession No.		3. Recipient's Catalog No.	
4. Title and Subtitle PRELIMINARY RESULTS OF DETERMINATION OF CHEMICAL ELEMENT CONCENTRATIONS IN THE AEROSOL OF VENUS CLOUDS				5. Report Date May 1986	
				6. Performing Organization Code	
7. Author(s) B.M. Andreychikov, L.M. Mukhin, B.N. Korchuganov, I.K. Akhmetshin, Ye.N. Tokarev, A.V. Medvedev, M.N. Gol'dfel'd, V.M. Faynboym, I.V. Kalyuzhnyy, et al.				8. Performing Organization Report No.	
				10. Work Unit No.	
9. Performing Organization Name and Address Leo Kanner Associates, Redwood City, California 94063				11. Contract or Grant No. NASw-4005	
				12. Type of Report and Period Covered Translation	
12. Sponsoring Agency Name and Address National Aeronautics and Space Administration, Washington, D.C. 20546				14. Sponsoring Agency Code	
15. Supplementary Notes Translation of "Predvaritel'nyye rezul'taty opredeleniy sodержaniya khimicheskikh elementov v aerozole oblakov venery", Pis'ma v AZh, vol. 12, No. 2, 1986, pp. 120-122. (UDC 523.4)					
16. Abstract An x-ray radiometric experiment is described along with the results of measurements of the elemental composition of aerosol in Venusian clouds. A preliminary analysis of the data showed that sulfur is present in the range of heights 63-47 km with mean content of 5.8 mg/m ³ and that chlorine is present in the height range 61-52 km with a mean content of 4.1 mg/m ³ . The results of measurements in the range 52-47 km may come to an agreement if phosphorus is present in the aerosol with a mean concentration of 7.7 mg/m ³ .					
17. Key Words (Selected by Author(s))				18. Distribution Statement Unclassified-Unlimited	
19. Security Classif. (of this report) Unclassified		20. Security Classif. (of this page) Unclassified		21. No. of Pages 6	22.

PRELIMINARY RESULTS OF DETERMINATION OF CHEMICAL ELEMENTS
CONCENTRATIONS IN THE AEROSOL OF VENUS CLOUDS

B.M. Andreychikov, L.M. Mukhin, B.N. Korchuganov, I.K.
Akhmetshin, Ye.N. Tokarev, A.V. Medvedev, M.N. Gol'dfel'd
V.M. Faynboym, A.V. Kalyuzhnyy, I.V. Petryanov, B.I.
Ogorodnikov and V.I. Skitovich

Institute of Space Investigations, USSR Academy of Sciences,
Moscow

In order to determine the concentrations of chemical elements /120*
in the aerosol of Venusian clouds on the "Vega-1" and "Vega-2" AMS
landing vehicles, x-ray radiometric instruments were installed.

The x-ray radiometric method of analysis makes it possible to
investigate a sample without destroying it and without changing the
conditions of its existence; this is especially important in analyz-
ing a material of unknown composition. The essence of the method
may be reduced to the following: during the interaction of electro-
magnetic or corpuscular radiation with the electron shells of
atomic nuclei, electrons escape from one of the shells closest to the
nucleus. An electron from a shell farther from the nucleus goes to
the empty site and this is accompanied by the emergence of an x-ray
photon with a frequency equal to the ratio of the difference in the
energies of these shells to Planck's constant (characteristic x-ray
radiation).

The characteristic x-ray radiation is transformed in a detector /121
into a current impulse and then with the help of an electronic cir-
cuit into a voltage impulse which is proportional to the energy of
the x-ray photon. The number of impulses of a given amplitude

* Numbers in the margin indicate pagination in the foreign text.

characterizes the concentration of the corresponding chemical element. In order to obtain a compact geometry of measurements, low energy capacity, mass and dimensions of the apparatus in the experiment being described, ionization of the atoms is done by isotope x-ray sources. The measurement was done continuously during the landing of the landing vehicle in a parachute in the Venusian cloud layer at altitudes ranging from 63-47 km. The aerosol was deposited on an ashless acidproof Petryanov filter which was specially created for the experiment and simultaneously analyzed. In order to ensure the maximum sensitivity of the measurements (maximum mass of the deposited aerosol), the filter had low (~ 0.1 mm water) standard resistance; this made it possible during the parachute landing of the vehicle in the Venusian clouds to pass ~ 2 m³ atmospheric gas through it. The filter's effectiveness in retaining particles more than 1 μ m in diameter was 100%.

The device combined four spectrometric channels which continuously analyzed the material being deposited on the filter. The intensity of the characteristic radiation in various sections of the spectral distribution was recorded. As the detectors we used sealed off proportional counters. With the help of two of them, measurements were done in the spectral region corresponding to the characteristic radiation of chlorine, sulfur and phosphorus. In order to separate the radiation of these elements in the aperture of one of the counters, a selective sulfur filter was used. The third counter measured the intensity of radiation in the region of the characteristic energy of iron, and the fourth, the intensity of primary radiation passing through the filter with the aerosol deposited on it. In order to excite characteristic radiation in the region of chlorine, sulfur and phosphorus, radio isotope sources based on radioactive nuclide ⁵⁵Fe were used, in the region of iron, ¹⁰⁹Cd, and for radioscoping of the filter, ³H/Zr. The construction of the device made it possible to reduce the hum caused by sources extraneous to the given experiment to the minimum (for example, a ¹⁰⁹Cd and ³H/Zr hum for determining the characteristic radiation of the element group Cl, P, S).

The device operated in the following way. After it was switched on on the command of the landing vehicle, the program timer of the device gave the command to depressurize the analytical cell in which the filter, sources and detectors of ionizing radiation were placed. The input and output valves of the analytical cell were discharged and under the effect of a high-speed thrust, the atmospheric gas began to pass through a filter of area 10 cm². During landing the presence and variations in x-ray radiation in the region of the characteristic energies of Cl, S, P were recorded along with variations in intensity in the region of characteristic radiation L - the zirconium series (a channel with a ³H/Zr source). Variations in intensity in the channel which measured the characteristic radiation of iron were above the statistical error of the measurements at altitudes of 54-47 km. /122

A preliminary analysis of the concentration of chemical elements in the aerosol in Venusian clouds showed that sulfur is present in the range of altitudes from 63-47 km and that its average concentration is 5.8 mg/m³ and that chlorine is present in the range of altitudes from 61-52 km in the amount 4.1 mg/m³. Chlorine and sulfur were previously recorded in the cloud aerosol of Venus with the help of x-ray radiometric spectrometers (Surkov et al., 1982).

The results of measurements in the range of altitudes from 52-47 km may correspond if phosphorus is present in the aerosol in average concentration 7.7 mg/m³.

Significant variations were observed in the concentrations of sulfur, chlorine and phosphorus in the aerosol with altitude; some of these may be related to phase transitions and (or) differentiations of these chemical elements in the cloud aerosol by altitude.

The preliminary data presented here will be refined and supplemented.

In conclusion, the authors would like to express their thanks to V.G. Shumkov, V.G. Rogozovets, V.F. Fateyev, B.V. Zatolokin, and B.K. Morozov for their help in the preparation of individual elements of the x-ray radiometric meter, to G.N. Krasovskiy for his constant support for and attention to this work and to A.V. D'yachkov for his help with information processing.

REFERENCES

Surkov, Yu.A., Kirnozov, F.F., Gur'yanov, V.I., Glazov, V.N., Dun-
chenko, A.G., Kurochkin, S.S., Rasputnyy, V.N., Kharitonova, E.G.,
Tatsiy, L.N. and Gimadov, V.D., Kosmich. issled.20, 435 (1982).