

N86-28558 5P 107

9204

## FLUID-ABSENT METAMORPHISM IN THE ADIRONDACKS.

John W. Valley, Department of Geology and Geophysics, The University of Wisconsin, Madison, WI 53706, USA W4560409

Introduction. Granulite facies metamorphism occurred over an area of  $> 20,000 \text{ km}^2$  in the Adirondack Highlands, N.Y., USA and has been dated at  $\sim 1.1$  by. (Grenville Orogeny, ref. 1). Peak metamorphic conditions as determined from numerous geothermometers and barometers in many rock types increase from about  $675^\circ$  at the orthopyroxene isograd in the NW to  $775^\circ\text{C}$  in the Central Adirondacks at  $7.6 \pm 0.5$  kbar (Fig. 1, ref. 2). Low values of water activity ( $\alpha\text{H}_2\text{O}$ ) are demonstrated by assemblages of tremolite + diopside + enstatite + quartz; phlogopite + quartz + enstatite + K-feldspar; annite + K-feldspar + magnetite + ilmenite; and unmelted K-feldspar + plagioclase + quartz. Quantitative estimates from mica and amphibole equilibria yield  $\alpha\text{H}_2\text{O} \approx 0.1$  (3,4,5). Muscovite + quartz assemblages in adjacent amphibolite facies rocks in the NW Adirondack Lowlands indicate high  $\alpha\text{H}_2\text{O}$  demonstrating that the amphibolite to granulite facies transition in the Adirondacks is the result of increasing temperature and decreasing  $\alpha\text{H}_2\text{O}$ .

Causes of low  $\alpha \text{H}_2\text{O}$ . Three processes can account for low  $\alpha \text{H}_2\text{O}$  in granulites: A) large amounts of  $\text{CO}_2$  infiltration where  $\text{H}_2\text{O}$ -rich fluids are diluted by  $\text{CO}_2$  and dehydration reactions for mica and amphibole are driven in the prograde direction (6); B) partial melting with or without the extraction of a magma and where  $\text{H}_2\text{O}$  is partitioned preferentially into the silicate liquid (7); and C) metamorphism of igneous or metamorphic rocks that were already "dry."

The recognition that one of these processes has influenced the formation of a given rock is important in understanding its history because the results of each process can be quite different. Only processes A) and B) can result in large amounts of mass transport such as is necessary to dehydrate a metasediment or cause LIL element depletions. Processes B and C on the other hand can lead to a metamorphic recrystallization in the absence of a free fluid phase where any hydrogen or carbon that may be in a rock during metamorphism is not in the form of an  $\text{H}_2\text{O} - \text{CO}_2$  fluid.  $\text{CO}_2$  infiltration, process A, cannot proceed under fluid-absent conditions.

Fluid-absent metamorphism. Fluid-absence has thus far been demonstrated in two rock types in the Adirondacks. In calc-silicates and marbles adjacent to massif-type anorthosite bodies low values of  $\delta^{18}\text{O}$  (-1.3 to 3) indicate that low P contact metamorphism preceded granulite metamorphism (8,9). In these rocks numerous wollastonite occurrences indicate low  $\alpha\text{CO}_2$  and in one rock the assemblage monticellite + forsterite + diopside + calcite + spinel indicates  $P \text{H}_2\text{O} + P \text{CO}_2 < 0.5$  kbar at  $P = 7.5$  kbar (Fig. 2). These results, plus the close preservation of pre-regional metamorphic  $\delta^{18}\text{O}$  values, indicate that granulite facies metamorphism was representative of process C where early shallow contact metamorphism volatilized marbles and subsequent granulite metamorphism was fluid-absent.

FLUID-ABSENT METAMORPHISM  
J.W. Valley

Many charnockites in the Adirondacks are also indicated to have been fluid-absent by low, buffered values of oxygen fugacity ( $fO_2$ ) and water fugacity ( $fH_2O$ , ie  $aH_2O$ ). Estimates of  $fO_2$  in Adirondack charnockites from coexisting magnetite + ilmenite (10) are mostly below the quartz + magnetite + fayalite buffer (Fig. 3) and are too low for the stable existence of a  $CO_2$ -rich metamorphic fluid (Fig. 4). These rocks typically contain orthopyroxene + K-feldspar assemblages which also indicate low  $fH_2O$ . Simultaneous solution of C-O-H fluid equations show that at such low  $fO_2$  and low  $fH_2O$  there cannot have been a free C-O-H fluid phase and thus metamorphism was fluid-absent in these rocks (11). Many of these charnockites are clearly indicated by field relations, textures, and geochemistry to be meta-igneous rocks and thus it is reasonable that they experienced fluid-absent metamorphism because they were already "dry." However, this has not been demonstrated in all instances and significant partial melting is likely to also be important.

Recognition of fluid-absence in some Adirondack rocks does not necessarily imply that this entire former portion of the deep crust was lacking in fluids. Assemblages of phlogopite + calcite + quartz and tremolite + calcite + quartz are common and at Adirondack P-T this requires  $P H_2O + P CO_2 \approx 6-8$  kbar (12) suggesting fluid saturation.

This result in no way conflicts with fluid-absence for other rocks if the fluid regime of the deep crust is complex. Sharp gradients in  $\delta^{18}O$  and in buffered fluid compositions (3,8) indicate that fluid conditions in the Adirondacks were locally heterogeneous. Thus it is likely that processes B, C, and possibly A all operated synchronously and possibly in close proximity. However, no evidence of large scale  $CO_2$  infiltration has yet been found in the Adirondacks and even in terranes where strong evidence of  $CO_2$  infiltration has been located in some rocks (6) it is premature to assume that all rocks were affected.

ORIGINAL PAGE IS  
OF POOR QUALITY




Figure 1. Values of peak metamorphic pressure and temperature in the Adirondacks (ref. 2).




Figure 2. Mineral equilibria buffering  $f_{CO_2}$  and  $f_{H_2O}$  at 780°C and 8 kbar. The dashed curve represents  $P_{H_2O} + P_{CO_2} = 8$  kbar and fluid-absent conditions plot below or to the left. The fluid-absent assemblage monticellite + forsterite + diopside + calcite + spinel plots at  $\log f_{CO_2} = 3.3$ ,  $\log f_{H_2O} < 2.5$  (ref. 9).




Figure 3. Log  $f_{O_2}$  - T plot at 8 kbar. Data points are from coexisting magnetite + ilmenite (10,13). Abbreviations are: H hematite, Q quartz, M magnetite, F fayalite, U ulvospinel, I ilmenite, W wustite, C graphite, Tr tremolite, Cc calcite, Di diopside.




Figure 4. C-O-H fluid equilibria at 700°C, 7 kbar. The outermost curved solid line represents graphite saturation at  $P_{fluid} = 7$  kbar. Solid isopleths below this curve are the maximum total pressure of all C-O-H fluids possible regardless of the presence or absence of graphite and thus the field below graphite saturation represents fluid-absence unless large amounts of some non C-O-H fluid are present (ref. 11).

FLUID-ABSENT METAMORPHISM  
J.W. Valley

References

1. Ashwal, L.D. and Wooden, J.L. (1983) Sr and Nd isotope geochronology, geologic history and origin of the Adirondack anorthosite. Geochim. Cosmochim. Acta 47, 1875-1886.
2. Bohlen, S.R., Valley, J.W. and Essene, E.J. (1985) Metamorphism in the Adirondacks. I. Petrology, Pressure and Temperature. J. Petrol., v. 26, pt. 4, in press.
3. Valley, J.W., McLelland, J., Essene, E.J. and Lamb, W. (1983) Metamorphic fluids in the deep crust: Evidence from the Adirondacks. Nature 301, 226-228.
4. Bohlen, S.R., Peacor, D.R. and Essene, E.J. (1980) Crystal chemistry of a metamorphic biotite and its significance for water barometry. Am. Mineral. 65, 55-62.
5. Powers, R.E. and Bohlen, S.R. (1985) The role of synmetamorphic igneous rocks in the metamorphism and partial melting of metasediments, Northwest Adirondacks. Contr. Min. Pet. 90, 401-409.
6. Newton, R.C., Smith, J.V. and Windley, B.F. (1980) Carbonic metamorphism, granulites and crustal growth. Nature 228, 45-50.
7. Fyfe, W.S. (1973) The granulite facies, partial melting, and the Archaean Crust. Phil. Trans. Roy. Soc. Land. 273A, 457-461.
8. Valley, J.W. and O'Neil, J.R. (1984) Fluid heterogeneity during granulite facies metamorphism in the Adirondacks: stable isotope evidence. Contr. Min. Pet. 85, 158-173.
9. Valley, J.W. (1985) Polymetamorphism in the Adirondacks: wollastonite at contacts of shallowly intruded anorthosite. The Deep Proterozoic Crust in the North Atlantic Provinces (Tobi and Touret, eds.), Reidel, in press.
10. Bohlen, S.R., Essene, E.J. and Hoffman, K.S. (1980) Feldspar and oxide thermometry in the Adirondacks: an update. Bull. geol. Soc. Amer. 91, 110-113.
11. Lamb, W. and Valley, J.W. (1984) Metamorphism of reduced granulites in a low-CO<sub>2</sub>, vapor-free environment. Nature 312, 56-58.
12. Valley, J.W. and Essene, E.J. (1980) Calc-silicate reactions in Adirondack marbles: the role of fluids and solid-solutions. Bull. geol. Soc. Amer. 91, 114-117, 720-815.
13. Anderson, D.J. and Lindsley, D.H. (1985) New (and Final!) models for Ti-magnetite/ilmenite geothermometer and oxygen-barometer. EOS-Trans. Amer. Geophys. Un. 66, 416.