

NOU-20000

9182

58

17

AS INDICATED BY SR, ND, AND PB ISOTOPIC SYSTEMATICS OF PROTEROZOIC INTRUSIVES.

¹L.D. Ashwal, ²J.L. Wooden, and ³R.F. Emslie

¹Lunar and Planetary Institute, 3303 NASA Road One, Houston, TX 77058, L8618929

²U.S. Geological Survey, Isotope Geology Branch, 345 Middlefield Road, CU717886

Menlo Park, CA 94025, ³Geological Survey of Canada, 601 Booth St.,

Ottawa, Ont., Canada K1A 0E8 CU 910235

Introduction. Sm-Nd and Rb-Sr isotopic compositions of mid-to late Proterozoic (~ 1.6 - 1.1 Ga) massif-type anorthosites and mafic intrusives in the eastern Canadian shield are correlated with geographic location (1,2). Complexes in the Grenville province have positive ϵ_{Nd} values and initial $^{87}Sr/^{86}Sr$ (I_{Sr}) generally less than 0.703, suggesting derivation from depleted mantle. In Labrador, similar complexes close to or northwest of a line roughly corresponding to the Grenville Front have negative ϵ_{Nd} values and $I_{Sr} > 0.703$. This contrast has been interpreted as reflecting either Sr enriched mantle under the Nain Province (1), or contamination of the Nain intrusives with older crustal components (2). Pb isotopic compositions, however, favor the latter (3). We speculate here on the possibility of using these Proterozoic intrusives as tracers to characterize the nature and extent of older basement types in Labrador.

Harp Lake and Mealy Mountains Complexes. Our data base includes Sr, Nd, and Pb isotopic compositions of 1.6-1.65 Ga anorthosites and related rocks from the Harp Lake (Nain Province) and Mealy Mountains (Grenville Province) complexes, and also of later (~ 1.4 Ga) mafic dikes which crosscut them (Fig. 1) (3-6). The anorthosite complexes are large (7000-8000 km²) composite massifs dominated by leucotroctolites and leuconorites, with minor ferrodiorites and gabbros (4). Both complexes are crosscut by an ENE-trending swarm of olivine diabase dikes with similar mineralogy and major and minor element chemistry (5,6). Mealy dikes yield a whole-rock Rb-Sr age of 1380 ± 54 Ma (Fig. 2) (6). Harp and Mealy dikes have been suggested as correlative with other gabbroic rocks of similar age in Labrador, including the Shabogamo, Michael, and Flowers River gabbros, the Seal Lake volcanics, and the Kiglapait layered intrusion (5,6).

Mealy anorthosites have $I_{Sr} = 0.7026 - 0.7032$ and $\epsilon_{Nd} = +1.2$ to $+3.0$ (computed for 1.65 Ga). Mealy dikes have $I_{Sr} = 0.7025 - 0.7028$ and $\epsilon_{Nd} = +3.8$ to $+5.6$ (computed for 1.4 Ga). Both are consistent with derivation from mantle sources depleted with respect to Rb/Sr and Nd/Sm. Pb isotope data for Mealy anorthosites and dikes form a linear array between the evolution curves for model mantle (7) and average crust (8). This array is essentially equivalent to that for Proterozoic carbonatites, which Tilton (9) interpreted as reflecting depleted mantle.

Harp Lake samples show a much larger variation in isotopic composition (Figs. 2, 3). The major rock units of the Harp Lake anorthosite complex have I_{Sr} between 0.7039 and 0.7066, and ϵ_{Nd} between -2.4 and -6.0 (at 1.65 Ga). Harp dikes have $I_{Sr} = 0.7032 - 0.7033$ and $\epsilon_{Nd} = -0.3$ to -2.3 (at 1.4 Ga). The Nd isotopic data for Harp dikes show a narrow range of $^{147}Sm/^{144}Nd$ (0.146 - 0.153), but plot along a ~ 3.3 Ga reference line, probably reflecting mixing with an Archean component (Fig. 3). Harp Lake anorthosites and related rocks show a wide scatter in Pb isotope compositions, and can be interpreted as reflecting mixing between a depleted Proterozoic mantle component and an older component with extremely unradiogenic Pb.

Likewise, Harp dikes and plagioclase separates therefrom have very low $^{207}\text{Pb}/^{204}\text{Pb}$ for Proterozoic mantle-derived materials, and must be interpreted as reflecting contamination with U-depleted crustal material.

Possible Contaminants. Although we cannot constrain the precise nature of the contaminant, its composition must be such that it has essentially no effect on the major or minor element chemistry of Harp Lake samples, assuming that Mealy Mountains samples can be considered uncontaminated starting materials. Harp dikes show no noticeable enrichments in "crustal" components such as SiO_2 , K_2O , Na_2O , etc. The most appropriate contaminant would have sufficiently low Rb/Sr, Sm/Nd, and U/Pb such that its isotopic compositions evolve to relatively low values of $^{87}\text{Sr}/^{86}\text{Sr}$, $^{143}\text{Nd}/^{144}\text{Nd}$, and $^{207}\text{Pb}/^{204}\text{Pb}$ by about 1.4 Ga.

The Harp Lake complex, and other Proterozoic intrusives in the Nain Province of Labrador were emplaced near the boundary between Aphebian (1.7 - 1.8 Ga) gneisses of the Churchill Province, and a belt of Archean gneisses with ages as old as 3.6 Ga, exposed along the eastern coast of Labrador (Fig. 1) (4,10). The Aphebian gneisses are considered to be reworked equivalents of late Archean (~ 2.7 Ga) materials similar to those in the Superior Province (11), and the older Archean gneisses of coastal Labrador have been correlated with the ancient gneiss terrane of West Greenland (e.g. 10).

Initial Sr and Nd isotopic compositions (at 1.4 Ga) of these gneisses are compared with those of Harp and Mealy dikes in Fig. 4. Data for the Churchill and Superior Provinces are McCulloch and Wasserburg's (11) composites for large areas of these terranes. The wide range in isotopic composition among Uivak gneisses is a likely consequence of early granulite facies metamorphism which produced low Rb/Sr and I_{Sr} in some areas (12,13). Among these possibilities, the most suitable contaminant for the Harp dikes would be Rb-depleted Uivak gneiss (component B, Fig. 4). Less than 10% of such a component added to Mealy dikes would produce the isotopic composition of the Harp dikes, assuming simple two-component mixing (Fig. 4). Although the Superior and Churchill Province composites would not appear to be appropriate contaminants, we cannot rule out as a possibility a Rb-depleted equivalent of such material, produced for example, by granulite facies metamorphism (component C, Fig. 4). This hypothetical component would be a less satisfactory contaminant in that it would require a larger degree of contamination (15 - 25%). Pb isotope systematics also favor an ancient gneiss-type contaminant (Uivak or Amitsoq gneiss) over late-Archean gneisses (Fig. 5).

Discussion. Although it remains speculative, it could be inferred from our data that the ancient gneiss complex exposed along coastal Labrador extends in the subsurface, beneath the Harp Lake anorthosite complex, and possibly as far west as the Labrador Trough, where the 1.45 Ga Shabogamo gabbros also have negative ϵ_{Nd} values of about -5 (14). Pb isotopic compositions of these, and other mafic intrusives in Labrador are clearly needed to evaluate this. It is possible, therefore, that the Archean North Atlantic craton is much more extensive than indicated by present surface exposures.

In any case, Proterozoic intrusives in the Grenville Province show no evidence of contamination with Archean crustal components. The Labrador segment of the Grenville Front, therefore, appears to mark the southern edge of the Archean craton in eastern North America. It is possible that this feature coincides with a suture, but if this is the case, it must be older than 1.6 Ga.

Fig. 1




Fig. 2




Fig. 3




Fig. 4





Fig. 5



References

1. Ashwal, L.D. and Wooden, J.L. (1983) Isotopic evidence from the eastern Canadian shield for geochemical discontinuity in the Proterozoic mantle. Nature **306**, 679-680.
2. Ashwal, L.D. and Wooden, J.L. (in press) Sm-Nd isotopic studies of Proterozoic anorthosites: Systematics and implications. In, The Deep Proterozoic Crust in the North Atlantic Provinces, (J. Touret, Ed.), Proc. NATO Adv. Study Inst., Reidel, Dordrecht.
3. Ashwal, L.D., Wooden, J.L., and Emslie, R.F. (1985) Nd, Sr, and Pb isotope systematics of mid-Proterozoic dikes from the Grenville and Nain Provinces of Labrador (abstr.) EOS **66**, 414.
4. Emslie, R.F. (1980) Geology and petrology of the Harp Lake complex, central Labrador: An example of Elsonian magmatism. Geol. Surv. Canada Bull. **293**, 136 pp.
5. Myers, R.E. and Emslie, R.F. (1977) The Harp dikes and their relationship to the Helikian geological record in central Labrador. Can. J. Earth Sci. **14**, 2683-2696.
6. Emslie, R.F., Loveridge, W.D., and Stevens, R.D. (1984) The Mealy dikes, Labrador: petrology, age, and tectonic significance. Can. J. Earth Sci. **21**, 437-446.
7. Zartman, R.E. and Doe, B.R. (1981) Plumbotectonics: the model. Tectonophysics **75**, 135-162.
8. Stacy, J.S. and Kramers, J.D. (1975) Approximation of terrestrial lead isotope evolution by a two-stage model. Earth Planet. Sci. Lett. **26**, 207-221.
9. Tilton, G.R. (1983) Archean crust-mantle geochemical differentiation. In, Workshop on A Cross Section of Archean Crust (L.D. Ashwal and K.D. Card, Eds.), pp. 92-94, L.P.I. Tech. Rept. 83-03, Lunar and Planetary Institute, Houston.
10. Collerson, K.D. and Bridgwater, D. (1979) Metamorphic development of early Archean tonalitic and trondhjemitic gneisses: Saglek area, Labrador. In, Trondhjemitic, Dacitic, and Related Rocks (F. Barker, Ed.) pp. 205-273, Elsevier, Amsterdam.
11. McCulloch, M.T. and Wasserburg, G.J. (1978) Sm-Nd and Rb-Sr chronology of continental crust formation. Science **200**, 1003-1011.
12. Collerson, K.D. (1983) The Archean gneiss complex of northern Labrador. 2. Mineral ages, secondary isochrons, and diffusion of strontium during polymetamorphism of the Uivak gneisses. Can. J. Earth Sci. **20**, 707-718.
13. Collerson, K.D., Kerr, A., Vocke, R.D., and Hanson, G.N. (1982) Reworking of sialic crust as represented in late Archean-age gneisses, northern Labrador. Geology **10**, 202-208.
14. Zindler, A., Hart, S.R., and Brooks, C. (1981) The Shabogamo intrusive suite, Labrador: Sr and Nd isotopic evidence for contaminated mafic magmas in the Proterozoic. Earth Planet. Sci. Lett. **54**, 217-235.