

NASA Technical Memorandum 87068

NASA-TM-87068 19850023843

Shock Tube Measurements of Growth Constants in the Branched-Chain Ethane-Carbon Monoxide-Oxygen System

Richard S. Brokaw
*Seven Beacon Hill
Strafford, Vermont*

and

Theodore A. Brabbs and Christopher A. Snyder
*Lewis Research Center
Cleveland, Ohio*

July 1985

LIBRARY COPY

SEP 23 1985

LANGLEY RESEARCH CENTER
LIBRARY, NASA
HAMPTON, VIRGINIA

NASA



NF00147

SHOCK TUBE MEASUREMENTS OF GROWTH CONSTANTS IN THE BRANCHED-CHAIN

ETHANE-CARBON MONOXIDE-OXYGEN SYSTEM

Richard S. Brokaw
Strafford, Vermont

Theodore A. Brabbs and Christopher A. Snyder
National Aeronautics and Space Administration
Lewis Research Center
Cleveland, Ohio 44135

SUMMARY

Exponential free-radical growth constants have been measured for ethane-carbon monoxide-oxygen mixtures by monitoring the growth of oxygen atom concentration as manifested by CO flame-band emission. Data were obtained over the temperature range of 1200 to 1700 K.

The data were analyzed using an ethane oxidation mechanism involving seven elementary reaction steps. Calculated growth constants were close to experimental values at lower temperatures, up to about 1400 K, but at higher temperatures computed growth constants were considerably smaller than experiment. In attempts to explain these results additional branching reactions were added to the mechanism. However, these additional reactions did not appreciably change calculated growth constants.


INTRODUCTION

Ethane is a simple hydrocarbon and is also an intermediate in the rich combustion of other hydrocarbons. Thus, in order to obtain a complete set of chemical reactions and rates for modeling hydrocarbon combustion it is desirable to know the pathways by which ethane reacts.


The shock tube has proved useful for obtaining elementary reaction rates from growth constants in the H₂-CO-O₂ (ref. 1) and CH₄-CO-O₂ (ref. 2) systems. Consequently an analogous study of the C₂H₆-CO-O₂ system was initiated in hopes of extracting elementary reaction rates for ethane oxidation.

Ethane Oxidation Scheme


When a mixture containing carbon monoxide, oxygen, and a small amount of ethane is subjected to a temperature and pressure pulse in a shock tube, small concentrations of atoms and free radicals may first be formed by processes such as






and



followed by



These atoms and free radical concentrations may then grow exponentially via the branched chain scheme



Reactions have been numbered to be compatible with notation used in the study of the $\text{H}_2\text{-O}_2\text{-CO}$ system (ref. 1)

Theory shows (refs. 1 to 5) that in such chain branched systems when depletion of reactants is negligible and temperature and pressure are constant, the atom and radical concentrations increase proportional to $\exp(\lambda t)$ (except very early in the reaction). The growth constant λ depends on the rate constants of the elementary chemical reactions and the concentrations of stable reactants - in this case CO , O_2 , and C_2H_6 .

This report presents experimental growth constants measured behind incident shocks for a range of pressures, temperatures, and gas compositions. These experimental growth constants are compared with values computed using rate constants from the literature. Calculated growth constants are close to experimental values at temperature below 1400 K, but at higher temperatures

computed growth constants are considerably smaller than the experimental values. Attempts to explain this result using additional chain branching reactions have not been successful.

EXPERIMENTAL ASPECTS

Growth constants were obtained by measuring the blue CO flame band emission behind incident shocks. The intensity of this radiation is proportional to the product of carbon monoxide and oxygen atom concentrations (ref. 6), and since little CO is consumed the light measures the increase of oxygen atom concentration with time. Details of the shock tube and associated optical and electronic equipment have been described elsewhere (ref. 7). Flame band radiation centered at 4100 Å was isolated by a high precision ultraviolet interference filter having a 100 Å band pass.

Gas mixtures contained small amounts of C₂H₆ with varying amounts of CO, O₂, and CO₂ diluted with argon. (Carbon dioxide was added to ensure vibrational relaxation of CO in mixtures with a large CO concentration.) Oxygen, carbon dioxide, ethane, carbon monoxide, and argon were high purity tank gases and were used without further purification.

It has been shown (refs. 7 and 8) that boundary-layer effects must be considered in analyzing data obtained behind incident shocks. Conditions behind the shocks, in the region of the experimental measurements, were obtained from a computer program (ref. 9) which integrates the equations of chemical change for a shocked gas accounting for the effects of boundary-layer buildup. The procedure has been described for both turbulent (ref. 7) and laminar (ref. 8) boundary layers. The data in this report were obtained with laminar boundary layers.

Exponential growth constants were obtained from plots of logarithm of observed light intensity versus gas time. The relation between gas and laboratory times was obtained from the computer calculations.

THEORETICAL CONSIDERATIONS

The analytic solutions of the differential equations describing the ignition kinetics in chain-branched systems involving hydrogen, oxygen, and carbon monoxide have been discussed in detail previously (refs. 3 to 5) and the solution has been presented for the methane-carbon monoxide-oxygen system (ref. 2). The solution for the system of reactions presented in the introduction is quite analogous and will merely be outlined here.

To obtain the analytic solution, the atom and radical concentrations - H, OH, O, and C₂H₅ - are assumed to be small in comparison to the concentrations of reactants C₂H₆, CO, and O₂ so that the reactant concentrations may be considered constant and reactions between chain carriers are unimportant. In addition, the temperature and pressure must be nearly constant over the time range of the experimental observations.

Subject to these assumptions, the chemical kinetics are described by a system of simultaneous first-order linear differential equations, one for each chain carrier. The solution of these equations shows that during the growth

period the chain-carrier concentrations grow as $\exp(\lambda t)$ (except very early in the reaction). During the growth period only reactions (2) to (10) are important. The growth constant λ is the positive root of a polynomial of a degree equal to the number of chain carriers. Thus with H, OH, O, and C_2H_5 as chain carriers, the following quartic equation is obtained:

$$\begin{vmatrix} -(v_9 + v_{10} + \lambda) & v_8 & v_6 & v_7 \\ 0 & -(v_8 + \lambda) & 0 & v_2 \\ 0 & v_8 & -(v_5 + v_6 + \lambda) & v_2 \\ v_{10} & 0 & v_5 & -(v_2 + v_7 + \lambda) \end{vmatrix} = 0 \quad (1)$$

where $v_2 \equiv k_2[O_2]$, $v_5 \equiv k_5[CO]$, $v_6 \equiv k_6[C_2H_6]$, $v_7 \equiv k_7[C_2H_6]$, $v_8 \equiv k_8[C_2H_6]$, $v_9 \equiv k_9[O_2]$, and $v_{10} \equiv k_{10}[M]$. The rows in this equation derive from the differential equations for the rates of formation of C_2H_5 , O, OH, and H, respectively. The columns derive from the rates of formation or destruction of C_2H_5 , O, OH, and H in the elementary chemical reactions.

This equation was used to calculate growth constants corresponding to the experimental conditions using rate constants taken from the literature. The rate constants and sources are listed in the appendix. Equation (1) was numerically differentiated to obtain the sensitivities of the calculated rate constants to the various reaction rates.

RESULTS AND DISCUSSION





The compositions of the three gas mixtures investigated are given in table I. Also shown are the sensitivities, $\partial \ln \lambda / \partial \ln v_i$ for the various reaction rates. The range of sensitivities shown is for the temperature range of the data for each composition. Mixture 1 shows sensitivity to the rates of reactions 2, 5, 6, and 8 while mixtures 2 and 3 are sensitive mainly to reactions 2 and 8.

The experimental results are set forth in table II and plotted in figures 1 to 3. Because the reactions in the ethane oxidation scheme are all bimolecular, the growth constant at a given temperature should be proportional to the pressure. Consequently, in figures 1 to 3 the experimental data are plotted as the logarithm of λ/P versus reciprocal temperature. Also shown as solid lines are values computed from equation (1) using rate constants taken from the literature, and set forth in the appendix.

For mixture 1 calculated growth constants are 25 to 50 percent lower than experimental values, with the larger discrepancies at the higher temperatures. For mixtures 2 and 3 calculation and experiment are in better agreement, particularly at temperatures below about 1400 K; but at the highest temperatures calculated values are 20 to 50 percent below experiment.




The agreement between theory and experiment can be improved by upward adjustments in k_5 and k_8 . However, the rate of reaction 5 is quite well established, and the increases required - fivefold or more - cannot be justified.

Other reactions were added to the scheme in hopes of improving the agreement between calculated and experimental growth constants. Additional reactions considered included



The introduction of another chain carrier - HO_2 or HCO - in schemes A to D increased mathematical complexity by transforming equation (1) to a quintic. However, these schemes had negligible effect on calculated growth constants.

Finally another reaction between atomic oxygen and ethane was added to the basic oxidation scheme:




Growth constants calculated assuming equal rate for reactions 8 and 8(a) were 23 to 24 percent larger for mixture 1 over the experimental temperature range. Although this is an improvement, it is not nearly large enough at the higher temperatures. If a larger activation energy were assumed for reaction 8(a) so as to increase the rate of change of growth constant with temperature, the rate constants for reaction 8(a) would be smaller and the increases in calculated growth constants would be less than 24 percent.

Thus a completely satisfactory theoretical analysis of the experimental growth constants has not been achieved. Perhaps the assumptions of the linearized analysis - negligible depletion of reactants, negligible heat release, negligible radical-radical reactions - are not valid at the higher temperatures.


APPENDIX

SPECIFIC REACTION RATE CONSTANTS


This appendix lists individual reactions used in calculations and indicates values and sources of the assumed rate constants. Bimolecular rate constants are expressed in $\text{cm}^3\text{mole}^{-1}\text{sec}^{-1}$. Temperatures are in degrees Kelvin and activation temperatures, E/R, are in degrees Kelvin.




This rate constant, from reference 8, was measured in the shock tube used in this investigation.




This rate constant (ref. 1) was determined in the shock tube used in this investigation.




Rate constant from reference 10.




Rate constant from reference 10.



Rate constant from reference 11.



Rate constant from reference 12.



Rate constant from reference 13.

REFERENCES

1. Brabbs, T.A.; Belles, F.E.; and Brokaw, R.S.: Shock Tube Measurements of Specific Reaction Rates in the Branched-Chain Hydrogen-Carbon Monoxide-Oxygen System. Thirteenth Symposium (International) on Combustion, Combustion Institute, 1971, pp. 129-135.
2. Brabbs, T.A.; and Brokaw, R.S.: Shock Tube Measurements of Specific Reaction Rates in the Branched Chain $\text{CH}_4\text{-CO-O}_2$ System. Fifteenth Symposium (International) on Combustion, Combustion Institute, 1975, pp. 893-900.
3. Brokaw, R.S.: Analytic Solutions to the Ignition Kinetics of the Hydrogen-Oxygen Reaction. Tenth Symposium (International) on Combustion, Combustion Institute, 1965, pp. 269-277.
4. Brokaw, R.S.: Ignition Kinetics of the Carbon Monoxide-Oxygen Reaction. Eleventh Symposium (International) on Combustion, Combustion Institute, 1967, pp. 1063-1072.
5. Brokaw, R.S.: Rate of Reaction Between Molecular Hydrogen and Molecular Oxygen. NASA TM-X-2707, 1973.
6. Clyne, M.A.A.; and Thrush, B.A.: Mechanism of Chemiluminescent Combination Reactions Involving Oxygen Atoms. Proc. Roy. Soc. (London), Ser. A, vol. 269, no. 1338, Sept. 25, 1962, pp. 404-418.
7. Belles, F.E.; and Brabbs, T.A.: Experimental Verification of Effects of Turbulent Boundary Layers on Chemical-Kinetic Measurements in a Shock Tube. Thirteenth Symposium (International) on Combustion, Combustion Institute, 1971, pp. 165-175.
8. Brabbs, T.A.; and Belles, F.E.: Experimental Study of Effects of Laminar Boundary Layers on Chemical Kinetic Measurements in a Shock Tube. Shock Tube Research. J.L. Stollery, A.G. Gaydon, and P.R. Owen, eds., Chapman and Hall, London, pp. 24-1 to 24-9.
9. Bittker, D.A.; and Scullin, V.J.: GCKP84 General Chemical Kinetics Code for Gas-Phase Flow and Batch Processes Including Heat Transfer Effects, NASA TP-2320, Sept. 1984.
10. Baker, R.R.; Baldwin, R.R.; and Walker, R.W.: Use of the Hydrogen + Oxygen Reaction in Determining The Velocity Constants of Elementary Reactions in Hydrocarbon Oxidation. Thirteenth Symposium (International) on Combustion, Combustion Institute, 1971, pp. 291-299.
11. Tanzawa, T.; and Klemm, R.B.: 179th National Meeting of the American Chemical Society, Houston, Texas, March 1980. Cited by Michael, J.V., Nineteenth Symposium (International) on Combustion, Combustion Institute, 1982, p. 59.
12. Baldwin, R.R.: Edinburgh Gas Kinetics Conference, 1975. Cited by Olson, D.B., and Gardiner, W.C. Jr., Combustion and Flame, Vol. 32, pp. 151-161, (1978).
13. Glaenger, K.; and Troe, J.: Thermal Decomposition Reactions of Nitro Compounds in Shock Waves. Helv. Chim. Acta, vol. 56, no. 2, 1973, pp. 577-585.

TABLE I. - MIXTURE COMPOSITION AND GROWTH CONSTANT SENSITIVITIES

Composition, mole percent	Gas mixture		
	1	2	3
C ₂ H ₆	0.0215	0.0190	0.0204
CO	3.04	10.33	10.08
O ₂	9.90	2.87	1.034
CO ₂	2.74	5.05	5.04
Calculated sensitivities, reaction i			
2	0.19 to 0.11	0.50 to 0.37	0.56 to 0.49
5	.25 to .26	.10 to .17	.06 to .10
6	.09 to .15	.02 to .03	.005 to .02
7	-.005 to -.001	-.02 to -.001	-.008 to -.001
8	.46 to .47	.41 to .43	.38 to .40
9	-.10 to .02	-.07 to -.006	-.015 to -.003
10	.12 to .03	.09 to .011	.025 to .006

TABLE II. - EXPERIMENTAL RESULTS

Temperature, K	Pressure, atm	Growth constant λ , sec^{-1}
Mixture 1		
1699	0.1642	17.11×10^3
1683	.1421	17.02
1677	.1485	17.41
1644	.1590	18.48
1617	.1584	16.28
1605	.1292	13.28
1603	.1340	12.06
1580	.1343	10.01
1483	.1347	9.29
1472	.1260	6.73
1409	.1160	7.63
1390	.1092	4.64
Mixture 2		
1602	0.1328	8.82×10^3
1558	.1292	8.20
1535	.1417	10.52
1529	.1510	9.32
1498	.1225	5.91
1477	.1310	7.15
1447	.1229	5.06
1438	.1252	4.31
1409	.1366	3.79
1377	.1188	3.91
1359	.1244	3.07
1337	.1152	3.45
1315	.1170	3.18
1287	.1136	2.46
1221	.1064	1.759
1208	.1035	1.918

TABLE II. - CONCLUDED

Temperature, K	Pressure, atm	Growth constant λ , sec ⁻¹
Mixture 3		
1605	0.1323	8.59x10 ³
1565	.1396	8.25
1536	.1603	7.68
1529	.1278	4.94
1501	.1501	6.03
1463	.1173	2.92
1443	.1272	4.01
1438	.1591	4.91
1432	.1270	3.15
1416	.1406	3.90
1408	.1363	3.07
1388	.1459	2.69
1355	.1157	1.803
1342	.1338	2.12
1321	.1301	2.39
1320	.1201	2.34




Figure 1. - Experimental and computed growth constants as a function of temperature, Mixture 1.




Figure 2. - Experimental and computed growth constants as a function of temperature. Mixture 2.




Figure 3. - Experimental and computed growth constants as a function of temperature. Mixture 3.

1. Report No. NASA TM-87068	2. Government Accession No.	3. Recipient's Catalog No.	
4. Title and Subtitle Shock Tube Measurements of Growth Constants in the Branched-Chain Ethane-Carbon Monoxide-Oxygen System		5. Report Date	
		6. Performing Organization Code 505-31-04	
7. Author(s) Richard S. Brokaw, Theodore A. Brabbs, and Christopher A. Snyder		8. Performing Organization Report No. E-2574	
		10. Work Unit No.	
9. Performing Organization Name and Address National Aeronautics and Space Administration Lewis Research Center Cleveland, Ohio 44135		11. Contract or Grant No.	
		13. Type of Report and Period Covered Technical Memorandum	
12. Sponsoring Agency Name and Address National Aeronautics and Space Administration Washington, D.C. 20546		14. Sponsoring Agency Code	
		15. Supplementary Notes Richard S. Brokaw, Seven Beacon Hill, Strafford, Vermont; Theodore A. Brabbs and Christopher A. Snyder, NASA Lewis Research Center.	
16. Abstract <p>Exponential free-radical growth constants have been measured for ethane-carbon monoxide-oxygen mixtures by monitoring the growth of oxygen atom concentration as manifested by CO flame-band emission. Data were obtained over the temperature range of 1200 to 1700 K. The data were analyzed using an ethane oxidation mechanism involving seven elementary reaction steps. Calculated growth constants were close to experimental values at lower temperatures, up to about 1400 K, but at higher temperatures computed growth constants were considerably smaller than experiment. In attempts to explain these results additional branching reactions were added to the mechanism. However, these additional reactions did not appreciably change calculated growth constants.</p>			
17. Key Words (Suggested by Author(s)) Kinetics Shock tube Ethane		18. Distribution Statement Unclassified - unlimited STAR Category 25	
19. Security Classif. (of this report) Unclassified	20. Security Classif. (of this page) Unclassified	21. No. of pages	22. Price*

End of Document