

General Disclaimer

One or more of the Following Statements may affect this Document

- This document has been reproduced from the best copy furnished by the organizational source. It is being released in the interest of making available as much information as possible.
- This document may contain data, which exceeds the sheet parameters. It was furnished in this condition by the organizational source and is the best copy available.
- This document may contain tone-on-tone or color graphs, charts and/or pictures, which have been reproduced in black and white.
- This document is paginated as submitted by the original source.
- Portions of this document are not fully legible due to the historical nature of some of the material. However, it is the best reproduction available from the original submission.

TIME DEPENDENT EFFECTS
AT THE
FOOT OF THE CHROMOSPHERE

NASA Grant NAGW-253

Final Report

For the period 1 November 1981 - 31 October 1984

Principal Investigator

Wolfgang Kalkofen

November 1984

Prepared for:

National Aeronautics and Space Administration
Washington, D.C. 20546

Smithsonian Institution
Astrophysical Observatory
Cambridge, Massachusetts 02138



The Smithsonian Astrophysical Observatory
is a member of the
Harvard-Smithsonian Center for Astrophysics

The NASA Technical Officer for this grant is Dr. D. Bohlin - Code EZ-7, Solar and Heliospheric Physics Office, NASA Headquarters, Washington, D.C. 20546.

(NASA-CR-174143) TIME DEPENDENT EFFECTS AT
THE FOOT OF THE CHROMOSPHERE Final Report
(Smithsonian Astrophysical Observatory) 4 p
HC A02/MF A01 CSCL 03B

N85-13762

Unclas

G3/92 24521

TIME DEPENDENT EFFECTS
AT THE
FOOT OF THE CHROMOSPHERE

The NASA grant NAGW-253 was awarded for a two-year period ending (with an extension) on October 31. It resulted in the publication or submission for publication of ten (10) scientific papers, dealing either with the structure of the upper solar photosphere near the temperature minimum or with numerical methods for solving the radiative transfer equation.

The purpose of the research was to study an apparent failure of empirical models to conserve energy and to resolve inconsistencies in the empirical temperatures at the temperature minimum. The model that was proposed as an explanation of the observed phenomena was a dynamical model, in which acoustic waves generated in the hydrogen convection zone travelled outward with amplitudes increasing nearly exponentially as a function of height. In the temperature minimum region, the amplitudes became sufficiently large for time averages of the fluctuating temperature to be significantly different from the ambient temperature. Thus, the mean temperature was depressed below the ambient temperature, and the apparent temperatures inferred from the time-averaged fluxes of lines in the ultra-violet part of the spectrum were elevated, with the enhancement increasing with the optical frequency of the line transition. These effects, which had been predicted, were indeed found. However, their magnitude was much less than what was needed in order to remove the difficulties in the empirical models.

The failure of the dynamical model is attributed to the assumption that the solar atmosphere is heated by plane waves emerging (horizontally) uniformly over the solar surface and with the energy flux required by the observed emission rate from the chromosphere. This flux gives wave amplitudes that are too small to explain the separation of the empirical temperatures observed in the lines of ionized calcium and magnesium, for example.

The remedy of the dynamical model is to require an energy flux that is large enough to lead to amplitudes for which the apparent temperatures match the empirical line temperatures in the UV lines. Consistency with the observations of chromospheric emission then requires that the emission take place not uniformly over the solar surface but in smaller spatial regions.

This modified dynamical model, which is suggested by the numerical modelling performed with the support of the NASA grant, is in agreement with the observations, which suggest that the solar chromosphere is heated preferentially in magnetic regions, such as magnetic flux tubes. In this case

the mechanical waves that heat the chromosphere are not pure acoustic modes but are slow-mode mhd waves.

Thus, the modified dynamical model of chromospheric heating is based on the generation of slow-mode waves in magnetic regions in the hydrogen convection zone and their propagation outward along magnetic flux tubes, with shocks forming in the low solar chromosphere. The properties of these waves are such that more energy flux can be generated in the convection zone, less is lost through radiation damping, and shock formation is delayed by the propagation in the flux tube geometry. This dynamical model has an excellent chance of describing the heating of the solar chromosphere. It will be proposed for study under another grant, to be submitted to NASA. Suffice it to note here merely that it is identical to the model discussed at a recent workshop held at Sacramento Peak Observatory, which is credited with having the best chance of explaining the chromospheric temperature structure.

IS ENERGY CONSERVED AT THE FOOT OF THE SOLAR CHROMOSPHERE?

(W. Kalkofen),
in: Second Cambridge Workshop on Cool Stars, Stellar Systems, and the Sun,
M. S. Giampapa and L. Golub, eds.
SAO Spec. Rep. 392, 1982.

CORE SATURATION IN A MOVING MEDIUM

(W. Kalkofen and P. Ulmschneider),
in: Methods in Radiative Transfer, W. Kalkofen, ed., Cambridge University
Press, 1984, p131.

APPARENT SOLAR TEMPERATURE ENHANCEMENT DUE TO LARGE AMPLITUDE WAVES

(W. Kalkofen, P. Ulmschneider and F. Schmitz),
Ap. J., Dec. 1984, in press.

ACOUSTIC WAVES IN THE SOLAR ATMOSPHERE VII; NON-GREY NON-LTE H- MODELS

(F. Schmitz, P. Ulmschneider and W. Kalkofen),
submitted to Astronomy and Astrophysics.

RADIATIVE TRANSFER IN AN EXPANDING SPHERICAL MEDIUM

(W. Kalkofen and R. Wehrse),
in: Cool Stars, Stellar Systems, and the Sun,
Springer Verlag, Berlin, 1984,
L. Baliunas and L. Hartmann, eds., p306.

RADIATIVE TRANSFER IN SPHERICAL ATMOSPHERES

(W. Kalkofen and R. Wehrse),
in: Methods in Radiative Transfer, W. Kalkofen, ed., Cambridge University
Press, 1984, p307.

RESONANCE LINES IN DUSTY GASEOUS NEBULAE

(R. Wehrse and W. Kalkofen),
submitted to Astronomy and Astrophysics.

AYRES' BIFURCATED SOLAR MODEL

(W. Kalkofen),
to appear in: Sacramento Peak Workshop on Chromospheric Structure and
Modeling, B. Lites, ed., 1985.

THE EQUILIBRIUM STRUCTURE OF THIN MAGNETIC FLUX TUBES I

(A. Ferrari, S. Massaglia, W. Kalkofen, R. Rosner, and G. Bodo),
in preparation.

THE EQUILIBRIUM STRUCTURE OF THIN MAGNETIC FLUX TUBES II

(W. Kalkofen, R. Rosner, A. Ferrari, and S. Massaglia),
in preparation.