

General Disclaimer

One or more of the Following Statements may affect this Document

- This document has been reproduced from the best copy furnished by the organizational source. It is being released in the interest of making available as much information as possible.
- This document may contain data, which exceeds the sheet parameters. It was furnished in this condition by the organizational source and is the best copy available.
- This document may contain tone-on-tone or color graphs, charts and/or pictures, which have been reproduced in black and white.
- This document is paginated as submitted by the original source.
- Portions of this document are not fully legible due to the historical nature of some of the material. However, it is the best reproduction available from the original submission.

AgRISTARS

"Made available under NASA sponsorship
in the interest of early and wide dis-
semination of Earth Resources Survey
Program information and without liability
for any use made thereof."

Early Warning and Crop Condition Assessment

E83-10377
OR-172921
EW-U3-04405

A Joint Program for
Agriculture and
Resources Inventory
Surveys Through
Aerospace
Remote Sensing

February 1983

RAIN-INDUCED SPRING WHEAT HARVEST LOSSES

ARMAND BAUER AND A. L. BLACK



USDA-ARS
NORTHERN GREAT PLAINS RESEARCH CENTER
MANDAN, ND 58554



(E83-10377) RAIN-INDUCED SPRING WHEAT
HARVEST LOSSES (Agricultural Research
Service) 34 p HC A03/MP A01 CSCL 02C

N83-32130

Unclas
G3/43 00377



Lyndon B. Johnson Space Center
Houston, Texas 77058

1. Report No. EW-U3-04405		2. Government Accession No.		3. Recipient's Catalog No.	
4. Title and Subtitle Rain-induced Spring Wheat Harvest Losses				5. Report Date February 1983	
				6. Performing Organization Code	
7. Author(s) Armand Bauer and A. L. Black				8. Performing Organization Report No.	
				10. Work Unit No.	
9. Performing Organization Name and Address USDA-ARS Northern Great Plains Research Center P.O. Box 459 Mandan, ND 58554				11. Contract or Grant No.	
				13. Type of Report and Period Covered Technical Report	
12. Sponsoring Agency Name and Address USDA -ARS Early Warning/Crop Condition Assessment Project 1050 Bay Area Blvd., Houston, TX 77058				14. Sponsoring Agency Code	
15. Supplementary Notes					
16. Abstract <p>When rain or a combination of rain and high humidity delay wheat harvest, losses can occur in grain yield and/or grain quality. Yield losses can result from shattering, from reduction in test weight, and in the case of windrowed grain, from rooting of sprouting grain at the soil: windrow contact. Losses in grain quality can result from reduction in test weight and from sprouting. Sprouting causes a degradation of grain proteins and starches, hence flour quality is reduced, and the grain price deteriorates to the value of feed grain.</p> <p>Although losses in grain yield and quality are rain-induced, these losses do not necessarily occur because a standing or windrowed crop is wetted by rain. Spike water concentration in hard red spring wheat must be increased to about 45-49% before sprouting is initiated in grain that has overcome dormancy. The time required to overcome this dormancy after the cultivar has dried to 12-14% water concentration differs with hard red spring cultivars.</p> <p>This study was conducted to evaluate and compare the effect of rain on threshing-ready standing and windrowed hard red spring wheat grain yield and quality. A goal is to develop capability to forecast the extent of expected loss of grain yield and quality from specific climatic events that delay threshing.</p>					
17. Key Words (Suggested by Author(s)) primary dehydration of wheat water imbibition rate spike wetting rate threshold spike water concentration			18. Distribution Statement		
19. Security Classif. (of this report)		20. Security Classif. (of this page)		21. No. of Pages	22. Price*

EW-U3-04405

RAIN-INDUCED SPRING WHEAT HARVEST LOSSES

PRINCIPAL INVESTIGATORS

Armand Bauer and A. L. Black

APPROVED BY

**Glenn O. Boatwright, Manager
Early Warning/Crop Condition Assessment Project
AgRISTARS Program**

**Houston, Texas
February 1983**

RAIN-INDUCED SPRING WHEAT HARVEST LOSSES

Armand Bauer and A. L. Black^{1/} ^{2/}

^{1/} Soil Scientists, Northern Great Plains Research Center, Mandan, ND.

^{2/} Funds for the project were provided by NASA (Johnson Space Center)
and USDA AgRISTARS EN/CCA.

Rain-Induced Spring Wheat Harvest Losses

Armand Bauer and A. L. Black

When rain or a combination of rain and high humidity delay wheat harvest, losses can occur in grain yield and/or grain quality. Yield losses can result from shattering, from reduction in test weight, and in the case of windrowed grain, from rooting of sprouting grain at the soil:windrow contact. Losses in grain quality can result from reduction in test weight and from sprouting. Sprouting causes a degradation of grain proteins and starches, hence flour quality is reduced (Gordon et al., 1977), and the grain price deteriorates to the value of feed grain.

Climatic conditions in North Dakota during harvest of hard red spring wheat are normally characterized by low rainfall, low humidity, and warm temperatures (Ramirez, 1972; 1973). These conditions favor rapid grain drying with little chance for biological damage to the grain. However, there is a 50% probability of receiving at least 0.1 inch rain per week during weeks 30 through 35 (last week in July to September) and a 15% probability for 0.60 inches during the same time period (Ramirez, 1973). Also, seeding occasionally is delayed in the spring, causing harvest to be extended into a cooler period of the year and hence a less favorable drying period.

Although losses in grain yield and quality are rain-induced, these losses do not necessarily occur because a standing or windrowed crop is wetted by rain (Wellington and Durham, 1958). Spike water

concentration in hard red spring wheat must be increased to about 45-49% before sprouting is initiated in grain that has overcome dormancy (Bauer and Black, 1983). The time required to overcome this dormancy after the cultivar has dried to 12-14% water concentration differs with hard red spring cultivars (Bauer and Black, 1983).

The spike can act as a water reservoir from which the grain can imbibe (absorb) water. The amount of water absorbed by saturated vegetative tissue of the spike (glumes, rachis, etc.) and in the interstitial areas of the spikelets is sufficient to increase the water concentration in the grain by 42 percentage units, assuming all the water is transferred from the vegetative parts to the grain. The grain makes up about 72% of the dry matter of the spike of hard red spring wheat (Bauer and Black, 1983).

Timgalen wheat grain (Australian soft white wheat) absorbed water at a rate of about 1.9 percentage units per hour in a linear manner from 14% water concentration to saturation at about 100% concentration (Gordon et al. 1977) when the spikes were misted 5 minutes every hour and in the interim kept in a high relative humidity environment. Water absorption rate by the spike (head) was about 6 times more rapid than the absorption rate by the grain over the first 10 hours of wetting. Spikes, including grain, became saturated at about 130-150% water concentration, oven-dry basis. The spikes therefore required about 11 to 12 hours to become saturated from an initial water concentration of about 14%. Spike water absorption rate of Wared hard red spring wheat was about 1 percentage unit per minute over a 50-minute period (Bauer and Black, 1983).

Grain of Olaf wheat absorbed water at a rate of 1.7 percentage units per hour in spikes equilibrated for 15 hours after being misted to 145% water concentration (Bauer and Black, 1983).

Hard red spring wheat is harvested by either the swath-combine method or by straight combining (simultaneous cutting and threshing) in the northern Great Plains of the United States and Canada. The advantages provided by the swath-combine method over straight combining are: (a) the crop is better protected from wind, hail, and frost when it is in windrows, b) green weeds are eliminated as a threshing problem, c) grain water concentration is equalized where field ripening is not uniform because of soil and topographic differences, d) the cost to artificially dry grain to assure safe storage is eliminated or reduced, and e) potential losses from sawfly infestations are reduced (Dodds, 1967). But the swath-combine method requires additional equipment, such as a swather and combine pickup attachment, as well as labor and fuel to operate the swather.

This study was conducted to evaluate and compare the effect of rain on threshing-ready standing and windrowed hard red spring wheat grain yield and quality. A goal is to develop capability to forecast the extent of expected loss of grain yield and quality from specific climatic events that delay threshing.

MATERIALS AND METHODS

Field experiments were conducted in 1979, 1980, and 1981 on privately-owned farm fields. The swath-combine and straight combine methods of harvesting were compared each year by measuring grain yield, test weight, and grain nitrogen (protein) concentration. Comparisons also were made among three swathing stubble heights in 1980, and between two swathing widths in 1981. Other measurements were made some years which are described under the specific year. Farm-sized swathers and combines were used in all harvesting procedures.

The combine-threshed grain was collected in bags, oven-dried at 156°F, cleaned, and weighed. Yield measurements were based on the total quantity from each plot calculated at 60 pounds per bushel. Test weight measurements were based on a quart volume randomly removed from the total sample. Grain nitrogen concentration was measured by a micro-Kjeldahl procedure (Shuman et al., 1973). All measurements are expressed in terms of oven-dry grain.

Following threshing, a square meter (1.2 square yards) area was vacuumed in the center of each plot to pick up grain that had fallen to the soil surface. The kernels were separated from the soil and straw, oven-dried, then counted and weighed.

When rainfall did not occur to wet the crop, water was sprinkler irrigated on the windrows and the standing crop to simulate rain. The amount of water applied by sprinkler was measured with cans placed on the area blanketed by the sprinklers. Rain was measured with a standard rain gauge.

1979

Plots were established on September 7 in a field seeded to a mixture of several hard red spring wheat cultivars. Each plot of standing and windrowed crop was an area 10.5 feet wide and 60 feet long. The swath stubble height was about 10 inches. Each treatment of six threshing dates and two harvesting methods (straight and swath-combine) was repeated three times. Except for 0.82 inches rain, all water applied on a given day was sprinkler-irrigated on the plots over a period of 45 minutes at a rate of about an inch per hour (Table 1).

1980

Plots were established in a field of Wared hard red spring wheat. Swathed plots were cut on August 7 at stubble heights of 4, 9, and 14 inches. Each plot of standing and windrowed crop was an area 10.5 feet wide and 45 feet long. Each treatment of six threshing dates and of harvesting methods (straight and swath-combine of three stubble cutting heights) was repeated three times. However, because of threshing problems, data of the first threshing date are not included in the text. The plot areas were wetted with more than 10 inches of water, with slightly more than 6 inches from rain (Table 1). Water sprinkled on the plots on a given day was applied over a period of 45 minutes at a rate of about 1.5 inches per hour.

The number of damaged kernels (sprouted and discolored) was determined from a 0.5 pint subsample randomly removed from the total sample. Each kernel was examined under magnification. The kernel was classed as sprouted when the coleoptile was visible, and as

discolored when dark blemishes were observed. When the kernel was both sprouted and discolored it was classed as sprouted. The examination of kernels was made by one person.

1981

The plots were established in a field of Olaf hard red spring wheat. Swathed plots were cut on August 18 leaving a stubble height of about 9 inches. All plots were 50 feet long. The standing crop plots and one set of plots of swathed crop were 10.5 feet wide, and the other set of plots of swathed crop was 15 feet wide. Each treatment of six threshing dates and of harvesting methods (straight and swath-combine of two swath widths) was repeated four times.

Water sprinkled on the plots was applied over a 2.5 to 2.75-hour period, usually beginning about 4 o'clock in the afternoon (Table 1). Following the second threshing, the windrows were covered with clear plastic sheeting at about 9:00 am (CDT) on the day after sprinkling. The plastic sheet was left on the windrows for 24 hours after the second threshing date, for 55 hours after the third threshing date, and for 78 hours after each of the fourth and fifth threshing dates.

The number of sprouted and of discolored kernels was determined under magnification from a constant volume subsample of about 200 kernels. The system to determine damage was identical to the one described above in 1980. The same person made the examinations in 1980 and 1981.

Straw water concentration was determined on a subsample collected at threshing, weighing the straw before and after oven drying at 156°F.

RESULTS AND DISCUSSION

The amount of water required to wet a crop is relatively small. To illustrate: the water in the grain and straw of a hard red spring wheat crop yielding 50 bushels per acre, saturated to 150% water concentration, weighs about 9000 pounds. (Straw and grain weight are about equal in a crop which yields 50 bushels per acre; Bauer and Zubriski, 1978). Nine thousand pounds of water is equivalent to 0.04 inches rain falling on one acre. (An inch of water over an acre weighs 113.26 tons). However, a standing crop of spring wheat seeded in 6 to 7 inch rows will not be saturated by 0.04 inches rain because not all the rain will strike the crop. If one were to assume that only 25% strikes the crop and that all the rain striking the crop is absorbed and none is evaporated, then the amount of water needed would be supplied by 0.16 inches rain.

Assuming a rainfall rate slow enough so that no runoff occurs, the amount of rain needed to saturate yard-wide windrows of a crop of various yields, cut with a swather of various widths, is shown in Figure 1. The amount of rain required to saturate a windrow is greater than is required for the standing crop because the windrow is concentrated in a smaller area. The quantity of tissue in a windrow will vary with the quantity produced per unit area, stubble cutting height, and with swath width. A yard-wide windrow of a 30 bushel per acre crop cut with an 18-foot swather can be saturated with about 0.21 inches rain, assuming all the rain striking the windrow is absorbed and none is evaporated. (Highest recorded state average wheat yield in North Dakota is 28.6 bushels per acre in 1971; Smith, 1978).

Grain Yield

Harvested grain yields differed between the straight and swath-combining methods when the windrows were formed with a 10.5-foot swath cut at about a 9-inch height only in 1979 (Table 2), and then only at the first and sixth threshing dates. On the first date, yields were higher on straight combined plots and on the sixth date on windrowed plots. (The low yield in the windrow at the first threshing date is attributed to experimental error). The amount of grain loss by shattering in 1979, as measured by the amount vacuumed from the plots (Table 3), generally reflected the same trend as the threshed yield in that shattering losses were highest with straight combining by the sixth threshing date.

Shattering losses in Olaf wheat in 1981 also were higher on straight combining than on windrows, at the fourth and fifth threshings (Table 3). But they were lower at the sixth threshing. The reason for the apparent less loss at the sixth threshing date is that more of the grain that had been shattered from the standing crop sprouted and became anchored to the soil, and therefore was not vacuumed up.

Shattering is caused by the transfer of energy to a wheat head by raindrop impact, whether the head is on a standing or windrowed crop. Less shattering is likely in the windrowed crop because fewer heads are exposed to direct hits. Wind too can cause shattering in the standing crop when the speed is great enough to cause heads to strike other heads or stems. The ease of shattering, however, differs among varieties.

Harvested grain yields in 1980 on 4-inch stubble averaged 5.2 bushels per acre higher than on the 9-inch stubble and 8.9 higher than on 14-inch stubble (Table 4), about 14 to 23% difference. Further, yields on 4-inch stubble were consistently higher than on 14-inch stubble at all threshing dates, and higher than on 9-inch stubble on all except the sixth threshing. The lower harvested yields on taller stubble is attributed to more extensive windrow:soil contact, thus allowing more of the sprouting grain to become anchored to the soil. The taller stubble is less rigid and bends more easily under windrow weight allowing more of the windrow to lie on the soil instead of being supported above the soil surface. Also, on the 14-inch stubble, much of the windrow fell through the stubble during the first rain, allowing extensive sprouting and head anchoring to occur. Windrows are pushed into and through stubble by the energy that is transferred from raindrop impact.

The volume and length of straw in the windrows is reduced as stubble cutting height increases. The lower the volume of straw in relation to stubble height the more easily the windrow will be forced down through the upright stubble by rain. In 1980, the average height of Wared wheat was 28 inches hence the proportion of stubble height to straw length was 1:6, 1:2, and 1:1 for 4-inch, 9-inch, and 14-inch stubble, respectively. The volume of tissue in the windrows from a given cutting height will vary with swath width. Hence crop height, stubble cutting height, and swath width (spike population) need to be considered to determine the ideal windrow volume to support on the stubble. However, consideration may need to be given

also to the ratio of material-other-than-grain to grain for maximum efficiency in separating grain from straw as these pass over the combine straw-walker (Reed et al., 1970). In 1980, 4.32 inches rain was received before the first windrows were picked up and threshed (Table 1). The initial 2.02 inches fell in one storm within 3 days after swathing, and this was followed by another 1.10 inches a week after the first storm.

Although the short stubble gave better support to windrows and provided a yield advantage over taller stubble in a wet harvest year like 1980, some of the yield advantage gained at harvest with short stubble may be offset in some succeeding years because short stubble has less snow-trapping capability than tall stubble. With less snow-trapping there is a reduction in the water storage potential for the next crop. This is illustrated from four years data developed at Mandan ^{1/}. On 14-inch stubble, each inch of stubble above 2 inches height increased the average available soil water content 0.15 inches in the upper 3 feet of soil from late autumn to spring seeding, a total of 1.80 inches, while on 8-inch stubble, each inch of stubble above 2 inches height increased the soil water content about 0.09 inches in the upper 3 feet over the same time period. The long-term average contribution to wheat grain yield from an inch of stored soil water at seeding is about 2.4 bushels per acre (Bauer, 1972).

Swath width, 10.5 versus 15 feet, had no effect of grain yield in 1981 (Table 5).

^{1/} Unpublished data, Bauer.

Test Weight

The effect of harvesting method on grain test weight was not consistent with years (Table 6). In 1979 there was no difference between methods. In 1980 the test weight was lower in the windrowed crop at all threshing dates. In 1981 test weights were lower in windrowed crop on the third through sixth threshing dates.

The difference in test weight between harvesting methods likely is associated with differences in rate of drying. The spikes of the standing crop dry faster than the spikes in windrows because of better air circulation. With more rapid drying, less water is imbibed by the grain. Seeds swell when they take up water and remain partially swelled after drying. This irreversible change in seed size is the cause of lowering of test weight (Ciha, 1981). The extent of swelling likely is related to the amount of water absorbed above an unknown minimum amount required to initiate expansion.

Because of the warm, dry weather that prevailed in 1979, the water sprinkled on the windrows and standing crop evaporated rapidly. Hence water uptake by the grain from sprinkling was minimal and no change in test weight occurred.

In 1980, no measurement of test weight was obtained before 4.32 inches of rain wetted the windrows, hence the change that occurred because of the rain is unknown. However, test weight was consistently higher in the standing crop than the windrowed crop. Also, test weight of grain from windrows on the 14-inch stubble was about 0.7 pounds per bushel lower than grain from 9-inch stubble windrows (Table 7), but did not differ from that of the 4-inch

stubble. A larger portion of the windrow on the 14-inch stubble made soil contact, and so drying rate likely was slower than in windrows on the 9-inch stubble.

After the second threshing in 1981, the windrows were covered with clear plastic sheets about 16 hours after wetting, to reduce the rate of drying in the windrows relative to the standing crop. The affect of this treatment is apparent in the lower test weights, shown in data in Table 6. Also, the lower test weight associated with the 15-foot swaths compared to the 10.5-foot swaths in 1981 (Table 5) appear to be a reflection of a slower drying rate in the larger windrows.

These data show that the length of time that water is present to be absorbed by the grain is a major factor in causing a decrease in test weight. A decrease in test weight does not necessarily occur with each rain event. The effect on test weight of length of time water can be absorbed by the grain is provided by the 1980 and 1981 data. Test weight of grain in windrows was lower than in the standing crop because windrows dried more slowly. Water available for absorption by the grain after rainfall ceases includes water adsorbed to and absorbed by spike tissue. In the case of windrows, additional water absorbed by the grain can come from the straw.

Grain Protein

Grain nitrogen concentration (protein) differed between harvesting methods in 1980 (Table 8), but not in 1979 and 1981. The difference at the fourth threshing in 1979 was not found at the other threshing dates. In 1980, the grain nitrogen was lower in the 9-inch stubble swathed crop than standing crop, an average of 0.20

percentage units over the five threshing dates. But, there was no difference in grain nitrogen concentration among the three swath, stubble-cutting heights as threshed in the field (Table 7), that is, before separation of "good" from "damaged" kernels. Further, the concentration was lower in the "good" kernels on the 9-inch stubble and was higher in the "damaged" kernels of the standing crop than any of the windrowed crop (Table 7). No reason can be given for this anomalous outcome.

Within any stubble height, nitrogen concentration in the "damaged" kernels averaged consistently higher than in the "good" kernels. The reason for the higher nitrogen concentration in "damaged" grain is that in the process of sprouting, carbohydrates are consumed in respiration and carbon dioxide is released. With the loss of carbohydrates, the apparent nitrogen concentration increases. Also one could expect protein synthesis taking place as germination begins. Proteins are not consumed during respiration but are utilized in the synthesis of other organic nitrogenous compounds (Meyer and Anderson, 1952).

Swath width, 10.5 versus 15 feet, had no effect on grain nitrogen concentration in 1981 (Table 5).

Grain Damage

Damage to grain, sprouting plus discoloration, was severe in the high rainfall year 1980, and in general increased with delayed threshing (Table 9, 10). At the initial threshing, damage was higher on the 9-inch than either the 4-inch or 14-inch stubbles. On the subsequent threshings, damage was as high or higher in grain on

4-inch stubble as on the taller stubbles. The relatively lower sprouting percentage on the 9- and 14-inch stubbles is likely a reflection of sprouted grain becoming anchored to the soil and not being picked up and included in the threshed grain. Damage was also observed in grain harvested by the straight combine method (Table 10), but the severity of damage was roughly 50% less. This outcome supports the premise that spikes on the standing crop dry faster than spikes in the windrows.

Damage to kernels was less severe in 1981 (Table 10) than 1980. Differences between harvesting methods were not significant until the fourth threshing. Covering the windrows with clear plastic sheets to reduce the drying rate also likely raised the temperature of the windrow and contributed to increased damage. More of the damage was a result of microbial activity, as reflected by discoloration, than sprouting. (Data are not shown).

Grain damage (sprout plus discoloration) of less than 3% difference was statistically significant between the 10.5 and 15-foot swaths in 1981 (Table 5).

Water in Straw

Water concentration in the straw of the 1981 standing crop was consistently higher than in the straw from windrows (Table 11). At the same time, the water concentration in the grain, while differing significantly, differed by no more than two percentage units between the windrowed and standing crop. Apparently the straw of a standing crop continued to absorb water from the soil after it was mature. This can affect the ease of threshing grain to be straight combined.

SUMMARY AND CONCLUSIONS

1. The amount of water needed to saturate a standing crop of wheat yielding 50 bushels per acre, grown in 6 to 7-inch spacing, is about 0.16 inches, provided that all the rain striking the tissue is absorbed by it and there is no loss by evaporation.
2. Yard-wide windrows of a 30-bushel per acre crop cut with an 18-foot swather can be saturated with about 0.21 inches rain, provided there is no runoff from the windrow and no water loss by evaporation.
3. Harvested grain yields differed between straight combining and windrows formed with a 10.5-foot swather, leaving a 9-inch stubble, in one year out of three and only at the sixth threshing date.
4. Harvested grain yields on 4-inch stubble averaged 14% and 23% higher than on 9-inch and 14-inch, respectively. Long stubble, less rigid, allowed a greater opportunity for contact between the windrow and soil. Also, short straw more readily fell between or was driven between the stubble by raindrop impact and thus made contact with the soil. More anchoring (sprouting grain rooting in soil) occurred on 9- and 14-inch stubble than 4-inch stubble.
5. After each rain, the standing crop lost some of its erectness. Hence the required cutting height of straight combined crop was lowered with each date to assure that all spikes were threshed.

6. Test weight of grain differed between straight combining and swathing some years. The reason likely is associated with drying rate. Spikes of the standing crop dried faster than spikes in windrows, likely because of better air circulation. Seeds swell when they take up water and remain at least partially swelled after subsequent drying.
7. Grain protein differed between cutting methods in one year of three. Protein in sprouted kernels was at a higher concentration than in kernels not observed to have sprouted. In the sprouting process, carbohydrates are consumed with respiration and carbon dioxide is released and lost. Proteins are not consumed in respiration. Hence, an apparent increase in protein is measured because of a decrease in carbohydrate concentration.
8. Sprouting was severe in a year when high rainfall occurred after the wheat had ripened in the windrow. In general, severity increased with increased delay in threshing. Sprouting also occurred in grain harvested by the straight combine method, but the severity of damage was about 50% less than in the windrowed crop.
9. Water concentration in the straw of the 1981 standing crop was consistently higher than in the straw in windrows at each threshing date, the smallest difference being 8 percentage units and the largest 27 percentage units. Simultaneously, the water concentration in the grain was higher in standing grain than windrowed by no more than two percentage units at any threshing date.


LITERATURE CITED

- Bauer, Armand. 1972. Effect of water supply and seasonal distribution on spring wheat yields. North Dakota Agr. Exp. Sta. Bull. 490.
- Bauer, Armand and A. L. Black. 1983. The water factor in harvest sprouting of hard red spring wheat. (Submitted to Agronomy Journal).
- Bauer, Armand and J. C. Zubriski. 1978. Hard red spring wheat straw yields in relation to grain yields. Soil Sci. Soc. Amer. J. 42:777-781.
- Ciha, Allan J. 1981. Environmental factors during seed development and their influence on pre-harvest sprouting in wheat. EW-U1-04115, JSC-17157, NASA, AgRISTARS/EW/CCA, Houston, TX.
- Dodds, M. E. 1967. Wheat harvesting -- machines and methods. p. 361-371. In K. F. Nielsen (ed.). Proc. Canadian Centennial Wheat Symposium. Modern Press, Saskatoon, Sask.
- Gordon, I. L., N. F. Derera, and L. N. Balaam. 1977. Selection against sprouting damage in wheat. I. Germination of unthreshed grain, with a standard wetting procedure. Aust. J. Agric. Res. 28:583-596.
- Meyer, Bernard S. and Donald B. Anderson. 1952. Plant Physiology, Second Edition. D. Van Nostrand Company Inc., New York. p. 704.
- Ramirez, J. M. 1972. The agro-climatology of North Dakota. Part 1. Air temperatures and growing degree days. North Dakota Coop. Ext. Bull. No. 15. North Dakota State University, Fargo, ND.

- Ramirez, J. M. 1973. The agro-climatology of North Dakota.
Part 2. Precipitation, forecast, probabilities and rainmaking.
North Dakota Coop. Ext. Bull. No. 16. North Dakota State
University, Fargo, ND.
- Reed, W. B., G. C. Zoerb, and F. W. Bigsby. 1970. A laboratory study
of grain-straw separation. ASAE Paper No. 70-604.
- Shuman, G. E., M. S. Stanley, and D. Knudsen. 1973. Automated total
nitrogen analysis of soil and plant samples. Soil Sci. Soc.
Amer. Proc. 37:480-481.
- Smith, Glenn S. 1978. Changes in North Dakota hard red spring wheat
varieties, 1900-1977. North Dakota Agr. Exp. Sta. Farm Res.
35(5):16-21.
- Wellington, P. S. and V. M. Durham. 1958. Varietal differences in
the tendency of wheat to sprout in the ear. Empire J. Exp.
Agric. 26:47-54.

ORIGINAL PAGE IS
OF POOR QUALITY

Figure 1. Inches of Water Needed to Wet Yard-wide Windrows of Wheat Tissue of Varying Grain Yield and Swath Width to 150% Water Concentration.



ORIGINAL PAGE IS
OF POOR QUALITY

Table 1. Interval between swathing and threshing, and amount of water applied to windrows and standing grain.

Method	Threshing date ^{1/}	Year					
		1979		1980		1981	
		Interval days ^{2/}	Water inches ^{3/}	Interval days ^{2/}	Water inches ^{3/}	Interval days ^{2/}	Water inches ^{3/}
Swath	1	10	0.82 ^{4/}	-	-	6	0.06
	2	13	1.57	18	4.32 ^{5/}	10	1.40
	3	17	2.32	21	5.42 ^{6/}	16	2.34
	4	20	3.08	29	6.76 ^{6/}	22	3.14 ^{9/}
	5	24	3.83	35	7.86 ^{6/}	28	4.39
	6	29	4.58	53	10.03 ^{8/}	34	5.79 ^{10/}
Straight	1	14	0.82	-	-	6	0.06
	2	17	1.57	21	5.42 ^{5/}	10	1.40
	3	20	2.33	28	7.86 ^{6/}	16	2.34
	4	24	3.08	35	8.96 ^{6/}	22	4.26 ^{9/}
	5	28	3.83	40	9.11 ^{7/}	28	5.46
	6	32	4.64	53	10.55 ^{8/}	34	6.91 ^{11/}

1/ See Table 2 for the threshing dates.

2/ From date of swathing to threshing.

3/ Sum of rainfall and water applied by sprinkler system.

4/ Except for the 0.82 inches, all water was applied by sprinkler.

5/ Of this total, 4.32 inches was rain.

6/ Of this total, 5.66 inches was rain.

7/ Of this total, 5.81 inches was rain.

8/ Of this total, 6.15 inches was rain.

9/ Windrows were covered with plastic sheets during a 1.12-inch rain event.

10/ Of this total, 0.19 inches was rain from 4 storms.

11/ Of this total, 1.31 inches was rain from 5 storms.

Table 2. Grain yield comparisons between swath and straight combine methods of cutting hard red spring wheat.

	Year					
	1979		1980		1981	
	Method		Method		Method	
Threshing ^{1/} date	Swath ^{2/} bu/ac ^{6/}	Straight	Swath ^{3/} bu/ac ^{6/}	Straight	Swath ^{4/5/} bu/ac ^{6/}	Straight
1	27.7 d	34.9 a	-	-	27.2 a	28.8 a
2	30.9 bc	32.6 ab	30.9 a	35.0 a	26.7 a	28.1 a
3	32.1 bc	33.1 ab	30.0 a	28.0 a	28.2 a	27.1 a
4	29.6 cd	31.7 bc	32.1 a	32.9 a	26.2 a	23.3 a
5	31.5 bc	31.3 bc	36.4 a	34.6 a	24.1 a	23.1 a
6	30.8 bc	23.7 e	34.5 a	34.8 a	22.0 a	22.9 a

1/ Threshing dates: 1979 Swath: 9/17, 9/20, 9/24, 9/27, 10/01, 10/04;
 1979 Straight: 9/21, 9/24, 9/27, 10/01, 10/04, 10/09;
 1980 Swath: 8/25, 8/28, 9/5, 9/11, 9/29;
 1980 Straight: 8/28, 9/4, 9/11, 9/16, 9/29;
 1981 Swath: 8/24, 8/28, 9/3, 9/9, 9/15, 9/21;
 1981 Straight: 8/24, 8/28, 9/3, 9/9, 9/15, 9/21.

2/ Stubble height 10 inches.

3/ Stubble height 9 inches.

4/ Stubble height 9 inches.

5/ Cut with 10.5 foot swather.

6/ Within a year, yield data followed by the same letter do not differ at the 5% confidence level.

ORIGINAL PAGE IS
 OF POOR QUALITY

Table 3. Grain vacuumed from the ground comparing swathed and straight combine methods of cutting hard red spring wheat.

Threshing ^{1/} date	1979 Method		Year 1980 Method		1981 Method	
	Swath bu/ac	Straight	Swath bu/ac	Straight	Swath bu/ac	Straight
1	4.2 b ^{2/}	2.0 bc	-	-	0.4 e ^{2/}	0.8 e
2	1.6 c	2.8 bc	1.4 a ^{2/}	2.4 a	0.6 e	0.7 e
3	2.8 bc	1.5 c	8.7 a	10.6 a	1.6 de	1.8 de
4	2.0 bc	1.8 c	4.5 a	7.6 a	1.8 de	4.8 ab
5	2.9 bc	3.6 bc	-	-	1.8 de	3.6 bc
6	3.0 bc	6.9 a	16.8 a	10.0 a	5.2 a	2.7 od

^{1/} See Table 2 for threshing dates.

^{2/} Within a year, data followed by the same letter do not differ at the 5% confidence level.

ORIGINAL PAGE IS
OF POOR QUALITY

Table 4. Effect of stubble height on grain yield, 1980.

Threshing date	Stubble height, inches		
	4	9 bu/ac	14
2	36.3 b ^{2/}	30.9 od	29.0 de
3	35.0 bc	30.0 de	29.9 de
4	41.5 a	32.1 bod	30.8 od
5	41.2 a	36.3 b	30.0 de
6	35.8 b	34.5 bc	26.2 e
Average	38.0 a ^{3/}	32.8 b	29.1 b

1/ See Table 2 for threshing dates.

2/ Among these three columns, yield data followed by the same letter do not differ at the 5% confidence level.

3/ Within this row, yield data followed by the same letter do not differ at the 5% confidence level.

ORIGINAL PAGE IS
OF POOR QUALITY

Table 5. Measurement comparisons between windrows from 10.5- and 15-foot swaths, 1981.

<u>Measurement</u>	<u>Swath width, feet</u>	
	<u>10.5</u>	<u>15</u>
Grain yield (bu/ac) ^{1/}	25.7 a	25.7 a
Test weight (lbs/bu)	55.9 a	55.1 b
Grain nitrogen (%N)	3.33 a	3.37 a
Number sprouted (%) ^{2/}	4 b	5 a
Number damaged (%) ^{2/}	10 a	7 b

1/ Within a row, the data followed by the same letter do not differ at the 5% confidence level.

2/ Percent of the total number of kernels in the sample.

ORIGINAL PAGE IS
OF POOR QUALITY

Table 6. Test weight comparisons between swathed and straight combine methods of cutting hard red spring wheat.

Threshing ^{1/} date	Year					
	1979		1980		1981	
	Method		Method		Method	
	Swath	Straight	Swath	Straight	Swath	Straight
	lbs/bu		lbs/bu		lbs/bu	
1	58.1 a ^{2/}	57.6 a	-	-	59.2 a ^{2/}	58.1 b
2	57.4 a	57.9 a	56.5 cd ^{2/}	57.4 nb	56.9 de	57.3 cd
3	57.4 a	57.4 a	56.2 d	58.0 a	55.2 g	57.6 bc
4	57.0 a	58.1 a	55.3 e	58.2 a	55.2 g	56.5 e
5	57.2 a	58.7 a	55.2 e	57.0 bc	54.4 h	56.3 ef
6	56.7 a	57.8 a	54.2 f	57.9 a	54.3 h	55.7 fg

^{1/} See Table 2 for threshing dates.

^{2/} Within a year, test weight data followed by the same letter do not differ at the 5% confidence level.

ORIGINAL PAGE IS
OF POOR QUALITY

ORIGINAL PAGE IS
OF POOR QUALITY

Table 7. Effect of stubble height on grain test weight and grain nitrogen (N) concentration, 1980.

<u>Measurement</u>	<u>Stubble height, inches</u>			<u>Standing</u>
	<u>4</u>	<u>9</u>	<u>14</u>	
Test weight (lbs/bu) ^{1/}	55.0 bc	55.5 b	54.8 c	57.7 a
Grain N, field (%) ^{2/} ^{5/}	2.25 b	2.32 b	2.34 b	2.52 a
Grain N, good (%) ^{3/}	2.26 ab	2.16 b	2.35 a	2.32 a
Grain N, damaged (%) ^{4/}	2.52 b	2.55 b	2.40 c	2.91 a

- ^{1/} Within any row, measurement data followed by the same number do not differ at the 5% confidence level.
- ^{2/} Grain N concentration of kernels as threshed in the field.
- ^{3/} Grain N concentration of kernels not discolored or sprouted.
- ^{4/} Grain N concentration of sprouted and discolored kernels.
- ^{5/} Grain N times 5.7 equals percent protein.

ORIGINAL PAGE IS
OF POOR QUALITY

Table 8. Grain nitrogen concentration comparisons between swathed and straight combine methods of cutting hard red spring wheat.

Threshing date	Year					
	1979		1980		1981	
	Method		Method		Method	
	<u>Swath</u>	<u>Straight</u>	<u>Swath</u>	<u>Straight</u>	<u>Swath</u>	<u>Straight</u>
	% N		% N		% N	
1	2.82 bc ^{2/}	2.79 bc	-	-	3.37 a ^{2/}	3.47 a
2	2.93 ab	2.95 ab	2.30	2.48	3.37 a	3.27 a
3	2.85 bc	2.84 bc	2.36	2.54	3.21 a	3.38 a
4	2.72 c	3.09 a	2.25	2.49	3.40 a	3.26 a
5	2.77 bc	2.86 bc	2.40	2.70	3.36 a	3.27 a
6	2.81 bc	2.96 ab	<u>2.31</u>	<u>2.42</u>	3.28 a	3.49 a
Avg.			2.32 b ^{3/}	2.52 a ^{3/}		

^{1/} See Table 2 for threshing dates.

^{2/} Within a year, nitrogen concentration data followed by the same letter do not differ at the 5% confidence level.

^{3/} Nitrogen concentration was lower in the windrows than in the standing grain.

ORIGINAL PAGE IS
OF POOR QUALITY

Table 9. Effect of stubble height and threshing date on percent damage to wheat grain, 1980.

Threshing ^{1/} date	Stubble height, inches			Average
	4	9 percent	14	
2 ^{2/}	9 f ^{3/}	24 cde ^{3/}	7 f ^{3/}	13 d ^{4/}
3	27 cd	25 cde	16 def	23 c
4	35 bc	31 c	12 ef	26 c
5	48 ab	28 cd	23 cde	33 b
6	50 a	45 ab	51 a	49 a
Average	34 a ^{5/}	30 a	22 b	

^{1/} See Table 2 for threshing dates.

^{2/} Dates: 8/25, 8/28, 9/5, 9/11, and 9/29

^{3/} Among these three columns data followed by the same letter do not differ at the 5% confidence level.

^{4/} Within this column, data followed by the same letter do not differ at the 5% confidence level.

^{5/} Within this row, data followed by the same letter do not differ at the 5% confidence level.

Table 10. Percent of damaged kernels in threshed grain in 1980 and 1981 as affected by date and cutting method.

Threshing date ^{1/}	1980		1981	
	Method		Method	
	Swath percent	Standing	Swath percent	Standing
1	-	-	6 d ^{3/}	9 bod ^{3/}
2	24	12	11 bc	10 bod
3	25	11	11 bc	8 cd
4	31	13	14 ab	7 cd
5	28	25	19 a	8 cd
6	<u>45</u>	<u>23</u>	<u>19</u> a	<u>9</u> bod
Average	30 a ^{2/}	17 b	13 a ^{2/}	8 b

^{1/} See Table 2 for threshing dates.

^{2/} Within a year, data of averages followed by the same letter do not differ at the 5% confidence level.

^{3/} Among these two columns, exclusive of the average values, data followed by the same letter do not differ at the 5% confidence level.

Table 11. Water concentration in the grain and straw at threshing as affected by cutting method, 1981.

Threshing date	Grain			Straw		
	Cutting method			Cutting method		
	Swath (feet)			Swath (feet)		
	10.5	15 % water	Standing	10.5	15 % water	Standing
1	8 ^{2/}	8	10	10 ^{2/}	9	37
2	9	11	13	8	8	17
3	10	12	11	-	-	-
4	9	10	11	6	4	31
5	13	12	14	11	13	23
6	<u>13</u>	<u>16</u>	<u>14</u>	<u>9</u>	<u>10</u>	<u>18</u>
	11 b ^{3/}	11 b	12 a	9 b ^{4/}	9 b	25 a

^{1/} See Table 2 for threshing dates.

^{2/} Statistically the water concentration did not differ in the grain or straw within a given harvest.

^{3/} Within this row, data followed by the same letter do not differ at the 5% confidence level.

^{4/} Within this row, data followed by the same letter do not differ at the 5% confidence level.