

General Disclaimer

One or more of the Following Statements may affect this Document

- This document has been reproduced from the best copy furnished by the organizational source. It is being released in the interest of making available as much information as possible.
- This document may contain data, which exceeds the sheet parameters. It was furnished in this condition by the organizational source and is the best copy available.
- This document may contain tone-on-tone or color graphs, charts and/or pictures, which have been reproduced in black and white.
- This document is paginated as submitted by the original source.
- Portions of this document are not fully legible due to the historical nature of some of the material. However, it is the best reproduction available from the original submission.

AgRISTARS

EW-U3-04397
JSC-18601

"Made available under NASA sponsorship
in the interest of early and wide dis-
semination of Earth Resources Survey
Program information and without liability
for any use made thereof."

Early Warning and Crop Condition Assessment

A Joint Program for
Agriculture and
Resources Inventory
Surveys Through
Aerospace
Remote Sensing

FEBRUARY 1983

A METEOROLOGICALLY-DRIVEN YIELD REDUCTION MODEL
FOR SPRING AND WINTER WHEAT

E83-10250
CR-170254

F. W. RAVET, W. J. CREMINS, T. W. TAYLOR,
P. ASHBURN, D. SMIKA AND A. AARONSON

(E83-10250) A METEOROLOGICALLY-DRIVEN YIELD
REDUCTION MODEL FOR SPRING AND WINTER WHEAT
(Department of Agriculture) 19 p
HC A02/MF A01

N83-23655



CSCI 02C

Unclas
00250

G3/43



U.S. DEPARTMENT OF AGRICULTURE
1050 BAY AREA BOULEVARD
HOUSTON, TEXAS 77058



Lyndon B. Johnson Space Center
Houston, Texas 77058

1. Report No. EW-U3-04397 JSC-18601		2. Government Accession No.		3. Recipient's Catalog No.	
4. Title and Subtitle A Meteorologically-Driven Yield Reduction Model for Spring and Winter Wheat				5. Report Date February, 1983	
				6. Performing Organization Code	
7. Author(s) F. W. Ravet, W. J. Cremins, T. W. Taylor, P. Ashburn, D. D. ... A. Aaronson				8. Performing Organization Report No.	
				10. Work Unit No.	
9. Performing Organization Name and Address U.S. Department of Agriculture EW/CCA and FAS/FCCAD 1050 Bay Area Blvd. Houston, Texas 77058				11. Contract or Grant No.	
				13. Type of Report and Period Covered Technical Memorandum	
12. Sponsoring Agency Name and Address Early Warning/Crop Condition Assessment, AgRISTARS 1050 Bay Area Blvd. Houston, Texas 77058				14. Sponsoring Agency Code	
15. Supplementary Notes A yield-reduction model for spring and winter wheat has been developed for large-area crop					
16. Abstract A yield reduction model for spring and winter wheat has been developed for large-area crop condition assessment. Reductions are expressed in percentage from a base yield and are calculated on a daily basis. The algorithm contains two integral components: a two-layer soil water budget model and a crop calendar routine. Yield reductions associated with hot, dry winds (Sukhovey) and soil moisture stress are determined. Input variables include evapotranspiration, maximum temperature and precipitation; subsequently crop-stage, available water holding percentage and stress duration are evaluated. No specific base yield is required and may be selected by the user; however, it may be generally characterized as the maximum likely to be produced commercially at a location. <p style="text-align: center;">ORIGINAL PAGE IS OF POOR QUALITY</p>					
17. Key Words (Suggested by Author(s)) Wheat Yield Reduction model Sukhovey				18. Distribution Statement	
19. Security Classif. (of this report) Unclassified		20. Security Classif. (of this page) Unclassified		21. No. of Pages 17	22. Price*

ORIGINAL PAGE IS
OF POOR QUALITY


EW-U3-04397
JSC-18601

A METEOROLOGICALLY-DRIVEN YIELD REDUCTION MODEL
FOR SPRING AND WINTER WHEAT

PRINCIPAL INVESTIGATORS

F.W. Ravet, W. J. Cremins, T. W. Taylor,
P. Ashburn, D. Smika, A. Aaronson

APPROVED BY



Glennis O. Boatwright, Manager
Early Warning/Crop Condition Assessment Project
AgRISTARS Program

Houston, Texas
February, 1983

A METEOROLOGICALLY DRIVEN YIELD REDUCTION
MODEL FOR SPRING AND WINTER WHEAT

F. W. Ravet, W. J. Cremins, T. W. Taylor, P. Ashburn, D. Smika, A. Aaronson

February 1983

WHEAT YIELD REDUCTION MODEL

PART 1.0 INTRODUCTION

1.1 PURPOSE

This paper documents a yield reduction model for winter and spring wheat. Reductions are expressed in percentage from a base yield and are calculated on a daily basis. A synopsis of the model logic and components is given.

1.2 SITUATION AND BACKGROUND

U.S. Department of Agriculture (USDA) policy is to provide American farmers and commodity analysts with timely information concerning world agricultural conditions. In 1978 USDA's Foreign Agricultural Service (FAS) created a Foreign Crop Condition Assessment Division (FCCAD) to aggressively pursue this policy. To further enhance this pursuit, U.S. government agencies involved in aerospace remote sensing coordinated their activities in the Agricultural and Resource Inventory Surveys Through Aerospace Remote Sensing (AgRISTARS).

An Early Warning/Crop Condition Assessment (EW/CCA) element of AgRISTARS seeks means to detect changes in production quantity and quality of commodities and renewable resources. The overall objective of the EW/CCA Project is to provide better capability for the USDA to identify environmental and agronomic events which significantly affect crop

condition and to determine extent and magnitude. Research conducted by EW/CCA augments and strengthens the capability of FCCAD. Improved crop condition information even based on subjective criteria is very useful in assessing crop loss and damage. As research provides better tools, subjective estimates will iteratively become more objective.

FCCAD operations call for assessments based on convergence of evidence from available data sources. Information sources include traditional reports from American embassies and consulates around the world and from information media. These are coupled to in-house use of agrometeorological crop condition indicator models and subjective analysis of remotely sensed satellite data.

The World Meteorological Organization (WMO) through its global telecommunications system provides timely exchange of meteorological data throughout the world. The U.S. Air Force collects meteorological data from an even larger network of stations and adapts the data to a gridded system. Environmental parameters, available daily, include in part maximum and minimum temperatures, type and quality of precipitation, solar radiation, snowfall and snow cover, wind direction and speed, vapor pressure and evapotranspiration.

EW/CCA and FCCAD developed crop stress indicator models and models that track crop phenology and soil moisture. This paper describes a yield reduction model which estimates daily stress impact on yield and particularly the impact of desiccating events as combinations of high temperature, low humidity, high wind speed and low soil moisture.

ORIGINAL PAPER IS OF POOR QUALITY

Quantity of water available at specific phases of plant growth and development greatly affects crop yield (Bauer and Young, 1969). Evaporation, as a measure of atmospheric demand, strongly influences crop condition. Primarily, it is a function of temperature, wind speed and humidity (Denmead and Shaw, 1962). Temperature is a major factor due to its effect on vapor pressure - an increase in temperature increases the evaporation rate and decreases the time a given quantity of water can effectively hydrate a plant.

Drought, a prolonged extension of desiccating conditions or soil moisture deficits that disrupt the water balance of plants, affects thousands of square kilometers annually (Ventskevich, 1961). Disruption of the water balance of plants may be induced either by a moisture deficit in the soil or by exceedingly rapid evaporation from plant surfaces. Hot, dry winds (called Sukhovey in the Soviet Union) may cause extreme yield reductions and/or plant death even though soil moisture conditions are optimal (Vitkevich, 1960). Sukhoveys have been roughly characterized as: 1) temperatures greater than 25°C, 2) relative humidity less than 25 percent and 3) windspeeds greater than 3-5 m/sec (CIA, 1974).

The operation of the yield reduction model does not require a specific base yield; rather, it measures yield loss. Therefore, any base yield can be selected by the user for final assessment. The base yield may however be generally characterized as the maximum likely to be produced commercially at a location.

PART 2.0 WHEAT YIELD REDUCTION MODEL

2.1 MODEL COMPONENTS

The yield reduction model contains two integral components; a crop calendar model and a soil moisture model. Both components require daily minimum and maximum temperatures. The crop calendar model uses the Robertson Biometeorological Time Scale (BMTS) (Figure 1) and the soil moisture model is the two-layer or crop moisture index model.

The crop calendar model requires actual or estimated planting dates. It identifies when 50 percent of the crop reaches a particular growth phase. Future refinements may integrate other crop phase increments into the model.

The two-layer soil moisture model requires long term monthly historical mean temperatures, daily rainfall and daily mean temperature and an estimate of the soil's available water-holding capacity.

There are two distinct modules in the operation of the yield reduction model; the ETP/Sukhovey module and the stress module. Sukhovey loss and damage can occur in a matter of minutes or hours while stress from diminishing soil moisture evolves over a longer period. Both can be very destructive.

2.2 PHENOLOGICAL GROWTH STAGE

The Robertson Biometeorological Time Scale measures phenological phase (Figure 1) and performs in the model as defined below and further described in Figure 1.


ROBERTSON'S BMTS	PHENOLOGICAL PHASE
0.0	PLANTING
1.0	EMERGENCE
1.5	TILLERING
2.0	JOINTING
2.5	FLAG LEAF
3.0	HEADING
3.5	MILK
4.0	DOUGH
5.0	RIPE

Attainment of maximum wheat yields is achieved through optimal environmental conditions during each phenological phase. However, if critical environmental factors such as soil, water and temperature are limiting, they reduce yield potential.

2.3 TWO-LAYER SOIL MOISTURE MODEL

The two-layer soil moisture model used by FCCAD and EW/CCA is similar to the Palmer two-layer model (Palmer, 1965). Atmospheric demand and soil water availability determine the amount of water withdrawn from the soil by both direct evaporation from the soil surface and transpiration by plants.

FIGURE 1. BMTS AND PHENOLOGICAL PHASES



ROBERTSON, G. W. - 1968

ORIGINAL SOURCE
OF POOR QUALITY

OF POOR QUALITY

Both soil moisture models assume the first inch of available water is held in the surface layer. The actual thickness of the surface layer is variable and depends on soil type, rooting depth and soil permeability.

The Palmer model assumes moisture is removed from the surface layer at a rate equal to potential evapotranspiration, and that moisture is removed from the subsurface layer at a fraction of the potential rate. The model assumes that moisture cannot be removed from the lower layer until the surface layer is completely dry. These assumptions are restrictive and do not adequately represent the soil water budget process.

Stress indicator models require a more accurate representation of the soil water budget, particularly in the surface layer. EW/CCA and FCCAD modified the two-layer model to allow a more gradual and realistic depletion of the surface layer and also allow moisture to be depleted from the lower layer before the surface layer is completely dry.

They also developed a moisture extraction function to allow depletion from the surface at the potential rate of less than or equal to 75 percent of surface capacity. Below 75 percent, the model extracts moisture from the surface at a reduced rate with the lower layer making up the remaining requirement. It extracts moisture from the lower layer at a fraction of potential and calculates this fraction as a ratio of actual water held to that level held at field capacity.

OF POOR QUALITY

Precipitation enters the model by first completely filling the surface layer and then the lower layer. When both layers reach capacity, excess precipitation becomes runoff and/or deep percolation and is lost from the model.

2.4 SUKHOVEY/ETP FUNCTION

This component of the model assigns empirical loss/damage magnitudes based on ETP and maximum temperature. The higher the temperature and ETP, the greater the yield loss (Table 1).

Adjustments are then made for soil moisture (as % of available water-holding capacity) (Table 2), crop phase (both for vulnerability and ETP) and duration of Sukhovey conditions (Table 3). An empirical scale assigns yield reduction by percent available soil water. Further adjustments are applied for crop phase (CSE). As the crop advances through growth and development, its water needs change. The crop becomes more or less vulnerable to environmental impact on yield. The calendar factor increases from small values during early phases of growth and development to a maximum value during early ripening (milk stage); then it decreases as the grain matures. The model calculates yield reduction as:

TABLE 1 - ETP/MAX TEMPERATURE YIELD REDUCTION

OF FISH QUALITY

MATRIX

		ETP													
		8	9	10	11	12	13	14	15	16	17	18	19	20	
M A X I M U M	26	0	0	0	0	0	0	0	0	0	1	2	3	4	
	27	0	0	0	0	0	0	0	1	2	3	4	5	6	
	28	0	0	0	0	0	1	2	3	4	5	6	7	8	
	29	0	0	0	1	2	3	4	5	6	7	8	9	10	
	30	0	1	2	3	4	5	6	7	8	9	10	11	12	
	31	2	3	4	5	6	7	8	9	10	11	12	13	14	
	32	4	5	6	7	8	9	10	11	12	13	14	15	16	
	33	6	7	8	9	10	11	12	13	14	15	16	17	18	
	34	8	9	10	11	12	13	14	15	16	17	18	19	20	
	35	10	11	12	13	14	15	16	17	18	19	20	21	22	
	36	12	13	14	15	16	17	18	19	20	21	22	23	24	
	37	14	15	16	17	18	19	20	21	22	23	24	25	26	
	38	16	17	18	19	20	21	22	23	24	25	26	27	28	
	39	18	19	20	21	22	23	24	25	26	27	28	29	30	
	40	20	21	22	23	24	25	26	27	28	29	30	31	32	
41	22	23	24	25	26	27	28	29	30	31	32	33	34		
42	24	25	26	27	28	29	30	31	32	33	34	35	36		
43	26	27	28	29	30	31	32	33	34	35	36	37	38		
44	28	29	30	31	32	33	34	35	36	37	38	39	40		
45	30	31	32	33	34	35	36	37	38	39	40	41	42		

OF POOR QUALITY

TABLE 2 - SOIL WATER AVAILABILITY FACTOR

% AWHC	YIELD REDUCTION FACTOR
5	1.25
10	1.25
15	1.25
20	1.20
25	1.15
30	1.10
35	1.05
40	1.00
45	0.99
50	0.98
55	0.97
60	0.96
65	0.95
70	0.94
75	0.93
80	0.92
85	0.91
90	0.90
95	0.89
100	0.88

SMIKA, D. - 1982

TABLE 3 - CONTINUOUS DAY ADJUSTMENT

DAY	ADJUSTMENT
1	1.0
2	0.5
3	0.25
4	0.125
5	0.0625
6	0.03125

ORIGINAL COPY
OF PROGRAM FILE

$$YR = (\text{Table 1 value}) * (\text{Table 2 value}) * (\text{Table 3 value}) * \text{CSE}$$

(Crop stage equation)

Prior to BMT 4.0 (milk) the model calculates crop phase effect as:

$$\text{CSE}_1 = 8.208 - 9.452X + 3.405 X^2 - 0.3666X^3$$

where X = BMT

This second equation determines the milk to ripe crop stage effect:

$$\text{CSE}_2 = 16.94 - 6.829X + 0.948X^2 - 0.052X^3$$

The model introduces a duration factor during each crop phase. The first occurrence of a harmful event receives full impact value. Values decrease one-half for the second occurrence during a crop phase, one-fourth for the third and so on (Table 3). These need not be consecutive. The occurrence counter resets to full impact at crop phases 2.5, 3.0, 3.5, 4.0 and 4.5.

An example of the scheme:

If max temperature = 37°C

If ETP = 15

then yield reduction factor (matrix) = 21

If % AWHC = 55

then AWHC factor = $.97 \times 21 = 20.37$

If crop phase is flowering the impact factor = 1.3 (from CSE₁)

then $1.3 \times 20.37 = 26.48$

If it is second day of event in the same crop phase; i.e., .5

then $.5 \times 26.48 = 13.24$


The event would cause a 13.24 point (%) yield reduction from a base yield figure.

2.5 SOIL MOISTURE/DEMAND FUNCTION


If ETP and temperature values do not fall in the ETP/Sukhovey module, the algorithm then defaults to the stress module. The stress module assesses yield reduction for small grains using ETP, available soil moisture and crop phase. The model assumes ETP to be the sum of the demands that are being placed on the plant by the environment and uses it to calculate a stress index. The stress index is a regression equation generated from the Moisture:ETP:Stress nomograph (Figure 2).

$$\text{Index} = .9998 + (.108 \text{ ETP}) + (.0145 \text{ AWC}) - (.3536 (\text{AWC})^5) - (.0001)$$


PERCENT
CAPACITY



ETP
m/m
DAY



SOIL MOISTURE
STRESS
INDEX



NOMOGRAPH
MOISTURE:ETP:STRESS

FIGURE 2

where

Index = the crop stress index (between 0 and 1)

ETP = the evapotranspiration potential

AWC = the percent of soil water available to the plant

A daily Yield Reduction Index is then adjusted for crop phase (See Figure 3). This is accomplished by multiplying the Index and the maximum daily reduction.

An example:

If maximum temperature = 28°C

ETP = 10

% AWC = 10

then Index = 1.10

and if the crop phase is at heading, then the stage factor is 3.4.

1.10 X 3.4 = 3.74 yield reduction.


The model calculates ETP using the albedo is specified as a function of crop phase.

ORIGINAL PLAN
OF FOUR QUARTERS

$$ETP = (A-1) ETP_1 + ETP_2$$

Phase	(A-1)
0 - .99	.9 - .14 phase
1 - 2.99	.76
3 - 5.00	.76 + .14 (phase - 3.0)

FIGURE 3 - DAILY YIELD REDUCTION



CROP PHASE

STEGMAN, E. C., AND H. BAUER. 1970. Canad. J. Ag. Eng. 12(2):92.

APPENDIX I

OF THE ...

SOIL MOISTURE EQUATION

Top Layer = Contains 1 inch of plant available water.

Lower Layer = Normally contains between 5 and 10 inches of available water.

$$L_s = S'_s - (ETP-P) D_f$$

$$L_u = (ETP-P-L_s) \frac{S'_u}{AWC} : L_u - S'_u$$

D_f = Surface moisture extraction function.

$$D_f = 1 \text{ if } P \leq ETP$$

$$D_f = (S'^s \div .75) : .1 - D_f - 1.$$

$$D_f = .1 \text{ if } D_f < .1 \text{ and } D_f = 1. : D_f = 1.$$

R = Excess P after both layers are filled.

ETP = ETP'(d) [Thornwaite, 1948]

If T less than 0°
ETP' = 0

If 0°C. T 26°C
ETP' = 1.6 $(\frac{10T}{I})^a$

If T 26°C
ETP' = Sin (T - 9.5) -.76

$$a = 6.75 \times 10^{-7} I^3 - 7.71 \times 10^{-5} I^2 + .01792I + .49239$$

$$I = \frac{12 \cdot 1.514}{(T/5)}$$

$$i=1$$

$$d = -0.767 \tan (.410117 \cos (.0172264(JDAY-172)))$$

APPENDIX II

DEFINITION OF TERMS

- L_s = Moisture loss from surface
- S'_s = Available water in surface layer at start
- P = Daily precipitation
- L_u = Loss from lower layer
- S'_u = Available moisture stored in lower layer
- AWC = Combined available water capacity; i.e., $\text{MAX}(S'_s + S'_u)$
- R = Runoff
- D_f = Surface moisture extraction function
- ETP = Evapotranspiration Potential - "The amount of water transpired in unit time by a short, green crop completely shading the ground, of uniform height and never short of water."
- d = Day length adjustment for ETP
- T = Average daily temp degree C
- I = Annual heat index
- JDAY = Julian date
- a = Coefficient
- CSE = Crop stage equation

LITERATURE CITED

- Bauer, A. 1972. Effect of Water Supply and Seasonal Distribution on Spring Wheat Yields. North Dakota State University Agricultural Experiment Station Bulletin 490. Fargo, ND.
- Bauer, A. and R. A. Young. 1969. Influence of Management and Environmental Factors on Extent of Soil Water Depletion by Spring Wheat. North Dakota Research Report 23.
- CIA. 1974. USSR Agricultural Atlas. Central Intelligence Agency, Washington, D.C.
- Denmead, O. T. and R. H. Shaw. 1962. Availability of Soil Water to Plants as Affected by Soil Moisture Content and Meteorological Conditions. Agron. J. 54:385.
- Palmer, W. C. 1965. Meteorological Drought. U.S. Weather Bureau Research Paper 45.
- Palmer, W. C. 1968. Keeping Track of Crop Moisture Conditions Nationwide: The New Crop Moisture Index. Weatherwise, pp. 156-161.
- Ravet, F. W. and J. R. Hickman. 1979. A Meteorologically Driven Wheat Stress Indicator Model. USDA/FAS Technical Memorandum #8, Houston, Texas.
- Robertson, G. W. 1968. A Biometeorological Time Scale for a Cereal Crop Involving Day and Night Temperatures and Photoperiod. Intern. J. Biometer. 12:191-233.
- Stegman, E. C., and A. Bauer. 1970. Computer Simulation Programming: An Aid to the Selection of Center Pivot Sprinkler Systems. Canad. J. Agric. Eng. 12(2):92.

Thorntwaite, C. W. 1948. An Approach Toward a Rational Classification of Climate. Geograph. Rev. 38:55-94.

Ventskevich, G. Z. Agrometeorology. Translated by National Science Foundation. Washington, D.C.

Vitkevich, V. I. 1960. Agricultural Meteorology. Translated by National Science Foundation. Washington, D.C.