

General Disclaimer

One or more of the Following Statements may affect this Document

- This document has been reproduced from the best copy furnished by the organizational source. It is being released in the interest of making available as much information as possible.
- This document may contain data, which exceeds the sheet parameters. It was furnished in this condition by the organizational source and is the best copy available.
- This document may contain tone-on-tone or color graphs, charts and/or pictures, which have been reproduced in black and white.
- This document is paginated as submitted by the original source.
- Portions of this document are not fully legible due to the historical nature of some of the material. However, it is the best reproduction available from the original submission.

We have examined several sets of coincident orbits where the ground track is nearly identical. In these cases the crust and upper mantle magnetic anomalies should be the same, but different ring current contributions would be expected because the orbits occur at different times, several weeks to months apart. Figures 1 and 2 show the results for two different sets. The polynomial fitting is based only upon the data between 50°N and 50°S latitudes. The best-fitting computed polynomial trends are then subtracted from the entire half-orbit to obtain the anomaly estimates. The higher the order of polynomial fit, generally the smaller are the magnitudes of the estimated anomalies. Our initial studies suggest that a third order polynomial provides the best anomaly estimate. The second order polynomial fit provides good consistency in the region of fitting, between 50 degrees north and south latitudes, however, the third degree provides a slightly better degree of consistency both within that same region as well as farther north and south beyond those bounds. Note how well the residuals from the third order polynomial agree with each other in both figures even though the original curves show considerable departures from each other presumably due to time-varying ring current effects. Thus a third order polynomial is the lowest polynomial order that appears to provide the best consistency of residual anomalies between coincident orbits. Because some half-revs yield residual crustal and upper mantle anomalies discordant with data from other nearby orbits, we, like Langel et al., 1982, delete values more than two standard deviations from the mean when interpolating data about a point.

The estimates of crustal and upper mantle magnetic anomalies above 50°N and below 50°S show considerable variation between coincident orbits, and among the residuals using different orders of polynomial fitting. Thus, for the immediate future we plan to concentrate our efforts on the region between

those two latitudes. However, we are retaining those values in our digital data library because with further examination, selection criteria for identifying valid crustal anomalies may be developed.

ORIGINAL PAGE IS
OF POOR QUALITY





Figure 1 Comparison of data from three coincident MAGSAT orbits. In the upper left hand corner are shown the MAGSAT residual anomaly profiles after removal of a core field represented by spherical harmonic coefficients through degree and order 13 (Langel et al., 1981) for half-orbits 1076, 1463, and 1851. The remaining panels show the residual anomalies remaining after subtraction from the aforementioned residuals of a polynomial trend of the degree indicated. The polynomial trends were computed only from data between 50 N and 50 S, although the continuation of those trends to higher latitudes enable residuals to be calculated over the range



ORIGINAL PAGE IS
OF POOR QUALITY

Figure 2 Comparison of data from two coincident MAGSAT orbits. The residuals after removal of core field effects for half-orbits 1054 and 2265 exhibit considerable departures from each other, particularly in the mid-latitudes. These departures are presumably owing to different ring current effects at the different times for the orbit passes. See caption for Fig. 1 for explanation of the profiles.

REFERENCES CITED

- Bowin, C., 1980, Why the Earth's Greatest Geoid Anomaly Is So Negative, *EOS*, 61, 17, 209.
- Bowin, C. 1981, Mass and topographic anomalies at the core-mantle boundary region from gravity and geoid observations, Abstracts of the 21st General Assembly IASPEI, London/Canada (July 21-30), p. B2.2.
- Bowin, C., in prep., Depth of principal mass anomalies contributing to the Earth's geoidal undulations and gravity anomalies.
- Bretherton, Francis P., Russ E. Davis and C.B. Fandry, 1976, A technique for objective analysis and design of oceanographic experiments applied to MODE-73, *Deep Sea Research*, 23, 559-582.
- Gandin, L.S., 1965, Objective analysis of meteorological fields, Israel Program for Scientific Translations.
- Langel, R., J. Berbert, T. Jennings, and R. J. Horner, 1981, MAGSAT data processing: An interim report for investigators, NASA TM 82160.
- Langel, R.A., J.D. Phillips and R.J. Horner, 1982, Initial scalar magnetic anomaly map from MAGSAT, *Geophys. Res. Lettrs.*, 9, 269-272.