brought to you by

NOTICE

THIS DOCUMENT HAS BEEN REPRODUCED FROM MICROFICHE. ALTHOUGH IT IS RECOGNIZED THAT CERTAIN PORTIONS ARE ILLEGIBLE, IT IS BEING RELEASED IN THE INTEREST OF MAKING AVAILABLE AS MUCH INFORMATION AS POSSIBLE

••Nade available under NASA sponsorship ••Nade available under NASA sponsorship ••Nade available under NASA sponsorship in the interest of early and wide disin the interest of early and without liability semination of Earth Resources Survey is semination of Earth Resources Survey in the interest of early and without liability semination of Earth Resources Survey is senior of the series of

E82-10240 CR-168756

INVESTIGATIONS OF MEDIUM WAVELENGTH MAGNETIC ANOMALIES IN THE EASTER! PACIFIC USING MAGSAT DATA

Christopher G. A. Harrison University of Miami Rosenstiel School of Marine & Atmospheric Science 4600 Rickenbacker Causeway Miami, Florida 33149

(E82-10240)INVESTIGATIONS OF MEDIUMN82-23013WAVELENGTH MAGNETIC ANOMALIES IN THE FESTERNPACIFIC USING MAGSAT Interim merort, Jul. -Sep. 1981 (Miami Univ.)2 p EC A02/MF A01UnclasCSCL 08G G3/4300240

November,1981 Interim Report for Period July-September 1981

RECEIVED

۰۰.

.

JAN 7, 1982SIS/902.6Prepared forM - 02.5GODDARD SPACE FLIGHT CENTER $T YF \in II$ Greenbelt, Maryland 20771

(MAGSAT Investigators)

QUAR'ERLY REPORT

JULY - SEPTEMBER 1981

The main body of research this quarter has been directed at the understanding of the long wavelength (~3000 km) anomalies appearing in the MAGSAT anomaly map. The existence of these anomalies resulted in controversy during the IAGA meeting during the summer. We cannot believe that these anomalies can be attributed to crustal sources but rather core sources that spherical harmonic models have failed to remove from the data. If true, this supposition has grave consequences for anybody attempting crustal magnitization modeling in large areas. Spectra of MAGSAT profiles of anomalous field tend to support the view that the spherical harmonic model is not performing as expected. As a result we are studying the details of spherical harmonic representations of the Earth's magnetic field.

During this quarter we also received the first of the INVESTIGATOR 'B' QUIET TIME tapes covering the time span NOVEMBER 1979 - FEDRUARY 1980. The data set seems to be error free. We have noted the change to an updated spherical harmonic model presumably corrected for the seasonal ionospheric currents which cr aminated the previous model.

Due to the attendance of the P.I. at the IAGA meeting and because of discussions concerning the wavelength of crustal and core anomalies, a paper is being prepared for the special issue of GRL on MAGSAT investigations, in which this question will be discussed.