

## N O T I C E

THIS DOCUMENT HAS BEEN REPRODUCED FROM  
MICROFICHE. ALTHOUGH IT IS RECOGNIZED THAT  
CERTAIN PORTIONS ARE ILLEGIBLE, IT IS BEING RELEASED  
IN THE INTEREST OF MAKING AVAILABLE AS MUCH  
INFORMATION AS POSSIBLE

Research on the Application of a Decoupling Algorithm  
for Structure Analysis

Final Report

NASA Grant NSG-1603

March 15, 1979 - March 14, 1980

(NASA-CR-163195) RESEARCH ON THE  
APPLICATION OF A DECOUPLING ALGORITHM FOR  
STRUCTURE ANALYSIS Final Report, 15 Mar.  
1979 - 14 Mar. 1980 (Houston Univ.) 78 p  
HC A05/MF A01

N80-24647

Unclas  
CSCL 20K G3/39 20995



Department of Electrical Engineering

Cullen College of Engineering

University of Houston

**Research on the Application of a Decoupling Algorithm  
for Structure Analysis**

**Final Report**

**NASA Grant NSG-1603**

**March 15, 1979 - March 14, 1980**

**Prepared by**

**Eugene D. Denman, Principal Investigator**

**University of Houston**

**Houston, Texas**

**for**

**NASA Langley Research Center**

**Hampton, Virginia 23665**

Table of Contents

1. Introduction . . . . .	1
2. Eigenprojectors of Matrices . . . . .	7
3. Lambda Matrices and Latent Projectors . . . . .	22
4. Projectors and the Sign Matrix . . . . .	33
5. Solvents of Matrix Polynomials . . . . .	41
6. Solution of a System of Differential Equations <sup>s</sup> . . . . .	49
7. Spectral Decomposition of Differential Equations . . . . .	61
8. Summary . . . . .	66
Appendix - Krylov Transformation . . . . .	68
References . . . . .	72

## Symbols

$A$	$m \times m$ matrix
$A(\lambda)$	$\lambda I - A$
$A(\lambda)$	lambda matrix or matrix polynomial
$J$	Jordan form matrix
$J_i$	pseudo-Jordan block
$\tilde{J}_i$	$n \times n$ matrix of Jordan blocks
$P_{i0}$	primary eigenprojector or matrix residue of $[A(\lambda)]^{-1}$
$\hat{P}_{i0}$	primary latent projector or matrix residue of $[A(\lambda)]^{-1}$
$P_{ij}$	$j > 0$ secondary eigenprojector
$\hat{P}_{ij}$	$j > 0$ secondary latent projector
$\Lambda_i$	coefficients of lambda matrix or matrix polynomial
$q$	number of pseudo-Jordan blocks in $A$
$r_i$	multiplicity of repeated eigenvalue $\lambda_i$
$\rho_i$	number of generalized eigenvectors for a repeated eigenvalue $\lambda_i$
$S$	sign matrix
$F_{i0}$	$m \times m$ matrix with $\text{sign}\{\text{diag}[0 \ 0 \ \dots \ \text{abs}(\lambda_i) \ \dots \ 0]\}$
$F_{ij}$	$m \times m$ matrix of superdiagonal elements for $J_i$
$R_i$	$i$ th solvent
$E$	diagonal matrix with $\pm 1$ on diagonals
$Q$	right eigenvector matrix
$P^+$	positive projector equal to sum of $P_{i0}$ with $\text{Re}(\lambda_i) > 0$
$P^-$	negative projector equal to sum of $P_{i0}$ with $\text{Re}(\lambda_i) < 0$
$I$	identity matrix
$y_i$	right eigenvector for $\lambda_i$
$z_i$	left eigenvector for $\lambda_i$

$\hat{y}_i$	right latent vector for $\lambda_i$
$\hat{z}_i$	left latent vector for $\lambda_i$
T	transformation matrix
$\rho$	scalar parameter
adj	adjoint of matrix
Badj	block adjoint of block matrix
det	determinant of matrix
Tr	trace of matrix
$\cdot \rangle \langle \cdot$	outer product
$\langle \cdot \cdot \rangle$	inner product
s	Laplace variable
$\Theta(t,0)$	state transition matrix

## Abstract

The mathematical theory for decoupling  $m$ th-order matrix differential equations is presented. It is shown that the decoupling procedure can be developed from the algebraic theory of matrix polynomials. The report discusses the role of eigenprojectors and latent projectors in the decoupling process and develops the mathematical relationships between eigenvalues, eigenvectors, latent roots and latent vectors. It is shown that the eigenvectors of the companion form of a matrix contains the latent vectors as a subset. The spectral decomposition of a matrix and the application to differential equations is given.

## 1. Introduction

The purpose of the material in this report is to formulate the algebraic theory of systems and application to spectral decomposition and decoupling of differential equations. The relationship between eigenvalues, eigenvectors, latent roots and latent vectors of matrix polynomials will be given. Since most of the equations of motion of vibrating systems are cast in second-order form, the algebraic properties of second-order matrix polynomials have an important role in the determination of solutions of vibration problems. Although the mathematical development will be in general form, the analysis includes second-order matrix polynomials.

The concept of scalar residues is well understood in complex variable theory and the inversion of Laplace transforms. The theory and use of matrix residues is not widely used and the relationship to eigenvectors and latent vectors has received little attention. Matrix residues, eigenprojectors and latent projectors are useful in analyzing matrix polynomials and time domain solutions to differential equations. Several papers have been published in recent years on matrix polynomials, see Dennis, Traub and Weber, [1], as well as a short paper by Denman, [2]. Some material on matrix residues has been given by Zadeh and Desoer, [3], and on projectors, Cullen [4]. Lancaster's book, [5], is an excellent source on latent roots and latent vectors of matrix polynomials which he denotes as lambda matrices.

It will be shown that eigenvalues and eigenvectors of the matrix companion form and latent roots and latent vectors of a matrix polynomial are related. The matrix residues of the inverse of a matrix polynomial  $\Lambda(\lambda) = I\lambda^m + A_1\lambda^{m-1} + \dots + A_m$ , which will be called latent projectors, are submatrices of the matrix residues of the inverse of  $(\lambda I - A_c)$  where  $A_c$  is in



block companion form. The latter residues will be referred to as eigenprojectors. It will be shown that latent projectors and eigenprojectors are useful for solving simultaneous differential equations.

The theory of Laplace transforms will be useful in introducing the concepts that are to follow. The material on Laplace transforms in most textbooks is limited to scalar problems and functions which is unfortunate since modern engineering problems are likely to be formulated as matrix problems due to the complexities of the systems to be analyzed. The extension of Laplace transforms to matrix functions is a simple task provided that the development of Laplace theory is based on algebraic functions rather than scalar functions.

Let  $f(t)$  be a scalar function for which the one-sided Laplace transform is given by

$$(1.1) \quad L[f(t)] = F(s) = \int_0^{\infty} f(t)e^{-st} dt$$

with the usual assumption that the integral exists. The inverse transform is defined as

$$(1.2) \quad L^{-1}[F(s)] = f(t) = \frac{1}{2\pi j} \int_{c-j\infty}^{c+j\infty} F(s)e^{st} ds$$

where  $c$  is properly defined to enclose all singularities of the integrand.

If  $F(s)$  has the property that

$$(1.3) \quad \lim_{s \rightarrow \infty} |F(s)| = 0$$

the inverse transform of  $F(s)$  is

$$(1.4) \quad L^{-1}[F(s)] = f(t) = \sum_{i=1}^n \text{residues of } [F(s)e^{st}] \Big|_{s=s_i} = \sum_{i=1}^n P_{i0} e^{s_i t}$$

where  $F(s)$  is a ratio of two scalar polynomials.

The Laplace transform method is valid for vector and matrix functions provided that certain restrictions are satisfied. Let  $A(s)$  be the matrix polynomial

$$(1.5) \quad A(s) = Is^m + A_1 s^{m-1} + \dots + A_m$$

where all coefficients  $A_i$  are  $n \times n$ . The inverse of  $A(s)$  is in general form

$$(1.6) \quad [A(s)]^{-1} = \frac{\text{adj}[A(s)]}{\det[A(s)]} = \frac{B(s)}{d(s)}$$

with  $\text{adj}$  the adjoint and  $\det$  the determinant of  $A(s)$  respectively. The characteristic equation of  $A(s)$  is given by  $d(s) = \det A(s)$  and will have a maximum of  $mn$  roots, Lancaster [5] calls these latent roots. The inverse transform of  $F(s) = [A(s)]^{-1}$  when the roots are distinct are

$$(1.7) \quad L^{-1}[F(s)] = f(t) = L^{-1}[[A(s)]^{-1}] = \sum_{i=1}^{mn} \text{residues of } [F(s)e^{st}] \Big|_{s=s_i}$$

which can be expressed as the matrix analog to (1.4) with

$$(1.8) \quad f(t) = \sum_{i=1}^{mn} \hat{P}_{i0} \exp(s_i t)$$

where the matrices  $\hat{P}_{i0}$  are matrix residues or latent projectors. It is obvious that the latent projectors are coefficients of the partial fraction

expansion

$$(1.9) \quad [A(s)]^{-1} = \sum_{i=1}^{mn} \frac{P_{i0}}{s-s_i}$$

The usefulness of the above approach to Laplace transforms can be illustrated by considering  $n$  simultaneous differential equations of  $m$ -th order, i.e.

$$(1.10) \quad A_0 \frac{d^m x}{dt^m} + A_1 \frac{d^{m-1} x}{dt^{m-1}} + \dots + A_m x = 0 \quad x(0) = c$$

where  $x(t)$  is a  $n$ th-order vector. It follows that

$$(1.11) \quad A(s)X(s) = A_m c \quad \dot{x}(0) = \ddot{x}(t_0) = \dots x^{(m)}(0) = 0$$

or

$$(1.12) \quad X(s) = [A(s)]^{-1} A_m c.$$

The time domain solution to (1.10) is then given by

$$(1.13) \quad x(t) = \sum_{i=1}^{mn} \hat{P}_{i0} A_m c \exp(s_i t)$$

when the latent roots are distinct.

The matrix polynomial given in (1.5) can also arise from the canonical form or companion matrix. It is not difficult to show that the matrix  $A(\lambda)$  given by

$$(1.14) \quad A(\lambda) = \begin{bmatrix} \lambda I & -I & 0 & \dots & 0 & 0 \\ 0 & \lambda I & -I & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \dots & \vdots & \vdots \\ 0 & 0 & 0 & \dots & \lambda I & -I \\ A_m & A_{m-1} & A_{m-2} & \dots & A_2 & \lambda I + A_1 \end{bmatrix}$$

has the same characteristic equation as (1.5) when  $A_0 = I$ . The root of the characteristic equation obtained from  $\det[A(\lambda)] = 0$  will be the eigenvalues of  $A$  and are equal to the latent roots of  $A(\lambda)$  when  $A_0 = I$ . The eigenvectors of  $A$  must be related to the latent vectors of  $A(\lambda)$ ; that relationship will be given later.

If  $z(t)$  is defined as the vector

$$(1.15) \quad [z(t)]^T = \{[x(t)]^T [\dot{x}(t)]^T \dots [x^{(m)}(t)]^T\}$$

then  $z(t)$  satisfies the equation

$$(1.16) \quad \dot{z}(t) = A z(t)$$

with

$$(1.17) \quad Ix^{(m)}(t) + A_1 x^{(m-1)}(t) + \dots + A_m x(t) = 0.$$

The solution vector  $z(t)$  is given by

$$(1.18) \quad z(t) = \sum_{i=1}^{mn} P_{i0} C \exp(\lambda_i t)$$

where  $P_{i0}$  are the matrix residues of  $[A(\lambda)]^{-1}$  which will be called eigenprojectors. It is assumed that (1.18) is for distinct eigenvalues of  $A$ . Since the eigenvalues of  $A$  and the latent roots of  $A(\lambda)$  are the same and the eigenvectors of  $A$  and the latent vectors of  $A(\lambda)$  are related, the eigenprojectors  $P_{i0}$  and the latent projectors  $\hat{P}_{i0}$  must be related. The vector  $C$  in (1.18) is obtained from (1.12) and the definition of the canonical form for the system.

## 2. Eigenprojectors of Matrices

Let  $A$  be defined as a  $m \times m$  matrix with eigenvalues  $\lambda_i$ , right eigenvectors  $y_i$  and left eigenvectors  $z_i$ . Define  $Q$  as a  $m \times m$  matrix constructed from the eigenvectors  $y_i$  such that

$$(2.1) \quad Q = [y_1 \ y_2 \ y_3 \ \dots \ y_m]$$

where it is assumed that the columns of  $Q$  are linearly independent and spans the  $C^{m \times m}$  space. The matrix  $Q$  has the property that a similarity transformation on  $A$  with  $Q$  will reduce  $A$  to the Jordan form

$$(2.2) \quad J = Q^{-1}AQ$$

with  $J = \text{diag}[J_1, J_2, \dots, J_p]$  with  $J_i$  a Jordan block. The Jordan form will be diagonal if  $A$  has  $m$  distinct eigenvalues or if  $A$  has  $m$  linearly independent eigenvectors satisfying  $[\lambda_i I - A]y_i = 0$ . If  $A$  has repeated eigenvalues and is defective, the Jordan blocks leading to the defectiveness of  $A$  will have one or more plus ones on the superdiagonal on a Jordan block. It will be necessary to utilize the chain rule for generating the generalized eigenvectors for the defective Jordan block.

The mathematical analysis will be simplified if all eigenvalues with the same values are considered as a pseudo-Jordan block with the plus ones on the superdiagonal. Assume that  $J_i$  and  $J_{i+1}$  are as shown with  $J_{i+1}$  having the same eigenvalues as  $J_i$  but where  $J_{i+1}$  has the plus ones on the superdiagonal. Although the definition

$$(2.3) \quad \begin{bmatrix} J_i & 0 \\ 0 & J_{i+1} \end{bmatrix} = \begin{bmatrix} \lambda_i & 0 & 0 & 0 \\ 0 & \lambda_i & 1 & 0 \\ 0 & 0 & \lambda_i & 1 \\ 0 & 0 & 0 & \lambda_i \end{bmatrix}$$

is not conventional. The two blocks  $J_i$  and  $J_{i+1}$  will be considered as a pseudo-Jordan block.

Let  $A$  have  $q$  values of  $\lambda_i$  with  $q$  pseudo-Jordan blocks as defined. Assume that  $m-k$  of the eigenvalues are distinct and  $q-m+k$  are repeated. Each repeated eigenvalue will have multiplicity  $r_i$  and the number of generalized eigenvectors for the repeated eigenvalues will be  $\ell_i$ , the number of plus ones on the superdiagonal of  $J_i$ . It will be assumed that the ones are located in the last  $\ell_i$  rows of the repeated eigenvalue pseudo-Jordan block. The term pseudo-Jordan block will be dropped in the following discussion and the term Jordan block will be utilized with  $J$  denoting a pseudo-Jordan block.

In addition to the above assumptions, let  $F_{ij}$  denote a  $m \times m$  matrix. The first subscript denotes the eigenvalue to which the  $F_{ij}$  matrix belongs and the second subscript is an index which has a maximum value equal to the number of generalized eigenvectors required for the eigenvalue  $\lambda_i$ ; this will be  $\ell_i$ . If  $\ell_i$  is the number of generalized eigenvectors for  $\lambda_i$ , then  $j = 0, 1, 2, \dots, \ell_i$  with  $j = 0$  for zero generalized eigenvectors. If  $\lambda_i$  is a distinct eigenvalue, then  $F_{i0}$  will be defined as

$$(2.4) \quad F_{i0} = \text{diag}[0, 0, \dots, 0, 1, 0, \dots, 0]$$

with the one located in the same row and column as  $\lambda_i$  is in  $J$ . The matrix

$F_{i0}$  for a repeated eigenvalue  $\lambda_i$  with multiplicity  $r_i = 3$  will have the form

$$(2.5) \quad F_{i0} = \text{diag}[0,0,\dots,0,1,1,1,0\dots,0]$$

with the one located in the same rows and columns as  $\lambda_i$  is in  $J$ .

To complete the definition of  $F_{ij}$ , assume that  $\lambda_i$  has multiplicity  $r_i$  with  $r_i - \ell_i$  linearly independent eigenvectors and  $\ell_i$  generalized eigenvectors. The associated Jordan block will have  $\ell_i$  ones on the superdiagonal with the ones located in the last  $\ell_i$  rows of  $J_i$ . The matrix  $F_{i0}$  will be as given in (2.5) but the set of matrices  $F_{i1}, F_{i2}, \dots, F_{i\ell_i}$  will now exist with  $F_{i1}$  having only the ones of the superdiagonal of  $J_i$  located on the superdiagonal of  $F_{i1}$ . The next matrix in the sequence  $F_{ij}$  will be generated by moving the ones on the superdiagonal of  $F_{i1}$  up one diagonal position by moving to the next columns of  $F_{i1}$ . To illustrate the construction of  $F_{ij}$ , let  $J$  be defined as in (2.3), then  $F_{i0}$ ,  $F_{i1}$  and  $F_{i2}$  are

$$(2.6) \quad F_{i0} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \quad F_{i1} = \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix} \quad F_{i2} = \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

The eigenprojectors of  $A$ , [6], can now be defined using the established notation. Assume that  $A$  has  $m-k$  distinct eigenvalues and  $q-m+k$  repeated eigenvalues. The primary eigenprojectors will be defined as

$$(2.7) \quad P_{i0} = Q F_{i0} Q^{-1} \quad i = 1, 2, \dots, q$$

with the primary eigenprojectors having the properties



$$(2.8a) \quad \sum_{i=1}^q P_{i0} = I$$

$$(2.8b) \quad P_{i0} P_{i0} = P_{i0}$$

$$(2.8c) \quad P_{i0} P_{j0} = 0 \quad i \neq j$$

These properties follow directly from (2.6) since  $\sum_{i=1}^q F_{i0} = I$ ,  $F_{i0} F_{i0} = F_{i0}$  and  $F_{i0} F_{j0} = 0$ . The primary eigenprojectors are idempotent matrices, i.e.  $P_{i0}^\alpha = P_{i0}$  where  $\alpha$  is a positive integer.

The spectral decomposition for  $A$  which is not defective is given by

$$(2.9) \quad A = \sum_{i=1}^q \lambda_i P_{i0}$$

which follows directly from the definition of the primary eigenprojectors.

If  $A$  is defective, a set of secondary eigenprojectors will be defined as the projectors constructed from the eigenvectors and the sequence of  $F_{ij}$  matrices. Let the secondary eigenprojectors be defined by

$$(2.10) \quad P_{ij} = Q F_{ij} Q^{-1} \quad j = 1, 2, \dots, \ell_i$$

with  $i$  set by the associated eigenvalue which is repeated but defective. It follows from the definition of  $F_{ij}$  that

$$(2.11a) \quad P_{i0} P_{ij} = P_{ij} \quad j = 1, 2, \dots, \ell_i$$

$$(2.11b) \quad P_{ij} P_{ij} = P_{i,j+1} \quad j \neq 0$$

$$(2.11c) \quad P_{ij} P_{i,j+1} = 0 \quad j \neq 0$$

The secondary eigenprojectors  $P_{i1}$  are required for the spectral decomposition of the most general matrix  $A$ .

If  $A$  has general form and is defective then

$$(2.12) \quad A = Q J Q^{-1} = Q \left[ \Lambda + \sum_{i=1}^q F_{ir} \right] Q^{-1}$$

but this is equal to

$$(2.13) \quad A = Q \left\{ \sum_{i=1}^q [F_{i0} \lambda_i + F_{i1}] \right\} Q^{-1}$$

or finally

$$(2.14) \quad A = \sum_{i=1}^q [P_{i0} \lambda_i + P_{i1}]$$

Although the secondary eigenprojectors  $P_{ij}$  with  $j > 1$  are not necessary for the spectral decomposition, it will be shown later that the partial fraction expansion of  $[A(\lambda)]^{-1} = [\lambda I - A]^{-1}$  can be expressed in terms of the eigenprojectors.

The procedure given for computing the eigenprojectors for a matrix has been based on the assumption that the right eigenvector matrix,  $Q$ , is known completely. The inverse of  $Q$  will have row vectors that are the left eigenvectors of  $A$  thus  $P_{i0}$  depends on the right and left eigenvectors,  $y_i$  and  $z_i$  respectively. The right and left eigenvectors for distinct eigenvalues are determined from the equations

$$(2.15) \quad [\lambda_i I - A] y_i = 0$$

$$(2.16) \quad z_i [\lambda_i I - A] = 0$$

or equivalently  $[\lambda_i I - A^T] z_i^T = 0$  for the left eigenvectors. If  $A$  is defective for an eigenvalue  $\lambda_i$ , the chain rules, [7]

$$(2.17) \quad [\lambda_i I - A] y_i^{k+1} = -y_i^k \quad k = 1, 2, \dots, \ell_i$$

$$(2.18) \quad [\lambda_i I - A^T] (z_i^{k+1})^T = -(z_i^k)^T \quad k = 1, 2, \dots, \ell_i$$

are used for the generalized eigenvectors where  $y_i^1$  and  $z_i^1$  are any one of the linearly independent eigenvectors for the repeated eigenvalues.

The primary eigenprojectors  $P_{i0}$  were defined earlier and are given by

$$(2.19) \quad P_{i0} = Q F_{i0} Q^{-1}$$

Let  $Q = Q_r$  be the matrix of right eigenvectors and  $Q_\ell$  be the matrix of left eigenvectors for distinct eigenvalues with

$$(2.20) \quad Q_r = [\bar{y}_1 \quad \bar{y}_2 \quad \dots \quad \bar{y}_{mn}] \quad Q_\ell = \begin{bmatrix} \bar{z}_1 \\ \bar{z}_2 \\ \vdots \\ \bar{z}_{mn} \end{bmatrix}$$

and let  $Q_\ell$  be scaled such that  $Q_\ell Q_r = I$ . It then follows for a distinct eigenvalue  $\lambda_i$  that

$$(2.21) \quad P_{i0} = Q_r F_{i0} Q_\ell = \bar{y}_i \langle \bar{z}_i = \bar{y}_i \bar{z}_i .$$

Since  $Q_r F_{i0} Q_l$  will be given by the outer product of  $\bar{y}_i$  and  $\bar{z}_i$ , the scaling of  $Q_l$  is equivalent to scaling  $\bar{y}_i$  or  $\bar{z}_i$  such that  $\langle \bar{z}_i, \bar{y}_i \rangle = \bar{z}_i \cdot \bar{y}_i = 1$ . Since any eigenvector can be multiplied by a constant then any set of arbitrary scaled eigenvectors can be used to compute  $P_{i0}$  provided that the arbitrary constants are removed by dividing by the scaling factor. If  $y_i$  and  $z_i$  are arbitrary eigenvectors the eigenprojectors for the distinct eigenvalues are given by

$$(2.22) \quad P_{i0} = \frac{y_i z_i}{z_i y_i}$$

The eigenprojectors for the repeated eigenvalues when  $A$  is not defective are determined from a simple extension of (2.21). Since the eigenvectors are linearly independent,  $P_{i0}$  is given by

$$(2.23) \quad P_{i0} = Q_r F_{i0} Q_l = \sum_{j=1}^{r_i} \frac{y_i^j z_i^j}{y_i^j z_i^j} = \sum_{j=1}^{r_i} \frac{y_i^j z_i^j}{z_i^j y_i^j}$$

where the superscript denotes the  $j$ th eigenvector belonging to the pseudo-Jordan block.

The eigenprojectors for the repeated eigenvalues when  $A$  is defective are computed in a similar manner to the repeated eigenvalues for the nondefective case. The primary eigenprojector for a defective Jordan block is given by (2.23). The secondary eigenprojectors can be computed from the eigenvectors by considering (2.10). Let  $\hat{F}_{ij}$  denote the subblock of  $F_{ij}$  for a Jordan block  $J_i$  where  $\hat{F}_{ij}$  is  $r_i \times r_i$ . Let  $\bar{Y}_i$  and  $\bar{Z}_i$  denote the rectangular matrices of right and left eigenvectors respectively of  $\lambda_i$ . Equation (2.10) can be rewritten as

$$(2.24) \quad P_{ij} = \bar{Y}_i \hat{F}_{ij} \bar{Z}_i = [y_1^{-1} y_1^{-2} \dots y_1^{-r_i}] \hat{F}_{ij} \begin{bmatrix} -1 \\ z_i \\ -2 \\ z_i \\ \vdots \\ -r_i \\ z_i \end{bmatrix}$$

where  $Q_r$  and  $Q_\ell$  have been properly scaled. Let  $f_{st}$  denote the elements of  $\hat{F}_{ij}$  with  $f_{st} = 1$  or  $0$  depending on the  $i$ th row and  $j$ th column of  $\hat{F}_{ij}$ . Equation (2.24) can then be written as

$$(2.25) \quad P_{ij} = \sum_{s=1}^{r_i} \sum_{t=1}^{r_i} \frac{y_1^{-s}}{z_i^s y_1^s} f_{ij} z_i^{-t} = \sum_{s=1}^{r_i} \sum_{t=1}^{r_i} \left[ \frac{y_1^s f_{st} z_i^t}{z_i^s y_1^s} \right]$$

Since  $\hat{F}_{ij}$  will be sparse, only a few terms of the summation are required.

An example will now be given to illustrate the computational procedure using the eigenprojectors. Let  $A$  be given by

$$A = \frac{1}{2} \begin{bmatrix} 5 & -1 & 0 & -2 \\ 0 & 4 & -1 & -1 \\ -2 & 0 & 5 & -1 \\ -1 & -1 & -2 & 6 \end{bmatrix}$$

with Jordan form

$$J = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 3 & 1 & 0 \\ 0 & 0 & 3 & 1 \\ 0 & 0 & 0 & 3 \end{bmatrix} \quad \begin{array}{l} \lambda_1 = 1 \\ \lambda_2 = 3 \\ r_2 = 3 \end{array} \quad \ell_2 = 2$$

The right eigenvector matrix  $Q_r$  is

$$Q_r = \begin{bmatrix} 1 & -1 & -1 & 1 \\ 1 & -1 & 1 & -1 \\ 1 & 1 & -1 & -1 \\ 1 & 1 & 1 & 1 \end{bmatrix} = [y_1 \ y_2^1 \ y_2^2 \ y_2^3]$$

and the left eigenvector matrix by

$$Q_\ell = \begin{bmatrix} 1 & 1 & 1 & 1 \\ -1 & -1 & 1 & 1 \\ -1 & 1 & -1 & 1 \\ 1 & -1 & -1 & 1 \end{bmatrix} = \begin{bmatrix} z_1 \\ z_1^1 \\ z_2^2 \\ z_2^3 \\ z_2^2 \end{bmatrix}$$

The eigenprojector  $P_{10}$  is found from  $y_1$  and  $z_1$  and is

$$P_{10} = \frac{y_1 z_1}{z_1 y_1} = \frac{1}{4} \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \end{bmatrix} \quad \hat{F}_{10} = 1$$

with the primary eigenprojector  $P_{20}$  for  $\lambda_2 = 3$  given by

$$P_{20} = \sum_{j=1}^3 \frac{y_2^j z_2^j}{z_2^j y_2^j} = \frac{1}{2} \frac{1}{z_2^1 y_2^1} + \frac{2}{z_2^2 y_2^2} + \frac{3}{z_2^3 y_2^3}$$

$$= \frac{1}{4} \begin{bmatrix} 3 & -1 & -1 & -1 \\ -1 & 3 & -1 & -1 \\ -1 & -1 & 3 & -1 \\ -1 & -1 & -1 & 3 \end{bmatrix} \quad \hat{F}_{20} = I_{3 \times 3}$$

As a check on the two eigenprojectors, note that  $P_{10} + P_{20} = I$ . The first secondary eigenprojector  $P_{21}$  is given by

$$P_{21} = \sum_{s=1}^2 \sum_{t=1}^2 \frac{y_2^s f_{1j} z_2^t}{z_2^s y_2^t} = \frac{y_2^1 z_2^2}{z_2^1 y_2^1} + \frac{y_2^2 z_2^3}{z_2^2 y_2^2}$$

$$= \frac{1}{4} \begin{bmatrix} 0 & 0 & 2 & -2 \\ 2 & -2 & 0 & 0 \\ -2 & 2 & 0 & 0 \\ 0 & 0 & -2 & 2 \end{bmatrix} \quad f_{12} = 1 \quad f_{23} = 1$$

The second secondary eigenprojector is determined from (2.25) with  $f_{13} = 1$  as the only nonzero element in  $\hat{F}_{22}$ . The eigenprojector  $P_{22}$  is then

$$P_{22} = \frac{y_2^1 z_2^3}{z_2^1 y_2^1} = \frac{1}{4} \begin{bmatrix} -1 & 1 & 1 & -1 \\ -1 & 1 & 1 & -1 \\ 1 & -1 & -1 & 1 \\ 1 & -1 & -1 & 1 \end{bmatrix}$$

A check will show that  $P_{20} P_{21} = P_{21}, P_{20} P_{22} = P_{22}$  and  $P_{21} P_{22} = 0$ .

The spectral decomposition of  $A$  is given by  $\lambda_1 P_{10} + \lambda_2 P_{20} + P_{21}$  which is

$$A = \frac{1}{4} \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \end{bmatrix} + \frac{3}{4} \begin{bmatrix} 3 & -1 & -1 & -1 \\ -1 & 3 & -1 & -1 \\ -1 & -1 & 3 & -1 \\ -1 & -1 & -1 & 3 \end{bmatrix} + \frac{1}{4} \begin{bmatrix} 0 & 0 & 2 & -2 \\ 2 & -2 & 0 & 0 \\ -2 & 2 & 0 & 0 \\ 0 & 0 & -2 & 2 \end{bmatrix}$$

$$= \frac{1}{4} \begin{bmatrix} 10 & -2 & 0 & -4 \\ 0 & 8 & -2 & -2 \\ -4 & 0 & 10 & -2 \\ -2 & -2 & -4 & 12 \end{bmatrix}$$

which agrees with  $A$  as given.

The eigenprojectors can also be computed from the inverse of  $A(\lambda)$  where  $A(\lambda) = \lambda I - A$ . The most general partial fraction expansion of  $[A(\lambda)]^{-1}$  is given by

$$(2.26) \quad [A(\lambda)]^{-1} = \sum_{i=1}^q \left\{ \frac{P_{i0}}{\lambda - \lambda_i} + \sum_{j=1}^{\ell_i} \frac{P_{ij}}{(\lambda - \lambda_i)^{j+1}} \right\}$$

The three cases, distinct eigenvalue, repeated eigenvalues with multiplicity  $r_i$  but not defective and the defective matrix must be discussed.

The three cases can be analyzed by considering the Jordan blocks for the three different eigenvalue cases. Rather than consider the mixed Jordan forms, consider the three individual Jordan forms. Assume first that

$J = \text{diag}[\lambda_1 \lambda_2 \lambda_3]$  thus

$$(2.27) \quad A = QJQ^{-1} = Q \begin{bmatrix} \lambda_1 & 0 & 0 \\ 0 & \lambda_2 & 0 \\ 0 & 0 & \lambda_3 \end{bmatrix} Q^{-1}$$

with  $[A(\lambda)]^{-1}$  given by

$$(2.28) \quad [A(\lambda)]^{-1} = [\lambda I - A]^{-1} = Q \begin{bmatrix} (\lambda - \lambda_1)^{-1} & 0 & 0 \\ 0 & (\lambda - \lambda_2)^{-1} & 0 \\ 0 & 0 & (\lambda - \lambda_3)^{-1} \end{bmatrix} Q^{-1}$$

It follows directly from  $(\lambda - \lambda_i) [A(\lambda)]^{-1}$  evaluated at  $\lambda_i$  that

$$(2.29) \quad (\lambda - \lambda_i) [\lambda I - A]^{-1} \Big|_{\lambda = \lambda_i} = Q F_{i0} Q^{-1} = P_{i0}$$

Consider the Jordan form with  $\lambda_1$  repeated 3 times but where  $A$  is not defective. The inverse of  $A(\lambda)$  is



$$(2.30) \quad [A(\lambda)]^{-1} = [\lambda I - A]^{-1} = Q \begin{bmatrix} (\lambda - \lambda_1)^{-1} & 0 & 0 \\ 0 & (\lambda - \lambda_1)^{-1} & 0 \\ 0 & 0 & (\lambda - \lambda_1)^{-1} \end{bmatrix} Q^{-1}$$

Since  $\lambda_1$  has multiplicity 3 then  $(\lambda - \lambda_1)^3 [A(\lambda)]^{-1}$  will be

$$(2.31) \quad (\lambda - \lambda_1)^3 [A(\lambda)]^{-1} = Q \begin{bmatrix} (\lambda - \lambda_1)^2 & 0 & 0 \\ 0 & (\lambda - \lambda_1)^2 & 0 \\ 0 & 0 & (\lambda - \lambda_1)^2 \end{bmatrix} Q^{-1}$$

It is obvious that (2.31) will be zero when evaluated at  $\lambda = \lambda_1$ . The first derivative of (2.31) with respect to  $\lambda$  will also be zero at  $\lambda = \lambda_1$  with the second derivative of (2.31) given by

$$(2.32) \quad \frac{d^2}{d\lambda^2} \{(\lambda - \lambda_1)^3 [A(\lambda)]^{-1}\} = Q \begin{bmatrix} 2 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 2 \end{bmatrix} Q^{-1} = 2QF_{10}Q^{-1}$$

The eigenprojector for the repeated eigenvalue matrix with multiplicity  $r_1$  but for  $A$  not defective is given by

$$(2.33) \quad P_{10} = QF_{10}Q^{-1} = \frac{1}{(r_1 - 1)!} \frac{d^{r_1 - 1}}{d\lambda^{r_1 - 1}} \{(\lambda - \lambda_1)^{r_1} [A(\lambda)]^{-1}\} \Big|_{\lambda = \lambda_1}$$

The defective matrix will be analyzed by considering the Jordan block with  $\lambda_1$  of multiplicity 3 and 2 generalized eigenvectors. Let  $[A(\lambda)]^{-1}$  be given by

$$(2.34) \quad [A(\lambda)]^{-1} = Q \begin{bmatrix} \lambda - \lambda_1 & -1 & 0 \\ 0 & \lambda - \lambda_1 & -1 \\ 0 & 0 & \lambda - \lambda_1 \end{bmatrix}^{-1} Q^{-1} = Q \begin{bmatrix} (\lambda - \lambda_1)^{-1} & (\lambda - \lambda_1)^{-2} & (\lambda - \lambda_1)^{-3} \\ 0 & (\lambda - \lambda_1)^{-1} & (\lambda - \lambda_1)^{-2} \\ 0 & 0 & (\lambda - \lambda_1)^{-1} \end{bmatrix} Q^{-1}$$

The sequence of evaluations of  $(\lambda - \lambda_1)^3 [A(\lambda)]^{-1}$  and the derivatives with respect at  $\lambda = \lambda_1$  will be

$$(2.35) \quad (\lambda - \lambda_1)^3 [A(\lambda)]^{-1} \Big|_{\lambda = \lambda_1} = Q \begin{bmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} Q^{-1} = QF_{12}Q^{-1} = P_{12}$$

$$(2.36) \quad \frac{d}{d\lambda} \{(\lambda - \lambda_1)^3 [A(\lambda)]^{-1}\} \Big|_{\lambda = \lambda_1} = Q \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix} Q^{-1} = QF_{11}Q^{-1} = P_{11}$$

$$(2.37) \quad \frac{1}{2} \frac{d^2}{d\lambda^2} \{(\lambda - \lambda_1)^3 [A(\lambda)]^{-1}\} \Big|_{\lambda = \lambda_1} = Q \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} Q^{-1} = QF_{10}Q^{-1} = P_{10}$$

The computational procedure for finding the eigenprojectors, usually referred to as matrix residues in the above, from the partial fraction expansion can now be summarized. If  $\lambda_i$  is distinct then

$$(2.38) \quad P_{i0} = (\lambda - \lambda_i) [A(\lambda)]^{-1} \Big|_{\lambda = \lambda_i}$$

with the primary eigenprojectors for the repeated eigenvalues  $\lambda_i$  of multiplicity  $r_i$  given by

$$(2.39) \quad P_{i0} = \frac{1}{(r_i - 1)!} \frac{d^{r_i - 1}}{d\lambda^{r_i - 1}} \{(\lambda - \lambda_i)^{r_i} [A(\lambda)]^{-1}\} \Big|_{\lambda = \lambda_i}$$

The secondary eigenprojectors are defined only for repeated eigenvalues with multiplicity  $r_i$  with  $A$  defective and requiring  $\ell_i$  generalized eigenvectors.

The secondary eigenprojectors are given by

$$(2.40) \quad P_{1, \ell_1 - j} = \frac{1}{j!} \frac{d^j}{d\lambda^j} [(\lambda - \lambda_1)^{\hat{k}_1} [A(\lambda)]^{-1}] \quad j = 0, 1, \dots, \ell_1$$

where  $[A(\lambda)]^{-1}$  has all common factors of  $\text{adj}[A(\lambda)]$  and  $\det[A(\lambda)]$  cancelled so that  $[A(\lambda)]^{-1}$  is a minimum polynomial with  $\hat{k}_1$  equal to the power of  $(\lambda - \lambda_1)^{\hat{k}_1}$  in the denominator.

The computation of the eigenprojectors by the residue method will be illustrated with the previous example. The inverse of  $[A(\lambda)]^{-1}$  is given by

$$[A(\lambda)]^{-1} = \frac{1}{(\lambda-1)(\lambda-3)^3} \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \lambda^3 + \begin{bmatrix} -7.5 & -0.5 & 0 & -1 \\ 0 & -8 & -0.5 & -0.5 \\ -1 & 0 & -7.5 & -0.5 \\ -0.5 & -0.5 & -1 & -7 \end{bmatrix} \lambda^2$$

$$+ \begin{bmatrix} 17.75 & 3.25 & 1.25 & 4.75 \\ 0.75 & 20.25 & 3.25 & 2.75 \\ 5.25 & 0.75 & 17.75 & 3.25 \\ 3.25 & 2.75 & 4.75 & 16.25 \end{bmatrix} \lambda + \begin{bmatrix} -13.25 & -4.75 & -3.25 & -5.75 \\ -2.75 & -15.25 & -4.75 & -4.25 \\ -6.25 & -2.75 & -13.25 & -4.75 \\ -4.75 & -4.25 & -5.75 & -12.25 \end{bmatrix}$$

The eigenprojector for  $\lambda = 1$  is

$$P_{10} = \frac{1}{4} \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \end{bmatrix} = (\lambda-1) [A(\lambda)]^{-1} \Big|_{\lambda=1}$$

The eigenvalue  $\lambda = 3$  will have a primary eigenprojector and two secondary eigenprojectors. The eigenprojectors  $P_{22}$ ,  $P_{21}$  and  $P_{20}$  are

$$P_{22} = (\lambda-3)^3 [A(\lambda)]^{-1} \Big|_{\lambda=3} = \frac{1}{4} \begin{bmatrix} -1 & 1 & 1 & -1 \\ -1 & 1 & 1 & -1 \\ 1 & -1 & -1 & 1 \\ 1 & -1 & -1 & 1 \end{bmatrix}$$

$$P_{21} = \frac{d}{d\lambda} \{ (\lambda-3)^3 [A(\lambda)]^{-1} \} \Big|_{\lambda=3} = \frac{1}{4} \begin{bmatrix} 0 & 0 & 2 & -2 \\ 2 & -2 & 0 & 0 \\ -2 & 2 & 0 & 0 \\ 0 & 0 & -2 & 2 \end{bmatrix}$$

$$P_{20} = \frac{1}{2} \frac{d}{d\lambda} \{ (\lambda-3)^2 [A(\lambda)]^{-1} \} \Big|_{\lambda=3} = \frac{1}{4} \begin{bmatrix} 3 & -1 & -1 & -1 \\ -1 & 3 & -1 & -1 \\ -1 & -1 & 3 & -1 \\ -1 & -1 & -1 & 3 \end{bmatrix}$$

The eigenprojectors agree with the previous values found from the eigenvectors.

The partial fraction expansion of  $[A(\lambda)]^{-1}$  is

$$(2.41) \quad [A(\lambda)]^{-1} = \frac{P_{10}}{\lambda-1} + \frac{P_{20}}{\lambda-3} + \frac{P_{21}}{(\lambda-3)^2} + \frac{P_{22}}{(\lambda-3)^3}$$

Two methods of computing the eigenprojectors (or matrix residues) have been discussed in this section. The first procedure given was based on the eigenvectors with the second method requiring the inversion of  $[A(\lambda)]^{-1}$  and evaluation of the residues. It has been shown that the two methods are equivalent although the numerical computations may not necessarily be comparable.

### 3. Lambda Matrices and Latent Projectors

The analysis in Section 2 was based on the assumption that the  $A$  matrix was in general form with  $m$  rows and columns. This assumption is valid when the differential equation describing the dynamics of a system are in the state variable form, first-order differential equations. The system equation may not always be in first-order form as it is common practice in some engineering disciplines to write the differential equations in  $m$ th-order form. If such is the practice, then lambda matrices will be encountered. This section considers lambda matrices or matrix polynomials, Gantmacher, [8].

Assume that  $\bar{A}(\lambda)$  is a matrix polynomial in  $\lambda$  of  $m$ th-order with  $n \times n$  coefficients of the form

$$(3.1) \quad \bar{A}(\lambda) = \bar{A}_0 \lambda^m + \bar{A}_1 \lambda^{m-1} + \dots + \bar{A}_{m-1} \lambda + \bar{A}_m$$

which Lancaster, [1], calls a lambda matrix. Dennis, Traub and Weber [3], make a distinction between  $A(\lambda)$  and  $A(X)$  where  $X$  is  $n \times n$  by calling the latter a matrix polynomial. The polynomial in (3.1) is commonly referred to as a matrix polynomial in control theory and that designation will be followed here.

The roots of  $\det[\bar{A}(\lambda)]$  are called latent roots and the vectors that satisfy  $[\bar{A}(\lambda_i)] \hat{y}_i$  are referred to as latent vector. This terminology will be followed in this work to avoid confusion with eigenvalues and eigenvectors. The concept of latent projectors will be introduced in this section with the latent projectors having an analogous role to eigenprojectors. It will be assumed that a latent root may have multiplicity  $r_i$  and that  $\bar{A}(\lambda)$  will be defective requiring  $\ell_i$  generalized latent vectors where a defective

lambda matrix has the same meaning as  $\Lambda$  being defective, there will not be  $r_i$  linearly latent vectors for the latent root  $\lambda_i$ .

If  $\Lambda_0$  is invertible, then (3.1) can be written as

$$(3.2) \quad \bar{\Lambda}(\lambda) = \bar{\Lambda}_0 [I\lambda^m + \Lambda_1 \lambda^{m-1} + \dots + \Lambda_{m-1} \lambda + \Lambda_m] = \bar{\Lambda}_0 \Lambda(\lambda)$$

where  $\Lambda_i = \bar{\Lambda}_0^{-1} \Lambda_i$ . The discussion that follows will focus on  $\Lambda(\lambda)$  although a complete treatment of lambda matrices should include the case when  $\Lambda_0$  is singular.

The latent roots of  $\Lambda(\lambda)$  will be denoted by  $\lambda_i$  with the right and left latent vectors, denoted by  $\hat{y}_i$  and  $\hat{z}_i$  respectively. The latent vectors for the latent roots  $\lambda_i$  satisfy

$$(3.3a) \quad \Lambda(\lambda_i) \hat{y}_i = 0$$

$$(3.3b) \quad \hat{z}_i \Lambda(\lambda_i) = 0$$

for the right and left latent vectors respectively when  $\lambda_i$  is distinct or  $\Lambda(\lambda_i)$  is not defective. If  $\Lambda(\lambda)$  is defective for  $\lambda_i$  then a chain rule must be employed. Lancaster and Webber, [9], have given the chain rule as

$$(3.4) \quad \Lambda(\lambda_i) \hat{y}_i^\ell + \frac{d\Lambda(\lambda_i)}{d\lambda} \hat{y}_i^{\ell-1} + \frac{1}{2} \frac{d^2\Lambda(\lambda_i)}{d\lambda^2} \hat{y}_i^{\ell-2} + \dots + \frac{1}{\ell-1} \frac{d^{\ell-1}\Lambda(\lambda_i)}{d\lambda^{\ell-1}} \hat{y}_i^1 = 0$$

with  $\hat{y}_i^1$  as a linearly independent latent vector. If  $\hat{y}_i^1$  is a linear independent latent vector than  $\hat{y}_i^2$  is given by

$$(3.5a) \quad \Lambda(\lambda_i) \hat{y}_i^2 = - \frac{d\Lambda(\lambda_i)}{d\lambda} \hat{y}_i^1$$

and  $\hat{y}_1^3$  by

$$(3.5b) \quad \Lambda(\lambda_1)\hat{y}_1^3 = -\frac{d\Lambda(\lambda_1)}{d\lambda}\hat{y}_1^2 - \frac{1}{2}\frac{d^2\Lambda(\lambda_1)}{d\lambda^2}\hat{y}_1^1$$

with all others computed by recursive use of (3.4). The chain rule for the left latent vector is similar except that  $\hat{z}_1^j$  is a premultiplier of the terms in (3.4).

The computation of the latent vectors is described in the example. Let  $\Lambda(\lambda)$  be

$$\Lambda(\lambda) = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \lambda^2 + \begin{bmatrix} -4.5 & 1.5 \\ 1.5 & -4.5 \end{bmatrix} \lambda + \begin{bmatrix} 5.5 & -3.5 \\ -3.5 & 5.5 \end{bmatrix}$$

which has latent roots  $\lambda_1 = 1$ ,  $\lambda_2 = 2$ ,  $\lambda_3 = 3$  and  $\lambda_4 = \lambda_3 = 3$  with  $\ell_3 = 1$ .

Let  $\lambda = 1$  then

$$\Lambda(1)\hat{y}_1 = \begin{bmatrix} 2 & -2 \\ -2 & 2 \end{bmatrix} \hat{y}_1 = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \quad \hat{y}_1 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

with  $\hat{y}_1$  as given for the latent vector. The latent vector for  $\lambda = 2$  is obtained from

$$\Lambda(2)\hat{y}_2 = \begin{bmatrix} 0.5 & -0.5 \\ -0.5 & 0.5 \end{bmatrix} \hat{y}_2 = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \quad \hat{y}_2 = \begin{bmatrix} -1 \\ -1 \end{bmatrix}$$

The linear independent latent vector for  $\lambda = 3$  is found from

$$\Lambda(3)\hat{y}_3^1 = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix} \hat{y}_3^1 = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \quad \hat{y}_3^1 = \begin{bmatrix} 1 \\ -1 \end{bmatrix}$$

and the generalized latent vector for  $\lambda = 3$ ,  $\ell_3 = 1$ , is computed from the

chain rule. Using (3.5a)

$$\Lambda(3)\hat{y}_3^2 = -\frac{d\Lambda(3)}{d\lambda}\hat{y}_3^1 = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix} \hat{y}_3^2 = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

or  $(\hat{y}_3^2)^T = [-1 \ 1]$ . The latent vectors  $\hat{y}_2$  and  $\hat{y}_3^2$  are chosen with signs opposite to  $\hat{y}_1$  and  $\hat{y}_3^1$  respectively for convenience. Since  $y_1$  is in a two-dimensional space, only two of the latent vectors are necessary to span the space. It should be noted that the term linearly independent latent vector is not proper terminology and will be dropped in favor of latent vector hereafter if  $\hat{y}_i$  satisfies  $\Lambda(\lambda_i)\hat{y}_i = 0$ .

The concept of distinct and repeated latent roots as well as a defective matrix polynomial will be clarified by relating  $\Lambda(\lambda)$  to the companion form of the  $m \times m$  matrix  $A$ . It is well known that the matrix

$$(3.6) \quad A = \begin{bmatrix} 0 & I & 0 & \dots & 0 \\ 0 & 0 & I & \dots & 0 \\ \vdots & \vdots & \vdots & \dots & \vdots \\ \vdots & \vdots & \vdots & \dots & \vdots \\ \vdots & \vdots & \vdots & \dots & \vdots \\ 0 & 0 & 0 & \dots & I \\ -A_m & -A_{m-1} & -A_{m-2} & \dots & -A_1 \end{bmatrix}$$

will have eigenvalues  $\lambda_i$  that are the same as the latent roots of  $\Lambda(\lambda)$  that is  $\det[A(\lambda)] = \det[\Lambda(\lambda)]$ . Furthermore, it can be shown that the latent vectors are subvectors of the eigenvectors of  $A$ . If  $y_i$  is an eigenvector of  $A$  for an eigenvalue  $\lambda_i$  then  $\hat{y}_i$  is a subvector of  $y_i$ . The eigenvector  $y_i$  of  $A$  is given by



$$(3.7) \quad A(\lambda_1)y_1 = \begin{bmatrix} \lambda_1 I & -I & 0 & \dots & 0 \\ 0 & \lambda_1 I & -I & \dots & 0 \\ \vdots & \vdots & \vdots & \dots & \vdots \\ \vdots & \vdots & \vdots & \dots & \vdots \\ 0 & 0 & 0 & \dots & -I \\ A_m & A_{m-1} & A_{m-2} & \dots & \lambda_1 I + A_1 \end{bmatrix} \begin{bmatrix} \hat{y}_1 \\ \lambda_1 \hat{y}_1 \\ \vdots \\ \vdots \\ \lambda_1^{m-1} \hat{y}_1 \end{bmatrix}$$

when  $\lambda_1$  is a distinct eigenvalue or  $A$  is not defective. It follows that the first  $n$  elements of  $y_1$  is a latent vector of  $A(\lambda)$ . Similarly if  $z_1$  is a left eigenvector of  $A$  then the left latent vector of  $A(\lambda)$  will be the last  $n$ -elements of the row vector  $z_1$  under the same restriction on  $\lambda_1$ .

There is a second relationship between  $A(\lambda)$  and  $A(\lambda)$  that will be useful in the development that follows. The inverse of  $A(\lambda)$  in companion form is given by

$$(3.8) \quad [A(\lambda)]^{-1} = \begin{bmatrix} \lambda I & -I & 0 & \dots & 0 \\ 0 & \lambda I & -I & \dots & 0 \\ \vdots & \vdots & \vdots & \dots & \vdots \\ \vdots & \vdots & \vdots & \dots & \vdots \\ 0 & 0 & 0 & \dots & -I \\ A_m & A_{m-1} & A_{m-2} & \dots & \lambda I + A_1 \end{bmatrix}^{-1} = [A(\lambda)]^{-1} B_{\text{adj}}[A(\lambda)]$$

where  $B_{\text{adj}}[A(\lambda)]$  denotes the block adjoint of  $A(\lambda)$ . The block adjoint is defined as the adjoint matrix of  $A(\lambda)$  with each block matrix of  $A(\lambda)$  treated as a scalar element. As an example the block adjoint of  $A(\lambda)$  with  $m = 3$  is

$$(3.9) \quad B_{\text{adj}}[A(\lambda)] = B_{\text{adj}} \begin{bmatrix} \lambda I & -I & 0 \\ 0 & \lambda I & -I \\ A_3 & A_2 & \lambda I + A_1 \end{bmatrix} = \begin{bmatrix} \lambda^2 I + A_1 \lambda + A_2 & \lambda I + A_1 & I \\ -A_3 & \lambda^2 I + \lambda A_1 & \lambda I \\ -\lambda A_3 & -\lambda A_2 - A_3 & \lambda^2 I \end{bmatrix}$$

The last column of  $B_{\text{adj}}[A(\lambda)]$  will always have the form shown with the last block equal to  $\lambda^{m-1}I$ .

Consider the inverse of  $[A(\lambda)]$  and let  $A(\lambda)$  have distinct latent roots. The partial fraction expansion of  $[A(\lambda)]$  was given by

$$(3.10) \quad [A(\lambda)]^{-1} = \sum_{i=1}^{mn} \frac{P_{i0}}{\lambda - \lambda_i}$$

where  $P_{i0} = (\lambda - \lambda_i) [A(\lambda)]^{-1}$  evaluated at  $\lambda = \lambda_i$ . It follows from (3.9) and (3.10) that the eigenprojector  $P_{i0}$  and the latent projector  $\hat{P}_{i0}$  are related since

$$(3.11) \quad P_{i0} = \hat{P}_{i0} B_{\text{adj}}[A(\lambda_i)] = \begin{bmatrix} \dots & \hat{P}_{i0} \\ \dots & \lambda_i \hat{P}_{i0} \\ \dots & \vdots \\ \dots & \lambda_i^{m-1} \hat{P}_{i0} \end{bmatrix}$$

where the first  $m-1$  columns of (3.11) are not important to the development. It was shown earlier that  $\sum_{i=1}^{mn} P_{i0} = I$  thus it follows from (3.11) that the latent projectors have the properties that

$$(3.12a) \quad \sum_{i=1}^{mn} \hat{P}_{i0} = 0$$

with

$$(3.12b) \quad \sum_{i=1}^{mn} \lambda_i^j \hat{P}_{i0} = 0 \quad j = 1, 2, \dots, m-2$$

$$(3.12c) \quad \sum \lambda_i^{m-1} \hat{P}_{i0} = I$$

where  $\hat{P}_{i0}$  are the primary latent projectors. If  $P_{i0}$  is a matrix residue of  $[A(\lambda)]^{-1}$  then  $P_{i0}$  is a matrix of  $[A(\lambda)]^{-1}$ .

The partial fraction expansion of  $[A(\lambda)]^{-1}$  for distinct latent roots can be obtained from (3.10) and (3.11) and is

$$(3.13) \quad [A(\lambda)]^{-1} = \sum_{i=1}^{mn} \frac{\hat{P}_{i0}}{\lambda - \lambda_i}$$

where

$$(3.14) \quad \hat{P}_{i0} = (\lambda - \lambda_i) [A(\lambda)]^{-1} \Big|_{\lambda = \lambda_i}$$

with  $\hat{P}_{i0}$  being primary latent projectors.

It was shown in Section 2 that the eigenprojectors for the  $A$  matrix with repeated eigenvalues are given by

$$(3.15) \quad P_{i, \ell_i - j} = \frac{1}{(\ell_i - r_i + j + 1)!} \frac{d^j}{d\lambda^j} \{ (\lambda - \lambda_i)^{r_i} [A(\lambda_i)]^{-1} \}$$

with  $j = 0, 1, 2, \dots, r_i$  and  $(\cdot)! = 1$  for  $(\cdot) \leq 0$ .

Using (3.11), it follows that

$$(3.16) \quad P_{i, \ell_i - j} = \frac{1}{j!} \frac{d^j}{d\lambda^j} \{ (\lambda - \lambda_i)^{r_i} \hat{P}_{i0} [A(\lambda)]^{-1} \}$$

Equation (3.16) agrees with the usual partial fraction expansion formula provided that  $[A(\lambda)]^{-1}$  is a minimum polynomial, factors common to  $\text{adj}[A(\lambda)]$  and  $\det[A(\lambda)]$  have been cancelled.

The numerical procedure for computing the latent projectors by the

residues will be given for the matrix polynomial

$$A(\lambda) = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \lambda^2 + \begin{bmatrix} -5 & 1 \\ 1 & -5 \end{bmatrix} \lambda + \begin{bmatrix} 6 & -3 \\ -3 & 6 \end{bmatrix}$$

which has latent roots  $\lambda_1 = 1$ ,  $\lambda_2 = \lambda_3 = \lambda_4 = 3$  and  $\ell_2 = 1$ . The inverse of  $A(\lambda)$  is

$$[A(\lambda)]^{-1} = \frac{1}{(\lambda-1)(\lambda-3)^3} \begin{bmatrix} \lambda^2 - 5\lambda + 6 & -\lambda + 3 \\ -\lambda + 3 & \lambda^2 - 5\lambda + 6 \end{bmatrix} = \frac{1}{(\lambda-1)(\lambda-3)^2} \begin{bmatrix} \lambda-2 & -1 \\ -1 & \lambda-2 \end{bmatrix}$$

thus

$$\hat{P}_{10} = (\lambda-1)[A(\lambda)]^{-1} \Big|_{\lambda=1} = \frac{1}{4} \begin{bmatrix} -1 & -1 \\ -1 & -1 \end{bmatrix}$$

Since  $\lambda = 3$  appears twice in the minimum form of  $[A(\lambda)]^{-1}$  and  $\ell = 1$ , there will be one primary and one secondary latent projector. The latent projectors are

$$P_{21} = \frac{1}{2} \begin{bmatrix} 1 & -1 \\ -1 & 1 \end{bmatrix}$$

$$P_{20} = \frac{1}{4} \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$$

The eigenprojector of  $A$  were constructed from the eigenvectors of  $A$  in Section 2. Since the eigenvectors of  $A$  have the latent vectors as subvectors, the latent projectors of  $A(\lambda)$  can be constructed from the latent vectors. Consider the distinct eigenvalues  $\lambda_i$  for which the eigenprojectors  $P_{i0}$  are given by

$$(3.17) \quad P_{i0} = \frac{y_i z_i}{z_i y_i}$$

Let the order of the matrix polynomial  $m$  be 2 then  $y_i$  satisfies the equation

$$(3.18) \quad \begin{bmatrix} \lambda_i I & -I \\ A_2 & \lambda_i I + A_1 \end{bmatrix} y_i = \begin{bmatrix} \lambda_i I & -I \\ A_2 & \lambda_i I + A_1 \end{bmatrix} \begin{bmatrix} \hat{y}_i \\ \lambda_i \hat{y}_i \end{bmatrix}$$

The left eigenvector  $z_i$  must satisfy

$$(3.19) \quad z_i \begin{bmatrix} \lambda_i I & -I \\ A_2 & \lambda_i I + A_1 \end{bmatrix} = \hat{z}_i [-\beta_i I] \begin{bmatrix} \lambda_i I & -I \\ A_2 & \lambda_i I + A_1 \end{bmatrix}$$

from which it follows that

$$(3.20) \quad z_i = \hat{z}_i [\lambda_i I + A_1 I]$$

thus the numerator of (3.17) is

$$(3.21) \quad y_i z_i = \begin{bmatrix} I \\ \lambda_i \end{bmatrix} \hat{y}_i \hat{z}_i [\lambda_i I + A_1 I]$$

$$= \begin{bmatrix} \hat{y}_i \hat{z}_i (\lambda_i I + A_1) & \hat{y}_i \hat{z}_i \\ -\hat{y}_i \hat{z}_i A_2 & \lambda_i \hat{y}_i \hat{z}_i \end{bmatrix}$$

The outer product  $z_i y_i$  is

$$(3.22) \quad z_i y_i = \hat{z}_i [\lambda_i I + A_1 I] \begin{bmatrix} I \\ \lambda_i \end{bmatrix} \hat{y}_i = \hat{z}_i \frac{dA(\lambda_i)}{d\lambda} \hat{y}_i$$

which gives for the distinct eigenvalue or latent root the eigenprojector

$$(3.23) \quad P_{10} = \frac{1}{\hat{z}_1 \frac{dA(\lambda_1)}{d\lambda} \hat{y}_1} \begin{bmatrix} \hat{y}_1 \hat{z}_1 (\lambda_1 I + A_1) & \hat{y}_1 \hat{z}_1 \\ -\hat{y}_1 \hat{z}_1 A_2 & \lambda_1 \hat{y}_1 \hat{z}_1 \end{bmatrix}$$

The latent projector, as given in (3.11) is the (1,2) block of  $P_{10}$  or

$$(3.24) \quad \hat{P}_{10} = \frac{\hat{y}_1 \hat{z}_1}{\hat{z}_1 \frac{dA(\lambda_1)}{d\lambda} \hat{y}_1}$$

for the distinct latent root. The eigenprojector for the repeated eigenvalue nondefective case was given as

$$(3.25) \quad P_{10} = \sum_{j=1}^r \frac{y_1^j z_1^j}{z_1^j y_1^j}$$

with the obvious extension to the latent projectors as given in (3.26)

$$(3.26) \quad \hat{P}_{10} = \sum_{j=1}^r \frac{\hat{y}_1^j \hat{z}_1^j}{\hat{z}_1^j \frac{dA(\lambda_1)}{d\lambda} \hat{y}_1^j}$$

Consider the matrix polynomial

$$A(\lambda) = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \lambda^2 + \begin{bmatrix} -4.5 & 0.5 \\ 0.5 & -4.5 \end{bmatrix} \lambda + \begin{bmatrix} 4.5 & -1.5 \\ -1.5 & 4.5 \end{bmatrix}$$

for which the latent projectors are to be found. The latent vectors for the latent roots  $\lambda_1 = 1$ ,  $\lambda_2 = 2$  and  $\lambda_3 = \lambda_4 = 3$  are

$$\hat{y}_1 = \begin{bmatrix} 1 \\ 1 \end{bmatrix} \quad \hat{y}_2 = \begin{bmatrix} -1 \\ 1 \end{bmatrix} \quad \hat{y}_3^1 = \begin{bmatrix} -1 \\ -1 \end{bmatrix} \quad \hat{y}_3^2 = \begin{bmatrix} 1 \\ -1 \end{bmatrix}$$

$$\hat{z}_1 = [-1 \ -1] \quad \hat{z}_2 = [1 \ -1] \quad \hat{z}_3^1 = [-1 \ -1] \quad \hat{z}_3^2 = [1 \ -1]$$

The two distinct latent projectors are

$$\hat{P}_{10} = \frac{\hat{y}_1 \hat{z}_1}{\hat{z}_1 \frac{d\Lambda(1)}{d\lambda} \hat{y}_1} = \frac{1}{4} \begin{bmatrix} -1 & -1 \\ -1 & -1 \end{bmatrix}$$

and

$$\hat{P}_{20} = \frac{\hat{y}_2 \hat{z}_2}{\hat{z}_2 \frac{d\Lambda(2)}{d\lambda} \hat{y}_2} = \frac{1}{4} \begin{bmatrix} -2 & -2 \\ -2 & -2 \end{bmatrix}$$

The latent projector for the repeated latent root is

$$\hat{P}_{30} = \frac{\hat{y}_3^1 \hat{z}_3^1}{\hat{z}_3^1 \frac{d\Lambda(3)}{d\lambda} \hat{y}_3^1} + \frac{\hat{y}_3^2 \hat{z}_3^2}{\hat{z}_3^2 \frac{d\Lambda(3)}{d\lambda} \hat{y}_3^2} = \frac{1}{4} \begin{bmatrix} 3 & -1 \\ -1 & 3 \end{bmatrix}$$

The partial fraction expansion of  $[\Lambda(\lambda)]^{-1}$  is then

$$[\Lambda(\lambda)]^{-1} = \frac{1}{4(\lambda-1)} \begin{bmatrix} -1 & -1 \\ -1 & -1 \end{bmatrix} + \frac{1}{4(\lambda-2)} \begin{bmatrix} -2 & 2 \\ 2 & -2 \end{bmatrix} + \frac{1}{4(\lambda-3)} \begin{bmatrix} 3 & -1 \\ -1 & 3 \end{bmatrix}$$

with  $\hat{P}_{10} + \hat{P}_{20} + \hat{P}_{30} = 0$ , and  $\lambda_1 \hat{P}_{10} + \lambda_2 \hat{P}_{20} + \lambda_3 \hat{P}_{30} = I$  as required.

The formulation of the latent projectors for the repeated latent root polynomial when  $\Lambda$  is defective in terms of the latent vectors remains as a problem. Attempts to formulate the latent projectors for the defective case have been unsuccessful. Future work will be devoted to this problem.

#### 4. Projectors and the Sign Matrix

It was shown in Section 2 that the primary eigenprojectors for the  $m \times m$   $A$  matrix are given by

$$(4.1) \quad P_{i0} = Q F_{i0} Q^{-1} \quad i = 1, 2, \dots, q$$

where  $F_{i0}$  is a diagonal matrix with ones along the diagonal of the  $i$ th pseudo-Jordan block. The secondary eigenprojectors were defined as

$$(4.2) \quad P_{ij} = Q F_{ij} Q^{-1} \quad j = 1, 2, \dots, l_i$$

when  $A$  has  $l_i$  generalized eigenvectors for the eigenvalue  $\lambda_i$ .

Assume that  $A$  has  $q_1$  eigenvalues with  $\text{Re}(\lambda_i) > 0$  and  $q_2$  eigenvalues having  $\text{Re}(\lambda_i) < 0$  and no eigenvalues with  $\text{Re}(\lambda_i) = 0$  so that  $q_1 + q_2 = q$ . Let  $P^+$  be denoted as the sum of the eigenprojectors with  $\text{Re}(\lambda_i) > 0$  and  $P^-$  the sum of the eigenprojectors with  $\text{Re}(\lambda_i) < 0$ ; that is

$$(4.3) \quad P^+ = \sum_{i=1}^{q_1} P_{i0} = Q \sum_{i=1}^{q_1} F_{i0} Q^{-1}$$

$$(4.4) \quad P^- = \sum_{i=q_1+1}^q P_{i0} = Q \sum_{i=q_1+1}^q F_{i0} Q^{-1}$$

where it has been assumed that the first  $q_1$  eigenvalues have  $\text{Re}(\lambda_i) > 0$ .

The sign of a matrix, denoted by  $S$ , will be defined as the matrix

$$(4.5) \quad S = Q \{\text{sign}[\text{Re}(\Lambda)]\} Q^{-1}$$



where  $\text{Re}(\Lambda)$  denotes the real part of the eigenvalues of  $A$  or the diagonal elements of the Jordan matrix  $J$ . Let  $E_1$  be the  $m \times m$  matrix with diagonal elements equal to 1 if  $\text{Re}(\lambda_i) > 0$  and zero for  $\text{Re}(\lambda_i) < 0$  and  $E_2$  be the complement to  $E_1$  such that  $E_1 + E_2 = I$ . The sign matrix can then be defined as

$$(4.6) \quad S = Q E_1 Q^{-1} - Q E_2 Q^{-1}$$

$$= Q \left[ \sum_{i=1}^q 1 F_{i0} - \sum_{i=q_1+1}^q 1 F_{i0} \right] Q^{-1}$$

therefore  $S$  is equal to

$$(4.7) \quad S = P^+ - P^-$$

Knowledge of the eigenprojectors is sufficient to construct the sign matrix, similarly it can be shown that knowledge of the sign matrix is sufficient to construct  $P^+$  and  $P^-$ . Assume that  $P^+$  is given as

$$(4.8) \quad P^+ = \frac{1}{2} (S+I) = \frac{1}{2} [Q E_1 Q^{-1} - Q E_2 Q^{-1} + Q Q^{-1}]$$

but since  $E_1 + E_2 = I$  then

$$(4.9) \quad P^+ = \frac{1}{2} Q [E_1 - E_2 + E_1 + E_2] Q^{-1} = Q E_1 Q^{-1} = Q \sum_{i=1}^q 1 F_{i0} Q^{-1}$$

It is not difficult to show that  $P^-$  is given by

$$(4.10) \quad P^- = \frac{1}{2} [I - S]$$

The computation of the sign of  $A$  is a rather simple task. Roberts [10], gave an iterative algorithm to compute  $S$  which is based on Newton's method for computing the square root of  $S^2 = I$ . The algorithm is

$$(4.11) \quad S(i+1) = \frac{1}{2} \{S(i) + [S(i)]^{-1}\} \quad S(0) = A$$

where the index  $i$  denotes the  $i$ th iteration. The algorithm will converge quadratically to  $S$  provided that  $A$  has no eigenvalues on the  $j\omega$  axis. The simplest test of convergence of (4.11) to the sign of  $A$  is to compute the trace of  $S^2(i)$  at each iteration. Since  $S^2(i)$  converges to  $I$ , then trace  $[S]$  will be  $mn$ .

Several accelerated versions of (4.11) have been described in the literature, Roberts [10], Hoskins and Walton, [11], and Mattheys, [12]. Numerous applications of the sign algorithm to system analysis have been given in the literature, [13]-[16].

The example below gives the sign of  $A$  where  $A$  is

$$A = \frac{1}{4} \begin{bmatrix} -1 & -1 & 9 & -3 \\ -3 & 1 & -1 & 7 \\ 9 & -3 & -1 & -1 \\ -1 & 7 & -3 & 1 \end{bmatrix}$$

with eigenvalues  $\lambda_1 = 1$ ,  $\lambda_2 = 3$ ,  $\lambda_3 = \lambda_4 = -2$  and  $\rho_3 = 1$ . The sign of  $A$  is

$$S = \begin{bmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{bmatrix}$$

which converged in 5 iterations. The trace of  $[S(1)]^2$  was

<u>iteration</u>	<u>trace</u>
1	6.90278
2	4.3857
3	4.01697
4	4.00006
5	4.00000

The positive and negative projectors,  $P^+$  and  $P^-$ , were found to be

$$P^+ = \frac{1}{2} \begin{bmatrix} 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \\ 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \end{bmatrix} = P_{10} + P_{20}$$

$$P^- = \frac{1}{2} \begin{bmatrix} 1 & 0 & -1 & 0 \\ 0 & 1 & 0 & -1 \\ -1 & 0 & 1 & 0 \\ 0 & -1 & 0 & 1 \end{bmatrix} = -P_{30}$$

The sign matrix of  $A$  when all eigenvalues have  $\text{Re}(\lambda_i) > 0$  will be  $I$  whereas the sign of  $A$  with  $\text{Re}(\lambda_i) < 0$  will be  $-I$ . Eigenvalues along the  $j\omega$  axis can be removed from the axis by an origin shift or by computing the sign of  $(A + \rho I)$  where  $\rho$  is a real number. The eigenvalues of  $A + \rho I$  will be  $\lambda_i + \rho$  since  $A + \rho I = Q(J + \rho I)Q^{-1}$ . Eigenvalues belonging to a Jordan block cannot be separated nor can eigenvalues along the  $j\omega$  axis be split by the described procedure.

A method of separating eigenvalues according to their magnitude is to compute a new matrix by the bilinear transformation

$$(4.12) \quad A_0 = (A - \rho I)(A + \rho I)^{-1}$$

where  $\rho$  has the same meaning as above. All of the eigenvalues with  $|\lambda_i| < \rho$  will be mapped into the half plane  $\text{Re}(\lambda_i) < 0$  with others mapped into the plane with  $\text{Re}(\lambda_i) > 0$ . This procedure is more general than the origin shifting method since the spectrum splitting will be according to the magnitudes of  $\lambda_i$ . The two methods, shifting and splitting can be combined if desired to isolate any circular region of the eigenvalue space. For example, the matrix  $A_0$  given by

$$(4.13) \quad A_0 = (A + \rho_1 I - \rho_2 I)(A + \rho_1 I + \rho_2 I)^{-1}$$

can be used to isolate eigenvalues inside a circle of radius  $\rho_2$  centered at  $\rho_1$ .

As an example of the bilinear transformation procedure, let

$$A = \frac{1}{2} \begin{bmatrix} 3 & -2 & -9 & 6 \\ -2 & 3 & 6 & -9 \\ -9 & 6 & 3 & -2 \\ 6 & -9 & -2 & 3 \end{bmatrix} \quad \text{trace} = 6$$

which has eigenvalues  $\lambda_1 = -1$ ,  $\lambda_2 = 2$ ,  $\lambda_3 = -5$  and  $\lambda_4 = 10$ . If the value of  $\rho = 4$  is selected then all eigenvalues inside of the circle  $\rho = 4$  will be mapped to the left half plane and those outside the circle will be in the right half plane. The sign of  $A$  is

$$S = \begin{bmatrix} 0 & -1 & 0 & 0 \\ -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 \\ 0 & 0 & -1 & 0 \end{bmatrix}$$

with projectors

$$P^+ = \frac{1}{2} \begin{bmatrix} 1 & -1 & 0 & 0 \\ -1 & 1 & 0 & 0 \\ 0 & 0 & 1 & -1 \\ 0 & 0 & -1 & 1 \end{bmatrix}$$

$$P^- = \frac{1}{2} \begin{bmatrix} 1 & 1 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 1 & 0 \end{bmatrix}$$

The projectors  $P^+$  and  $P^-$  are idempotent matrices and satisfy the properties of the eigenprojector given earlier. It is not difficult to show that the positive and negative projectors can be used for the spectral decomposition of  $A$ . Since  $P^+ = Q E_1 Q^{-1}$  with  $E_1 = \text{diag}[1 \ 1 \ 1 \ 1 \ . \ . \ 0 \ 0 \ 0]$  with the ones in the first  $q_1$  locations. The product  $AP^+$  will have the eigenvalues with  $|\lambda_i| > \rho$  whereas  $AP^-$  will have eigenvalues  $|\lambda_i| < \rho$  with all other eigenvalues zero. Using the example

$$A^+ = AP^+ = \frac{1}{4} \begin{bmatrix} 5 & 5 & -15 & 15 \\ -5 & 5 & 15 & -15 \\ -15 & 15 & 5 & -5 \\ 15 & -15 & -5 & 5 \end{bmatrix} \quad \text{trace} = 5 = \lambda_4 - \lambda_3$$

$$A^- = AP^- = \frac{1}{4} \begin{bmatrix} 1 & 1 & -3 & -3 \\ 1 & 1 & -3 & -3 \\ -3 & -3 & 1 & 1 \\ 3 & -3 & 1 & 1 \end{bmatrix} \quad \text{trace} = 1 = \lambda_2 - \lambda_1$$

The sum of  $AP^+$  and  $AP^-$  must be  $A$  since  $P^+ + P^- = I$ .

It is obvious that  $P^+$  and  $P^-$  can be decomposed into eigenprojectors for the eigenvalues  $\lambda_1, \lambda_2, \lambda_3$  and  $\lambda_4$ . The details for the determination of the eigenprojectors have been covered in Section 2 and will not be covered at this point.

The positive and negative projectors,  $P^+$  and  $P^-$ , have been defined in (4.3) and (4.4). It follows from (4.3) and (3.11) that  $P^+$  is also given by

$$(4.13) \quad P^+ = \sum_{i=1}^q P_{i0} = \begin{bmatrix} \sum_{i=1}^q \hat{P}_{i0} (\lambda_i I + A_1) & \sum_{i=1}^q \hat{P}_{i0} \\ -\sum_{i=1}^q \hat{P}_{i0} \Lambda_2 & \sum_{i=1}^q \lambda_i \hat{P}_{i0} \end{bmatrix}$$

for the companion form when  $A(\lambda)$  is a second-order polynomial.

Similarly,  $P^- = I - P^+$  thus

$$(4.14) \quad P^- = -\sum_{i=q_1+1}^q P_{i0} = \begin{bmatrix} -\sum_{i=q_1+1}^q \hat{P}_{i0} (\lambda_i I + A_1) & -\sum_{i=q_1+1}^q \hat{P}_{i0} \\ \sum_{i=q_1+1}^q \hat{P}_{i0} \Lambda_2 & -\sum_{i=q_1+1}^q \lambda_i \hat{P}_{i0} \end{bmatrix}$$

The secondary eigenprojectors and latent projectors are not needed in the decomposition of  $A$  into  $A^+$  and  $A^-$ . This can be shown from the definition of  $A^+$  and  $A^-$ .

The individual eigenprojectors  $P_{i0}$  can be computed by repeated use of the sign algorithm. Assume that  $n = 4$  with  $\lambda_1 = 1, \lambda_2 = 2, \lambda_3 = 3$  and  $\lambda_4 = 4$ . The bilinear transformation can be carried out first with  $\rho = 1.5$

to separate  $\lambda_1$  from the other eigenvalues. If  $S_1$  denotes the sign of  $A_{01}$  with

$$(4.14) \quad A_{01} = (A - 1.5I)(A + 1.5I)^{-1}$$

then  $P_{10}$  will be given by

$$(4.15) \quad P_{10} = \frac{1}{2} (I - S_1) = P_1^-$$

with  $P^+ = \sum_{i=2}^3 P_{i0}$ . The next step in the procedure is to compute  $A_{02}$  with  $\rho_2 = 2.5$  and compute the sign of  $A_{02}$ . The negative projector of  $A_{02}$  will be the sum of  $P_{10}$  and  $P_{20}$  or

$$P_2^- = P_{10} + P_{20} = \frac{1}{2} (I - S_2) .$$

Thus  $P_{20} = P_2^- - P_{10}$ . This process can be continued until each eigenprojector has been found. Since the eigenprojectors also give the latent projectors as the upper right block, the latent projectors will also be known when  $A$  is in companion form.

A method of computing the projectors of a matrix has been discussed in this section. It has been shown that the eigenprojector for any general matrix can be computed from the sign of a matrix. If  $A$  is in companion form, the latent projectors can also be found from the eigenprojectors.

## 5. Solvents of Matrix Polynomials

The concept of matrix polynomials was introduced in Section 1 of this report where it was shown that  $\Lambda(\lambda)$  arises when  $n$  simultaneous equations of  $m$ th-order are used to define the time-behavior of a dynamic system. The eigenprojectors for a matrix in companion form was discussed in Section 2 and the latent projectors were described in Section 3. The application of lambda matrices to the dynamics of systems has been described by Frazer, Duncan and Collar, [17], in their book Elementary Matrices and Some Applications to Dynamics and Differential Equations. The concept of solvents or matrix roots of a matrix polynomial will be given in this section. It will be shown in the next section that solvents are useful in solving sets of differential equations.

Let  $A(\lambda)$  be defined as a  $m$ th-order matrix polynomial with  $n \times n$  matrix coefficients. The associated  $mn \times mn$   $A$  matrix, which will be called the block companion matrix, is given by

$$(5.1) \quad A = \begin{bmatrix} 0 & I & 0 & \dots & 0 \\ 0 & 0 & I & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \dots & I \\ -\Lambda_m & -\Lambda_{m-1} & -\Lambda_{m-2} & \dots & -\Lambda_1 \end{bmatrix}$$

The eigenvector matrix  $Q$  for  $A$  will always have the form

$$(5.2) \quad Q = \begin{bmatrix} Q_1 & Q_2 & \dots & Q_m \\ Q_1 \Lambda_1 & Q_2 \Lambda_2 & \dots & Q_m \Lambda_m \\ Q_1 \Lambda_1^2 & Q_2 \Lambda_2^2 & \dots & Q_m \Lambda_m^2 \\ \vdots & \vdots & \ddots & \vdots \\ Q_1 \Lambda_1^{m-1} & Q_2 \Lambda_2^{m-1} & \dots & Q_m \Lambda_m^{m-1} \end{bmatrix}$$



when  $A$  has distinct eigenvalues or  $\Lambda(\lambda)$  has distinct latent roots. The submatrices  $Q_j$  will be a matrix of latent vectors  $\hat{y}_j$  for the latent roots  $\lambda_j$ . It will be assumed that  $Q_j$  exists and is invertible; under the above assumptions,  $R_j = Q_j \Lambda_j Q_j^{-1}$  is a solvent of the matrix polynomial and satisfies the equation, [1],

$$(5.3) \quad R_j^m + A_1 R_j^{m-1} + \dots + A_m = 0 \quad j = 1, 2, \dots, m$$

The proof of this is straightforward if  $\Lambda(\lambda)$  is considered. Let  $\Lambda(\lambda)$  be defined as the block matrix  $\Lambda I - A$ ; it then follows that

$$(5.4) \quad \begin{bmatrix} \Lambda_j & -I & 0 & 0 & 0 \\ \cdot & 0 & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot \\ 0 & 0 & 0 & \Lambda_j & -I \\ A_m & A_{m-1} & A_{m-2} & A_2 & \Lambda_j + A_1 \end{bmatrix} \begin{bmatrix} Q_j \\ Q_j \Lambda_j \\ \cdot \\ \cdot \\ Q_j \Lambda_j^{m-1} \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ \cdot \\ \cdot \\ 0 \end{bmatrix}$$

where (5.3) is given by the last row of (5.4) provided that  $\Lambda_j Q_j = Q_j \Lambda_j$  which must hold from the first row of (5.4).

The block matrices  $\Lambda_j$  are  $n \times n$  diagonal matrices constructed from a subset of the latent roots of  $\Lambda(\lambda)$  or the eigenvalues of  $A$ . Each  $R_j$  is defined in terms of  $n$  latent roots  $\lambda_i$  and  $n$  latent vectors  $\hat{y}_i$ , as an example, consider the matrix polynomial  $\Lambda(\lambda)$  with

$$\Lambda(\lambda) = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \lambda^2 + \begin{bmatrix} -5 & 2 \\ 2 & -5 \end{bmatrix} \lambda + \begin{bmatrix} 7 & -5 \\ -5 & 7 \end{bmatrix}$$

which has latent roots  $\lambda_1 = 1$ ,  $\lambda_2 = 2$ ,  $\lambda_3 = 3$  and  $\lambda_4 = 4$ . The right latent vectors of  $A(\lambda)$  are

$$\hat{y}_1 = \begin{bmatrix} 1 \\ 1 \end{bmatrix} \quad \hat{y}_2 = \begin{bmatrix} -1 \\ -1 \end{bmatrix} \quad \hat{y}_3 = \begin{bmatrix} -1 \\ 1 \end{bmatrix} \quad \hat{y}_4 = \begin{bmatrix} 1 \\ -1 \end{bmatrix}$$

The solvents are constructed from the latent roots and latent vectors by forming the  $2 \times 2$  matrices  $Q_j \Lambda_j Q_j^{-1}$  where  $Q_j = \{\hat{y}_j\}$  such that  $Q_j$  is invertible with  $j$  indicating a subset of latent vectors. Noting that  $\{y_1, y_2\}$  is singular, then

$$R_1 = [\hat{y}_1 \hat{y}_3] \text{diag}[\lambda_1 \lambda_3] [\hat{y}_1 \hat{y}_3]^{-1} = \begin{bmatrix} 2 & -1 \\ -1 & 2 \end{bmatrix}$$

$$R_2 = [\hat{y}_2 \hat{y}_4] \text{diag}[\lambda_2 \lambda_4] [\hat{y}_2 \hat{y}_4]^{-1} = \begin{bmatrix} 3 & -1 \\ -1 & 3 \end{bmatrix}$$

It can be shown that  $R_1$  and  $R_2$  satisfy (5.3).

The eigenprojectors for the companion matrix  $A$  are given by

$$(5.5) \quad P_{i0} = \hat{P}_{i0} \begin{bmatrix} \dots & I \\ \dots & \lambda_i I \\ \dots & \lambda_i^2 I \\ \dots & \cdot \\ \dots & \cdot \\ \dots & \lambda_i^{m-1} I \end{bmatrix}$$

where the first  $m-1$  block columns are not important in the development.

Suppose that the latent projectors are known and  $A$  has distinct eigenvalues.

It can be shown that if  $Q_j$  is defined as

$$(5.6) \quad Q_{j+1} \equiv \sum_{i=jn+1}^{(j+1)n} \hat{P}_{i0} \quad j = 0, 1, 2, \dots, m-1$$

and

$$(5.7) \quad Q_{j+1} \Lambda_{j+1} \equiv \sum_{i=jn+1}^{(j+1)n} \lambda_i \hat{P}_{i0}$$

then  $R_j$  is defined by the latent projectors of  $\Lambda(\lambda)$  provided that  $Q_j$  is invertible. If  $Q_j$  is singular, the latent projectors are reordered until a set is found for  $Q_j^{-1}$  to exist.

The solvents for repeated roots can be defined by noting that  $\Lambda_j$  may include several Jordan blocks. Assuming that the multiplicity  $r_i$  is less than  $n$ , and that full Jordan blocks are included in  $\Lambda_j$ ,  $Q_j$  is defined

as in (5.6). The product  $Q_j \Lambda_j$  must be modified since  $\Lambda_j$  is no longer diagonal but may include the ones on the super diagonal of  $\Lambda_j$  due to the included Jordan block. Assuming that the Jordan block is defective, then  $Q_j \tilde{J}_j$  will be defined as

$$(5.8) \quad Q_j \tilde{J}_j = \sum_{i=jn+1}^{(j+1)n} [\lambda_i \hat{P}_{i0} + \hat{P}_{i1}]$$

The matrix polynomial

$$A(\lambda) = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \lambda^2 + \begin{bmatrix} -3.5 & 0.5 \\ 1.5 & -4.5 \end{bmatrix} \lambda + \begin{bmatrix} 2.5 & -0.5 \\ -3.5 & 5.5 \end{bmatrix}$$

has latent roots  $\lambda_1 = 1$ ,  $\lambda_2 = 3$ ,  $\lambda_3 = 2$  and  $\lambda_4 = 2$  with  $\ell_3 = 1$ . The Jordan block for  $\lambda = 2$  is then

$$J_3 = \begin{bmatrix} 2 & 1 \\ 0 & 2 \end{bmatrix}$$

The latent vectors of  $A(\lambda)$  are

$$\begin{aligned} \hat{y}_1 &= \begin{bmatrix} 1 \\ 1 \end{bmatrix} & \hat{y}_2 &= \begin{bmatrix} 1 \\ -1 \end{bmatrix} & \hat{y}_3^1 &= \begin{bmatrix} 1 \\ 1 \end{bmatrix} & \hat{y}_3^2 &= \begin{bmatrix} 1 \\ -1 \end{bmatrix} \\ \hat{z}_1 &= [1 \ 0] & \hat{z}_2 &= [1 \ -1] & \hat{z}_3^1 &= [1 \ -1] & \hat{z}_3^2 &= [-1 \ -1] \end{aligned}$$

The first solvent is given by

$$R_1 = Q_1 A Q_1^{-1} = \sum_{i=1}^2 \hat{P}_{i0} \lambda_i \left[ \sum_{i=1}^2 \hat{P}_{i0} \right]^{-1} = \begin{bmatrix} 2 & -1 \\ -1 & 2 \end{bmatrix}$$

which gives  $A(R_1) = 0$  as required. The second solvent requires the primary latent projector and the secondary latent projector which are

$$\hat{P}_{30} = \frac{1}{2} \begin{bmatrix} 1 & 1 \\ 3 & -1 \end{bmatrix} = Q_2 \quad \hat{P}_{31} = \frac{1}{2} \begin{bmatrix} -1 & 1 \\ -1 & 1 \end{bmatrix}$$

with  $Q_2 J_2 = \hat{P}_{30} \lambda_3 + \hat{P}_{31}$  or

$$Q_2 J_2 = \frac{1}{2} \begin{bmatrix} 2 & 2 \\ 6 & -2 \end{bmatrix} + \frac{1}{2} \begin{bmatrix} -1 & 1 \\ -1 & 1 \end{bmatrix} = \frac{1}{2} \begin{bmatrix} 1 & 3 \\ 5 & -1 \end{bmatrix}$$

The second solvent is

$$R_2 = Q_2 J_2 Q_2^{-1} = \begin{bmatrix} 1 & 3 \\ 5 & -1 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 3 & -1 \end{bmatrix}^{-1} = \frac{1}{2} \begin{bmatrix} 5 & -1 \\ 1 & 3 \end{bmatrix}$$

which gives  $A(R_2) = 0$  as required.

The extension to higher order polynomials is a simple matter and does not require additional analysis. Each solvent,  $R_1, R_2, \dots, R_m$ , is found by

the procedure given in the preceding work.

The solvents can be determined by the sign algorithm provided that the spectrum of  $A$  has the required distribution. The first step in the procedure is to establish  $A$  for the matrix polynomial and assume that the multiplicity of the repeated eigenvalues are less than  $n$ . Let  $|\lambda_1|, \dots, |\lambda_n|$  have magnitudes less than  $\rho_1$ , and compute the sign of  $A_{D1}$  where  $A_{D1} = (A - \rho_1 I)(A + \rho_1 I)$ . The sign of  $A_{D1}$  will have  $n$  eigenvalues of  $-1$  and  $m-n+1$  eigenvalues. The sign of  $A_{D1}$  can be arranged in the form

$$(5.9) \quad S_1 = \text{Sign}(A_{D1}) = Q \begin{bmatrix} -I_{n \times n} & 0 \\ 0 & I_{m-n+1 \times m-n+1} \end{bmatrix} Q^{-1}$$

by row-column interchanges of the sign as computed by (4.11). Let the eigenvalue matrix (5.9) be denoted by  $J_{I1}$  then

$$(5.10) \quad (S_1 + J_{I1}) = [QJ_{I1} + J_{I1}Q]Q^{-1}$$

Suppose that  $Q$  is partitioned as

$$(5.11) \quad Q = \begin{bmatrix} Q_1 & Q_{12} \\ Q_{21} & Q_{22} \end{bmatrix} = \begin{bmatrix} Q_1 & Q_2 & \dots & Q_m \\ Q_1 \Lambda_1 & Q_2 \Lambda_2 & \dots & Q_m \Lambda_m \\ \vdots & \vdots & \ddots & \vdots \\ Q_1 \Lambda_1^{m-1} & Q_2 \Lambda_2^{m-1} & \dots & Q_m \Lambda_m^{m-1} \end{bmatrix}$$

which when substituted into (5.10) gives

$$(5.12) \quad [S_1 + J_{I1}] = 2 \begin{bmatrix} Q_1 & 0 \\ 0 & Q_{22} \end{bmatrix} Q^{-1}$$

The similarity transformation  $[S_1 + J_1]A[S_1 + J_1]^{-1}$  then gives

$$(5.13) \quad [S_1 + J_{11}]A[S_1 + J_{11}]^{-1} = \begin{bmatrix} Q_1 & 0 \\ 0 & Q_{22} \end{bmatrix} \begin{bmatrix} \Lambda_1 & 0 \\ 0 & \Lambda_{22} \end{bmatrix} \begin{bmatrix} Q_1^{-1} & 0 \\ 0 & Q_{22}^{-1} \end{bmatrix} \\ = \begin{bmatrix} R_1 & 0 \\ 0 & \bar{R}_2 \end{bmatrix}$$

The similarity transformation  $T$  required to reduce  $A$  to the block diagonal form is given by

$$(5.17) \quad T = \frac{1}{(2)^{m-1}} \begin{bmatrix} I & -R_2^{-1} & \dots & (-1)^{m+1} R_m^{-(m+1)} \\ R_1 & -I & \dots & (-1)^{m+1} R_m^{-(m+1)+1} \\ R_1^2 & R_2 & \dots & (-1)^{m+1} R_m^{-(m+1)+2} \\ \vdots & \vdots & \ddots & \vdots \\ \vdots & \vdots & \ddots & \vdots \\ R_1^{m-1} & R_2^{m-2} & \dots & (-1)^{m+1} I \end{bmatrix}$$

where  $R_i$  is a solvent of  $A(\lambda)$ .

The spectral decomposition of  $A$  will now be shown using the second example in this section. Let  $A(\lambda)$  be defined as

$$A(\lambda) = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \lambda^2 + \begin{bmatrix} -3.5 & 0.5 \\ 1.5 & -4.5 \end{bmatrix} \lambda + \begin{bmatrix} 2.5 & -0.5 \\ -3.5 & 5.5 \end{bmatrix}$$

with  $\lambda_1 = 1$ ,  $\lambda_2 = 3$ ,  $\lambda_3 = \lambda_4 = 2$  and  $\ell_3 = 1$ . The companion form  $A$  is

$$A = \begin{bmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ -2.5 & 0.5 & 3.5 & -0.5 \\ 3.5 & -5.5 & -1.5 & 4.5 \end{bmatrix}$$

with solvents

$$R_1 = \begin{bmatrix} 2 & -1 \\ -1 & 2 \end{bmatrix} \quad R_2 = \begin{bmatrix} 2.5 & -0.5 \\ 0.5 & 1.5 \end{bmatrix}$$

From (5.17), T is

$$T = \frac{1}{2} \begin{bmatrix} 1 & 0 & 3/8 & 1/8 \\ 0 & 1 & -1/8 & 5/8 \\ 2 & -1 & -1 & 0 \\ -1 & 2 & 0 & -1 \end{bmatrix}$$

with

$$A_m = A_2 = T A T^{-1} = \begin{bmatrix} 2 & -1 & 0 & 0 \\ -1 & 2 & 0 & 0 \\ 0 & 0 & 2.5 & -0.5 \\ 0 & 0 & 0.5 & 1.5 \end{bmatrix}$$

The characteristic equation for the upper block is  $\lambda^2 - 4\lambda + 3$  and the lower block has  $\lambda^2 - 4\lambda + 4$  as its characteristic equation. The eigenvalues  $\lambda_1 = 1$  and  $\lambda_2 = 3$  are in the upper block with  $\lambda_3 = 2$  and  $\lambda_4 = 2$  in the lower block. The computations for this example were checked by the sign algorithm with a shift of  $-1.8$  and  $\rho_1 = 0.5$ . The two solvents,  $R_1$  and  $R_2$  computed by the sign algorithm, agree with the values given.

## 6. Solution of a System of Differential Equations

The mathematical tools developed in the previous sections will now be applied to the time domain analysis of systems. Assume that the system has been characterized in the first-order form, usually called the state variable form, with states  $x(t)$  such that  $x(t)$  satisfies the differential equation

$$(6.1) \quad \frac{dx(t)}{dt} = Ax(t) + Bu(t)$$

where  $A$  is a general  $m \times m$  matrix,  $B$  is  $m \times k$ ,  $x(t)$  is a  $m \times 1$  vector and  $u(t)$  is  $k \times 1$ . Let  $Z(t)$  denote the outvector with

$$(6.2) \quad Z(t) = Cx(t)$$

where  $Z(t)$  is  $k \times 1$  and  $C$  is  $k \times m$ . The vector  $u(t)$  will be considered as the input to the system or a control vector if the system is a control system. It will be assumed that  $A, B$  and  $C$  are constant matrices which will be referred to as the system triplet.

It will be assumed that the system is stable, all eigenvalues of  $A$  have  $\text{Re}(\lambda_i) < 0$  except for distinct eigenvalues along the  $j\omega$  axis and multiple eigenvalues at the origin,  $\lambda_i = 0$ .

The solution to (6.1) can be expressed as

$$(6.3) \quad x(t) = \Theta(t, t_0)x(t_0) + \int_{t_0}^t \Theta(t, \tau)Bu(\tau)d\tau$$



where  $\Theta(t, t_0)$  denotes\* the state transition matrix, (STM). The state-transition matrix satisfies the differential equation

$$(6.4) \quad \frac{d\Theta(t, t_0)}{dt} = A \Theta(t, t_0) \quad \Theta(t_0, t_0) = I = \Theta(t, t)$$

where  $\Theta(t, t_0)$  is  $m \times m$  and  $I$  is the identity matrix. There are numerous methods of obtaining  $\Theta(t, t_0)$ , several of the methods will be described.

Since  $A$  is a constant matrix, the Laplace transform of (6.4) can be taken with

$$(6.5) \quad L\left[\frac{d}{dt} \Theta(t, 0)\right] = s\Theta(s) - \Theta(0, 0) = s\Theta(s) - I$$

thus the transform of (6.4) is

$$(6.6) \quad \Theta(s) = [sI - A]^{-1} = Q[sI - J]^{-1}Q^{-1}$$

where  $J$  is the Jordan form. Noting that  $s$  in the Laplace domain is equivalent to  $\lambda$  in the eigenvalue domain,  $\Theta(s)$  is equivalent to  $[\lambda I - A]^{-1} = [A(\lambda)]^{-1}$ .

It therefore follows that  $\Theta(s)$  can be expressed as

$$(6.7) \quad \Theta(s) = \sum_{i=1}^q \frac{P_{i0}}{s - s_i} + \sum_{j=1}^{\ell} \frac{P_{ij}}{(s - s_i)^j}$$

The inverse Laplace transform of  $\Theta(s)$  is  $L^{-1}[\Theta(s)]$  or

$$(6.8) \quad \Theta(t, 0) = Q[\exp(Jt)]Q^{-1} = \sum_{i=1}^q \{P_{i0} \exp(s_i t) + \sum_{j=1}^{\ell} \frac{P_{ij} t^j}{j!} \exp(s_i t)\}$$

\*The normal use of  $\Phi(t, t_0)$  as the state-transition matrix will not be made due to the use of  $\phi$  as the modal matrix, [17].

where  $s_i$  is a eigenvalue of  $A$  or a root of the characteristic equation  $\det[sI-A] = 0$ .

The Laplace transform is usually taken with  $t_0 = 0$  rather than on arbitrary value of  $t_0$ . The state transition matrix  $\Theta(t, t_0)$  can be found by using the semigroup property

$$(6.9) \quad \Theta(t, t_0) = \Theta(t, 0)\Theta(0, t_0)$$

where  $\Theta(0, t_0) = [\Theta(t_0, 0)]^{-1}$  if  $t_0 > 0$ . It follows that  $\Theta(t, t_0)$  is given by

$$(6.10) \quad \Theta(t, t_0) = Q\{\exp[J(t-t_0)]\}Q^{-1} = \sum_{i=1}^q P_{i0} \exp[s_i(t-t_0)] \\ + \sum_{j=1}^{\ell_i} \frac{P_{ij}(t-t_0)^j}{j!} \exp[s_i(t-t_0)]$$

The analytical solution to (6.1) can be expressed as

$$(6.11) \quad x(t) = Q \exp[J(t-t_0)]Q^{-1}x(t_0) + Q \int_{t_0}^t \exp[J(t-\tau)]Q^{-1}Bu(\tau) d\tau$$

where  $Q$  is the eigenvector matrix and  $J$  is the Jordan form. If the eigenprojectors  $P_{i0}$  and  $P_{ij}$  are used, then

$$(6.12) \quad x(t) = \sum_{i=1}^q \{P_{i0} \exp[s_i(t-t_0)]\} + \sum_{j=1}^{\ell_i} \frac{P_{ij}(t-t_0)^j}{j!} \exp[s_i(t-t_0)]x(t_0) \\ + \int_{t_0}^t \sum_{i=1}^q \{ \exp[s_i(t-\tau)]P_{i0} + \sum_{j=1}^{\ell_i} \frac{(t-\tau)^j}{j!} \exp[s_i(t-\tau)]P_{ij} \} Bu(\tau) d\tau$$

where the order of the operation under the integral has been reversed for convenience. This reordering is permissible since  $\exp(\cdot)$  is a scalar.

The system defined in (6.1) is said to be controllable if the input  $u(t)$  (or the control) can drive the initial states  $x(t_0)$  to the origin  $x(t) = 0$  (or to an arbitrary value  $x(t) = x_f$ ). The usual test of controllability is to examine the matrix

$$(6.13) \quad Q_c = [B, AB, A^2B, \dots, A^m B]$$

to determine if  $Q_c$  is invertible. The system is controllable if  $\det Q_c \neq 0$  or if  $Q_c$  is invertible. Controllability of a system can also be measured by considering the product  $P_{ij}B$  for  $i = 1, 2, \dots, q$  and  $j = 1, \dots, l_1$  for the eigenprojectors. The system mode  $\exp(\lambda_1 t)$  is not controllable if

$$(6.14) \quad P_{ij}B = 0$$

for the primary and secondary eigenprojectors of the eigenvalue  $\lambda_1$ . This test implies that  $u(t)$  cannot drive the mode  $\exp(\lambda_1 t)$  as the mode is following the natural response rather than a forced response.

The partial fraction expansion method of determining  $\Theta(t, t_0)$  for all  $t$  is not a computationally efficient process and would be used only when the analytical form of  $\Theta(t, t_0)$  is desired. For computational purposes, the state-transition matrix  $\Theta(t, 0)$  can be determined more efficiently by expressing  $\Theta(t, 0)$  in the form

$$(6.15) \quad \Theta(t, 0) = \exp At = I + \sum_{j=1}^{\infty} \frac{A^j t^j}{j!}$$

The series is then used to compute  $\Theta(t, 0)$  for a small value of  $t$  with trun-

cation of the series when the change in  $\Theta(t,0)$  is beyond the accuracy of the digital computer. Assume that the small  $t$  is taken as  $\Delta t$ , with  $\Theta(\Delta t,0)$  determined from (6.15). The semigroup properties is utilized to find  $\Theta(t,0)$  for  $t = k\Delta t$  by the operation

$$(6.16) \quad \Theta(t,0) = [\Theta(\Delta t,0)]^k = \Theta(\Delta t,0) \Theta(t-\Delta t,0)$$

when  $A$  is constant.

The solution vector  $x(t)$  depends upon the integral as well as the state transition matrix. The expression in (6.3) can be given in a recursive form which is more convenient and which has computational advantages. Let (6.3) be written as

$$(6.17) \quad x(t) = \Theta(t,t_0) x(t_0) + \Gamma(t,t_0)$$

where  $\Theta(t,t_0)$  has been computed by means of (6.10) and where  $\Gamma(t,t_0)$  is the integral. The solution vector at  $t+\Delta t$  can be expressed as

$$(6.18) \quad x(t+\Delta t) = \Theta(t+\Delta t,t)x(t) + \Gamma(t+\Delta t,t)$$

Substituting (6.17) into (6.18),

$$(6.19) \quad x(t+\Delta t) = \Theta(t+\Delta t,t)\Theta(t,t_0)x(t_0) + \Gamma(t+\Delta t,t) + \Theta(t+\Delta t,t)\Gamma(t,t_0)$$

or

$$(6.20) \quad x(t+\Delta t) = \Theta(t+\Delta t,t_0)x(t_0) + \Gamma(t+\Delta t,t_0)$$

Knowledge of  $\Gamma(t+\Delta t, t)$  is required along with  $\Theta(t+\Delta t, t)$  or  $\Theta(\Delta t, 0)$ , since  $A$  is constant, to find  $x(t+\Delta t)$ . The vector  $\Gamma(t, t_0)$  satisfies the equation

$$(6.21) \quad \frac{d\Gamma(t, t_0)}{dt} = A \Gamma(t, t_0) + Bu(t) \quad \Gamma(t_0, t_0) = 0$$

and can be found a 4th-order Runge-Kutta algorithm. Since  $\Gamma(t+\Delta t, t)$  is required for (6.20), the value is found from (6.21) by integrating (6.21) from  $t$  to  $t+\Delta t$  for all  $t$ . Equation (6.19) is then used in a recursive manner with  $\Theta(t+\Delta t, t) = \Theta(\Delta t, 0)$  for constant  $A$ . The series given in (6.15) is used to compute  $\Theta(\Delta t, 0)$  which remains constant thereafter.

If  $u(t)$  is slowly varying and  $\Delta t$  is small, a reasonably good approximation may be obtained from the z-transform, [3], [18], of (6.1). The Laplace transform of (6.1) is

$$(6.22) \quad [sI-A]X(s) = Bu(s)$$

where the initial condition  $x(0)$  has been neglected. Equation (6.22) can be written as

$$(6.23) \quad x(s) = [sI-A]^{-1}Bu(s) = G(s) U(s)$$

Taking the z-transform of  $G(s)$  with a zero-order-hold gives

$$(6.24) \quad G(z) = Z\left\{\frac{1-e^{-sT}}{s} [sI-A]^{-1}\right\} B$$

which is found to be

$$(6.25) \quad G(z) = [zI - \exp(AT)]^{-1} [\exp(AT) - I] A^{-1} B$$

where  $T = \Delta t$ . The associated difference equation of (6.1) is obtained from (6.25) with\*

$$(6.26) \quad x(k+1) = [\exp(AT) - I] A^{-1} B U(k) + \exp(AT) x(k)$$

with  $t = kT$ ,  $(k+1)T = t + \Delta t$  and  $x(0)$  the initial condition vector. The validity of (6.26) can be shown by noting that

$$(6.27) \quad \Theta(t, 0) = \exp(At)$$

$$(6.28) \quad \int_0^T \Theta(t, \tau) B \, d\tau = \int_0^T \exp[A(t-\tau)] B \, d\tau \\ = [\exp(AT) - I] A^{-1} B$$

Equation (6.26) is valid provided that the assumption of slowly varying  $u(t)$  and small  $\Delta t$  is not too strict.

The eigenprojectors can also be utilized to separate the solutions into modes if desired. Recalling that  $[sI - A]^{-1}$  is equivalent to  $[\lambda I - A]^{-1}$  and that  $[\lambda I - A]^{-1}$  can be expanded as a partial fraction, (6.24) can be reformulated as

$$(6.29) \quad G(z) = [A(0)]^{-1} + \sum_{i=1}^q \frac{z-1}{s_i} \left\{ \frac{P_{i0}}{z - e^{s_i T}} + \sum_{j=1}^{\ell_i} \frac{P_{ij} T^j}{(z - e^{s_i T})^{j+1}} e^{j s_i T} \right\}$$

which can be written as a series of transfer functions  $G_{ij}$  with

\*The argument of  $x(\cdot)$  and  $u(\cdot)$  is  $(k+1)T$  with  $T$  being dropped for convenience hereafter.

$$(6.30) \quad G(z) = G_{00}(z) + \sum_{i=1}^q [G_{i0}(z) + \sum_{j=1}^{\ell_i} G_{ij}(z)]$$

The vector  $x(z)$  is then given by

$$(6.31) \quad x(z) = G_{00}(z)B U(z) + \sum_{i=1}^q [G_{i0}(z) + \sum_{j=1}^{\ell_i} G_{ij}(z)]B U(z)$$

Equation (6.31) defines a set of transfer functions that can be placed in parallel with each having  $u(z)$  as an input and each block can be implemented separately by a matrix difference equation.

Recalling that the eigenprojectors are defined by the right and left eigenvectors, the eigenprojectors need not be stored in the digital computer. The eigenprojectors are stored and used in (6.29) to construct (6.29) at each  $T$ .

The discussion in this section has been based on  $A$  having a general form. Assume that  $A$  is the companion form of the matrix polynomial  $A(\lambda)$  with

$$(6.32) \quad I \frac{d^m \hat{x}}{dt^m} + A_1 \frac{d^{m-1} \hat{x}}{dt} + \dots + A_m \hat{x} = u(t)$$

The canonical form of  $x(t)$  is the same as given in (6.1) but with

$$(6.33) \quad \dot{\hat{x}}(t) = \begin{bmatrix} \dot{\hat{x}}(t) \\ \hat{x}(t) \\ \vdots \\ \hat{x}^{(m)}(t) \end{bmatrix} = \begin{bmatrix} 0 & I & 0 & \dots & 0 \\ 0 & 0 & I & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ -\Lambda_m & -\Lambda_{m-1} & -\Lambda_{m-2} & \dots & -\Lambda_1 \end{bmatrix} \begin{bmatrix} \hat{x}(t) \\ \hat{x}(t) \\ \vdots \\ \hat{x}^{(m-1)}(t) \end{bmatrix}$$

$$+ \begin{bmatrix} 0 \\ 0 \\ \vdots \\ I \end{bmatrix} u(t) = A x(t) + B u(t)$$

Rather than use (6.33), the solution vector  $\hat{x}(t)$  can be determined directly from (6.32). Taking the Laplace transform of (6.32) gives

$$(6.34) \quad [Is^m + A_1 s^{m-1} + \dots + A_{m-1} s + A_m] \hat{x}(s) = U(s)$$

where the initial conditions have been neglected. Equation (6.34) can be rewritten as

$$(6.35) \quad \hat{X}(s) = [A(s)]^{-1} U(s) = G(s) U(s)$$

which can be expanded into a partial fraction.

The z-transform of (6.35) can now be found provided that the zero-order-hold is included and  $u(t)$  is slowly varying. Letting

$$(6.36) \quad G(s) = \frac{1-e^{-sT}}{s} G(s)$$

gives

$$(6.37) \quad G(z) = Z\left\{\frac{1-e^{-sT}}{s} [A(s)]^{-1}\right\} = Z\left\{\frac{1-e^{-sT}}{s} G(s)\right\}$$

The z-transform of (6.37) is



$$(6.38) \quad G(z) = [A(0)]^{-1} + \sum_{i=1}^q \frac{(z-1)}{s_i} \left\{ \frac{\hat{p}_{i0}}{s_i^T} + \sum_{j=1}^{\ell} \frac{\hat{p}_{ij}^T e^{js_i^T}}{(z-e^{s_i^T})^{j+1}} \right\}$$

from which the difference equation for  $\hat{x}(k+1)$  can be determined.

The expression in (6.38) would be implemented on a digital machine in parallel form rather than as a series representation. As an example, let

$$A(s) = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} s^2 + \begin{bmatrix} 5 & -2 \\ -2 & 5 \end{bmatrix} s + \begin{bmatrix} 7 & -5 \\ -5 & 7 \end{bmatrix}$$

with roots  $\lambda_1 = -1$ ,  $\lambda_2 = -2$ ,  $\lambda_3 = -3$  and  $\lambda_4 = -4$ . The latent projectors are

$$\hat{p}_{10} = \frac{1}{2} \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix} \quad \hat{p}_{20} = \begin{bmatrix} -1 & -1 \\ -1 & -1 \end{bmatrix} \quad \hat{p}_{30} = \frac{1}{2} \begin{bmatrix} 1 & -1 \\ -1 & 1 \end{bmatrix} \quad \hat{p}_{40} = \frac{1}{2} \begin{bmatrix} -1 & 1 \\ 1 & -1 \end{bmatrix}$$

with  $A(0)$  obtained from  $A(s)$ . The transfer function  $G(z)$  is

$$G(z) = G_{00}(z) + G_{10}(z) + G_{20}(z) + G_{30}(z) + G_{40}(z)$$

where

$$G_{00}(z) = \frac{1}{24} \begin{bmatrix} 7 & 5 \\ 5 & 7 \end{bmatrix}$$

$$G_{10}(z) = \frac{z-1}{2(z-e^{-T})} \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix} = \frac{1}{2} \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix} \frac{1-z^{-1}}{1-e^{-T}z^{-1}}$$

$$G_{20}(z) = \frac{z-1}{2(z-e^{-2T})} \begin{bmatrix} -1 & -1 \\ -1 & -1 \end{bmatrix} = \frac{1}{2} \begin{bmatrix} -1 & -1 \\ -1 & -1 \end{bmatrix} \frac{1-z^{-1}}{1-e^{-2T}z^{-1}}$$

$$G_{30}(z) = \frac{z-1}{2(z-e^{-3T})} \begin{bmatrix} 1 & -1 \\ -1 & 1 \end{bmatrix} = \frac{1}{2} \begin{bmatrix} 1 & -1 \\ -1 & 1 \end{bmatrix} \frac{1-z^{-1}}{1-e^{-3T}z^{-1}}$$

$$G_{40}(z) = \frac{z-1}{2(z-e^{-4T})} \begin{bmatrix} -1 & 1 \\ 1 & -1 \end{bmatrix} = \frac{1}{2} \begin{bmatrix} -1 & 1 \\ 1 & -1 \end{bmatrix} \frac{1-z^{-1}}{1-e^{-4T}z^{-1}}$$

The implementation as a difference equation would then be

$$\hat{X}_0(k+1) = \frac{1}{24} \begin{bmatrix} 7 & 5 \\ 5 & 7 \end{bmatrix} u(k+1)$$

$$\hat{X}_1(k+1) = \frac{1}{2} \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix} (u(k)-u(k-1)) + e^{-T} \hat{X}_1(k)$$

$$\hat{X}_2(k+1) = \frac{1}{2} \begin{bmatrix} -1 & -1 \\ -1 & -1 \end{bmatrix} (u(k)-u(k-1)) + e^{-2T} \hat{X}_2(k)$$

$$\hat{X}_3(k+1) = \frac{1}{2} \begin{bmatrix} 1 & -1 \\ -1 & 1 \end{bmatrix} (u(k)-u(k-1)) + e^{-3T} \hat{X}_3(k)$$

$$\hat{X}_4(k+1) = \frac{1}{2} \begin{bmatrix} -1 & 1 \\ 1 & -1 \end{bmatrix} (u(k)-u(k-1)) + e^{-4T} \hat{X}_4(k)$$

with

$$\hat{X}(k+1) = \sum_{k=0}^4 \hat{X}_k(k+1)$$

The parameter  $T$  is the integration step size (or sampling rate) with  $t = kT$ . Different step sizes can be selected for the modes  $\exp(-s_1 T)$  if better accuracy is desired for the higher frequency modes.

The use of the eigenprojectors and the latent projector for solving the state equation of (6.1) has been described. The extension to the solution of  $n$  mth-order differential equations has been given.

## 7. Spectral Decomposition of Differential Equations

The increasing complexity of modern systems generally requires a large number of algebraic operations on system equations and the corresponding solutions to these equations to characterize the dynamics of the systems. It is therefore essential that the analysis of large scale systems be carried out on subsystems or subsets of the equations. A dissertation by K. S. Yoo, [15], used the concept of mode decoupling to analyze an optimal control system. The overhead for the decoupling procedure was moderately high but the total computational task for the analysis was decreased when compared to the usual procedure. Popeeva and Lupas, [16], published several papers describing the procedure, the referenced paper was the first reference to Yoo's work in the open literature. The original idea for decoupling, or order reduction, was probably due to Roberts, [10].

Consider the system equation given in Section 3 with

$$(7.1) \quad \frac{dx(t)}{dt} = Ax(t) + Bu(t)$$

Let  $T$  denote the  $m \times m$  transformation matrix

$$(7.2) \quad T = \frac{1}{(2)^{m-1}} \begin{bmatrix} I & -R_2^{-1} & R_3^{-2} & \dots & (-1)^m R_m^{-m+1} \\ R_1 & -I & R_3^{-1} & \dots & (-1)^m R_m^{-m+2} \\ R_1^2 & R_2 & I & \dots & (-1)^m R_m^{-m+3} \\ \cdot & \cdot & \cdot & \dots & \cdot \\ \cdot & \cdot & \cdot & \dots & \cdot \\ R_1^{m-1} & R_2^{m-2} & R_3^{m-3} & \dots & I \end{bmatrix}$$

where  $I$  is the  $n \times n$  identity matrix and  $R_i$  is the solvent of

$$(7.3) \quad R_1^m + A_1 R_1^{m-1} + \dots + A_{m-1} R_1 + A_m = 0$$

Define the vector  $v(t)$  by the transformation

$$(7.4) \quad v(t) = T^{-1}x(t)$$

or  $x(t) = T v(t)$ . Assume that  $A$  is the companion form for the  $m$ th-order differential equation, it follows that  $v(t)$  must satisfy

$$(7.5) \quad \frac{dv(t)}{dt} = T^{-1}A T v(t) + T^{-1}B u(t)$$

Consider the matrix  $T^{-1}AT$  when  $m = 3$  and  $T$  is as given in (7.2). It follows that

$$(7.6) \quad A_B = T^{-1}AT = \begin{bmatrix} A_{B1} & 0 & 0 \\ 0 & A_{B2} & 0 \\ 0 & 0 & A_{B3} \end{bmatrix} = \begin{bmatrix} R_1 & 0 & 0 \\ 0 & R_2 & 0 \\ 0 & 0 & R_3 \end{bmatrix}$$

or in general for the  $m$ th-order differential equation

$$(7.7) \quad A_B = \text{diag}[R_1, R_2, R_3, \dots, R_m]$$

Recalling that the solvents  $R_k$  were given by

$$(7.8) \quad R_k = Q_k J_k Q_k^{-1} = \left[ \begin{matrix} (j+1)n \\ \vdots \\ i=jn+1 \end{matrix} \right] [\lambda_1 \hat{P}_{10} + \hat{P}_{11}] \left\{ \begin{matrix} (j+1)n \\ \vdots \\ i=jn+1 \end{matrix} \right\} \hat{P}_{10}^{-1}$$

$A_B$  and  $T$  are completely known provided that the latent projectors are known.

The solution to (7.5) is obtained from the  $m$  equations

$$(7.9) \quad \frac{dv_1(t)}{dt} = R_1 v(t) + B_1 u(t)$$

where

$$(7.10) \quad \bar{B} = T^{-1}B = \begin{bmatrix} B_1 \\ B_2 \\ \cdot \\ \cdot \\ B_m \end{bmatrix}$$

with initial conditions  $v_1(t_0)$  obtained from (7.4) with  $x(t_0)$  given. The  $m \times 1$  vector  $x(t)$  as given in (6.33) is

$$(7.11) \quad x(t) = \begin{bmatrix} \hat{x}(t) \\ \dot{\hat{x}}(t) \\ \cdot \\ \cdot \\ \hat{x}^{(m-1)}(t) \end{bmatrix}$$

and  $x(t) = T v(t)$ . The solution to the  $m$ th-order differential equation of (6.32) is

$$(7.12) \quad \hat{x}(t) = \frac{1}{(2)^{m-1}} \{v_1(t) - R_2^{-1}v_2(t) + R_3^{-2}v_3(t) - \dots\}$$

The decoupling procedure described is useful in solving a large set of differential equations as it allows the set to be decoupled into a number of equations. A  $m$ th-order differential equations with  $n \times n$  coefficients can be

solved by generating  $m$  first-order equations with  $n \times n$  coefficients. These equations can be reduced further if desired by decoupling any of the  $m$  first-order to several first order differential equations with coefficients less than  $n \times n$ . The limit to the decoupling procedure will be  $mn$  first-order equations.

As an example of the above let  $A$  be the general matrix

$$A = \begin{bmatrix} -168 & 48 & 279 & -218 & 8 & 50 \\ -136 & 32 & 233 & -180 & 14 & 36 \\ -80 & 16 & 95 & -58 & 8 & 18 \\ -60 & 12 & 75 & -50 & 8 & 14 \\ -40 & 8 & 50 & -32 & 3 & 10 \\ -20 & 4 & 25 & -16 & 2 & 4 \end{bmatrix}$$

which has eigenvalues  $\lambda_1 = 1$ ,  $\lambda_2 = -2$ ,  $\lambda_3 = -5$ ,  $\lambda_4 = -12$ ,  $\lambda_5 = -24$  and  $\lambda_6 = -40$ . This matrix can be reduced to the companion form by the Krylov transformation

$$(7.13) \quad A_c = K A K^{-1}$$

where the structure of  $K$  is given in the Appendix. The associated matrix polynomial  $A(\lambda)$  is

$$A(\lambda) = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \lambda^3 + \begin{bmatrix} 702.051 & -734.624 \\ 591.229 & -618.051 \end{bmatrix} \lambda^2 + \begin{bmatrix} 7688.79 & -6761.64 \\ 6568.29 & -5763.7 \end{bmatrix} \lambda + \begin{bmatrix} 9481.76 & -8521.03 \\ 8095.49 & -7263.08 \end{bmatrix}$$

The solvents for  $A(\lambda)$  are

$$R_1 = \begin{bmatrix} 1 & -2 \\ 3 & -4 \end{bmatrix} \quad \det[\lambda I - R_1] = \lambda^2 + 3\lambda + 2 = (\lambda + 1)(\lambda + 2)$$

$$R_2 = \begin{bmatrix} -15 & 6 \\ -5 & -2 \end{bmatrix} \quad \det[\lambda I - R_2] = \lambda^2 + 17\lambda + 60 = (\lambda + 5)(\lambda + 12)$$

$$R_3 = \begin{bmatrix} 360 & -480 \\ 320 & -424 \end{bmatrix} \quad \det[\lambda I - R_3] = \lambda^2 + 64\lambda + 960 = (\lambda + 24)(\lambda + 40)$$

The system equation  $\dot{x}(t) = Ax(t) + Bu(t)$  can be solved by considering the reduced equations

$$(7.14) \quad \dot{v}_1(t) = R_1 v(t) + B_1 u_1(t)$$

provided that the overhead for computing the Krylov transformation and that for finding the solvents are acceptable as to efficiency and accuracy.



## 8. Summary

The mathematical analysis in this report had two objectives: the first to bring together the mathematical tools for understanding matrix polynomials, with the second of applying these tools to decoupling of system equations. The study of matrix polynomials, or lambda matrices, is justified as vibrating systems are generally defined with second-order matrix polynomials, see [19]. Although there are several books and journal articles on matrix polynomials, a library search did not reveal a single source of the material that is complete. It is essential that the mathematics of matrix polynomials be understood before efficient algorithms for analyzing vibrating systems can be developed. Present algorithms are not capable of handling large space systems--systems with the number of modes greater than 1000.

The report is incomplete in several areas. The algorithm for determining latent projectors from latent vectors for matrix polynomials with repeated roots was not fully developed. Algorithms for efficient computation of latent roots and latent vectors have not been devised. A thorough literature search has not revealed the availability of a computer program for that purpose. In addition, the decoupling scheme was presented but the computational algorithm for that task was not described. Work will continue in these areas.

The damping of large-space structures is an important engineering design task that must be addressed and algorithms must be developed for that purpose. The analysis of matrix polynomials as well as a comprehensive understanding of how damping affects the overall mathematical structure is an integral part of designing large space structures. Modification of matrix polynomials,

which is necessary for the inclusion of damping, was not considered in the report. Studies will begin in that area with the emphasis on second order polynomials.

This report is only a beginning for the several tasks described above. The development of algorithms for analysis, and design of large space structures will be addressed during the next year of work.

## Appendix Krylov Transformation

The Krylov transformation is a useful algorithm that transforms a general  $m \times m$  matrix  $A$  to the companion form. Let  $A_c$  denote the companion form, then

$$(A.1) \quad A_c = K A K^{-1}$$

The Krylov transformation is a similarity transformation that leaves the eigenvalues invariant but changes the eigenvectors of  $A$  from  $Q$  to  $Q_c$ , where

$$(A.2) \quad Q = \begin{bmatrix} Q_{11} & Q_{12} & \dots & Q_{1m} \\ Q_{21} & Q_{22} & \dots & Q_{2m} \\ \vdots & \vdots & \dots & \vdots \\ Q_{m1} & Q_{m2} & \dots & Q_{mm} \end{bmatrix}$$

and

$$(A.3) \quad Q_c = \begin{bmatrix} Q_{11} & Q_{12} & \dots & Q_m \\ Q_{11}\Lambda_1 & Q_{12}\Lambda_2 & \dots & Q_{1m}\Lambda_m \\ \vdots & \vdots & \dots & \vdots \\ Q_{11}\Lambda_1^{m-1} & Q_{12}\Lambda_2^{m-1} & \dots & Q_{1m}\Lambda_m^{m-1} \end{bmatrix}$$

when  $A$  has distinct eigenvalues. The similarity transformation in (A.1) and the structure of (A.2) and (A.3) requires that  $K$  satisfy

$$(A.4) \quad Q_c = KQ$$

Now from (A.2) - (A.4), if  $m = 3$

$$(A.5) \quad \begin{bmatrix} Q_{11} & Q_{12} & Q_{13} \\ Q_{11}\Lambda_1 & Q_{12}\Lambda_2 & Q_{13}\Lambda_3 \\ Q_{11}\Lambda_1^2 & Q_{12}\Lambda_2^2 & Q_{13}\Lambda_3^2 \end{bmatrix} = \begin{bmatrix} K_{11} & K_{12} & K_{13} \\ K_{21} & K_{22} & K_{23} \\ K_{31} & K_{32} & K_{33} \end{bmatrix} \begin{bmatrix} Q_{11} & Q_{12} & Q_{13} \\ Q_{21} & Q_{22} & Q_{23} \\ Q_{31} & Q_{32} & Q_{33} \end{bmatrix}$$

The  $n \times n$  matrix  $Q_{11}\Lambda_1$  is from (A.5)

$$(A.6) \quad Q_{11}\Lambda_1 = K_{21}Q_{11} + K_{22}Q_{21} + K_{23}Q_{31}$$

or

$$(A.7) \quad Q_{11}\Lambda_1 Q_{11}^{-1} = K_{21} + K_{22}Q_{21}Q_{11}^{-1} + K_{23}Q_{31}Q_{11}^{-1} = A_{B1}$$

where  $A_{B1}$  is the matrix that would be in the upper diagonal from the similarity transformation  $T^{-1}AT$  given in Section 7. The matrices  $Q_{21}Q_{11}^{-1}$  and  $Q_{31}Q_{11}^{-1}$  are  $-R_{12}$  and  $R_{13}$  as defined in Section 7 but for the general matrix with

$$(A.8) \quad T = \frac{1}{(2)^2} \begin{bmatrix} I & -R_{12} & R_{13} \\ R_{21} & -I & R_{23} \\ R_{31} & -R_{32} & I \end{bmatrix} = \frac{1}{4} \begin{bmatrix} I & -Q_{12}Q_{11}^{-1} & Q_{13}Q_{11}^{-1} \\ Q_{21}Q_{11}^{-1} & -I & Q_{23}Q_{11}^{-1} \\ Q_{31}Q_{11}^{-1} & -Q_{32}Q_{11}^{-1} & I \end{bmatrix}$$

Analysis of the other equations in (A.5) gives the set of equations

$$(A.9) \quad \begin{aligned} A_{B1} &= K_{21} + K_{22}R_{12} + K_{23}R_{13} \\ R_{12}A_{B2} &= K_{21}R_{12} + K_{22} + K_{23}R_{32} \\ R_{13}A_{B3} &= K_{21}R_{13} + K_{22}R_{23} + K_{23} \end{aligned}$$

$$\begin{aligned}
 (A.10) \quad I &= K_{11} + K_{12}R_{21} + K_{13}R_{31} \\
 R_{12} &= K_{11}R_{12} + K_{12} + K_{13}R_{31} \\
 R_{13} &= K_{11}R_{13} + K_{22}R_{23} + K_{23}
 \end{aligned}$$

$$\begin{aligned}
 (A.11) \quad A_{B1}^2 &= K_{31} + K_{32}R_{21} + K_{33}R_{31} \\
 R_{12}A_{B2}^2 &= K_{31}R_{12} + K_{32} + K_{33}R_{32} \\
 R_{13}A_{B3}^2 &= K_{31}R_{13} + K_{32}R_{23} + K_{33}
 \end{aligned}$$

The Krylov matrix is given from (A.9)-(A.11) as

$$(A.12) \quad \begin{bmatrix} K_{11} & K_{12} & K_{13} \\ K_{21} & K_{22} & K_{23} \\ K_{31} & K_{32} & K_{33} \end{bmatrix} \begin{bmatrix} I & R_{12} & R_{13} \\ R_{21} & I & R_{23} \\ R_{31} & R_{31} & I \end{bmatrix} = \begin{bmatrix} I & R_{12} & R_{13} \\ A_{B1} & R_{12}A_{B2} & R_{13}A_{B3} \\ A_{B1}^2 & R_{12}A_{B2}^2 & R_{13}A_{B3}^2 \end{bmatrix}$$

or

$$(A.13) \quad \begin{bmatrix} K_{11} & K_{12} & K_{13} \\ K_{21} & K_{22} & K_{23} \\ K_{31} & K_{32} & K_{33} \end{bmatrix} = \begin{bmatrix} I & R_{12} & R_{13} \\ A_{B1} & R_{12}A_{B2} & R_{13}A_{B3} \\ A_{B1}^2 & R_{12}A_{B2}^2 & R_{13}A_{B3}^2 \end{bmatrix} \begin{bmatrix} I & R_{12} & R_{13} \\ R_{21} & I & R_{23} \\ R_{31} & R_{32} & I \end{bmatrix}^{-1}$$

$$= \begin{bmatrix} I & 0 & 0 \\ A_{11} & A_{12} & A_{13} \\ A_{11}^2 + A_{12}A_{21} + A_{13}A_{31} & A_{11}A_{12} + A_{12}A_{22} + A_{13}A_{32} & A_{11}A_{13} + A_{12}A_{23} + A_{13}A_{33} \end{bmatrix}$$

$$= \begin{bmatrix} E_R & I \\ E_R & A \\ E_R & A^2 \end{bmatrix} = K$$

where  $E_R = [I \ 0 \ 0]$ . The similarity transformation given in (A.1) is required to show that the Krylov matrix is as given in (A.13).

The general form of (A.13) is

$$(A.14) \quad K = \begin{bmatrix} E_R I \\ E_R A \\ \cdot \\ \cdot \\ E_R A^{m-1} \end{bmatrix}$$

where  $A$  is the general matrix to be reduced to the companion form,  $A_c$ .

The inverse of all  $Q_{ii}$  matrices must exist. Row-column interchanges of  $A$  can be made in most cases to assure the existence of  $A_c$ . The algorithm will not have good accuracy when  $m$  is large particularly when the matrix is stiff; the eigenvalues have a large spread in magnitude.

## References

1. J. E. Dennis, Jr., J. F. Traub and R. P. Weber, The Algebraic Theory of Matrix Polynomials, SIAM J. Num. Anal., Vol. 13, pp. 831-845, (1976).
2. E. D. Denman, Matrix Polynomials, Roots, and Spectral Factors, Appl. Math. and Comp., Vol. 3, pp. 359-368, (1977).
3. L. A. Zadeh and C. A. Desoer, Linear System Theory, McGraw-Hill Book Company, N. Y., (1963).
4. C. G. Cullen, Matrices and Linear Transformations, Addison-Wesley, Reading, Mass., (1966).
5. P. Lancaster, Lambda-Matrices and Vibrating Systems, Pergamon Press, N. Y., (1966).
6. F. V. Atkinson, Multiparameter Eigenvalue Problems, Vol. I, Academic Press, N. Y. (1972).
7. C. T. Chen, Introduction to Linear System Theory, Holt, Rinehart, and Winston, Inc., N. Y., (1970).
8. F. R. Gantmacher, The Theory of Matrices, Chelsea Publishing Co., N. Y., (1959).
9. P. Lancaster and P. N. Webber, Jordan Chains for Lambda Matrices, Lin. Alg. and Appl., Vol. 1, pp. 563-569, (1968).
10. J. D. Roberts, Linear Model Reduction and Solutions of Algebraic Riccati Equations by Use of the Sign Function, Cambridge U. Rept. CUED/B - CONTROL, TR-13 (1971).
11. W. D. Hoskins and D. J. Walton, A Faster Method of Computing the Square Root of a Matrix, IEEE Trans. Auto. Cont. AC-23, pp. 494-495, (1978).
12. R. L. Mattheys, Stability Analysis via the Extended Matrix Sign Function, Proc. IEE.
13. A. N. Beavers, Jr., and E. D. Denman, The Matrix Sign Function and Computations in Systems, Appl. Math. and Comp., Vol. 2, pp. 63-94, (1976).
14. A. Halbersberg and Y. Bar-Ness, Solution of the Discrete Regulator Problem Using the Matrix Sign Function, Electronic Letters, Vol. 14, pp. 286-289, (1978).
15. K. S. Yoo, Development of a Numerical Algorithm for Uncoupling Constant Coefficient State Equations of Control Theory, Dissertation, Dept. of Elec. Eng., U. of Houston, 1974.

16. C. Popeeva and L. Lupas, Decomposition and Reduction of Linear Systems by the Matrix Sign Function, Revue Roumaine des Sciences Tech., Serie Electrotechnique et Energetique, Vol. 20, pp. 567-576, (1975).
17. R. A. Frazer, W. J. Duncan, and A. R. Collar, Elementary Matrices and Some Applications to Dynamics and Differential Equations, Cambridge Press, London, (1938).
18. E. I. Jury, Theory and Application of the Z-transform Method, John Wiley and Sons, N. Y., (1964).
19. E. C. Pestel and F. A. Leckle, Matrix Methods in Elastomechanics, McGraw-Hill Book Co., N. Y., (1963).
20. R. T. Gregory and D. L. Karney, A Collection of Matrices for Testing Computational Algorithms, Wiley-Interscience, N. Y., (1969).