

N O T I C E

THIS DOCUMENT HAS BEEN REPRODUCED FROM
MICROFICHE. ALTHOUGH IT IS RECOGNIZED THAT
CERTAIN PORTIONS ARE ILLEGIBLE, IT IS BEING RELEASED
IN THE INTEREST OF MAKING AVAILABLE AS MUCH
INFORMATION AS POSSIBLE

NASA

Technical Memorandum **80637**

X-Ray Spectrum of Kepler's SNR

(NASA-TM-80637) X-RAY SPECTRUM OF KEPLER'S
SNR (NASA) 11 p HC A02/MF A01 CSCL 03B

N80-17974

G3/93 Unclass
12887

**R. H. Becker, E. A. Boldt,
S. S. Holt, P. J. Serlemitsos,
and N. E. White**

JANUARY 1980

National Aeronautics and
Space Administration

Goddard Space Flight Center
Greenbelt, Maryland 20771



λ-RAY SPECTRUM OF KEPLER'S SNR

R.H. Becker¹, E.A. Boldt, S.S. Holt, P.J. Serlemitsos, and N.E. White¹

Laboratory for High Energy Astrophysics
NASA/Goddard Space Flight Center
Greenbelt, Maryland 20771

ABSTRACT

Observations made with the Solid State Spectrometer (SSS) aboard the Einstein Observatory confirm Kepler's SNR as an X-ray source with an intensity between 1-3 keV of 7.2×10^{-11} ergs/cm²-s. The X-ray spectrum is similar to those of Cas A and Tycho, with strong line emission from the helium-like species of Si, S and Ar. Direct comparisons to Tycho's SNR suggest a distance to Kepler's SNR of ≥ 5 kpc.

¹Also Dept. Physics & Astronomy, Univ. of Maryland

I. INTRODUCTION

Historical SNR are generally known to be X-ray sources. Radio and optical measurements indicate that Kepler's SNR shares many properties in common with other such remnants and therefore is a likely candidate for X-ray emission. Recently two independent groups (Bunner 1979; Tuohy et al. 1979) have reported X-ray emission from the vicinity of Kepler's SNR, but the relatively large field of view of their respective instruments precluded positive identification. In this paper, we report on the results of an observation of Kepler's SNR with the Solid State Spectrometer (SSS) at the focus of the HEAO-2 (Einstein) X-ray telescope and discuss the results in the context of the emission observed from Cas A and Tycho's SNR with the same instrument.

II. EXPERIMENT AND ANALYSIS

For a detailed description of the SSS detector and electronics we refer the reader to Joyce et al. (1978). To summarize, the experiment consists of a cooled Si(Li) detector, which can be placed at the focus of the Einstein X-ray telescope. So located, the SSS has an effective area of $\sim 100 \text{ cm}^2$ between 1-3 keV which drops monotonically to $\sim 10 \text{ cm}^2$ at 4.5 keV. The field of view of the SSS is a ~ 6 arc min diameter circle. The SSS has an energy resolution of $\sim 160 \text{ eV}$ (FWHM).

The spectral analysis procedures have been discussed previously by Holt et al. (1979) and Becker et al. (1979a). In general, theoretical X-ray spectra are folded through the experiment response and compared to the observational data. Model parameters are varied to minimize the chi-square between the data and the model. In this case, we utilized a model which predicts the emission expected from a hot plasma in collisional equilibrium (Raymond and Smith 1977, 1979).

The extent of the radio emission from Kepler's SNR is ~ 3 arc min (Hermann and Dickel 1973), somewhat smaller than the SSS field of view, so that these measurements should reflect the total X-ray emission from Kepler's SNR.

III. RESULTS AND DISCUSSION

The SSS pulse height spectrum of Kepler's SNR is shown in Figure 1. As in the case of other SNR, the spectrum cannot be well represented by a single temperature, but requires at least two thermal components. The best estimate for the kT of the softer component is controlled by the line emission and is 0.54 keV. The data are not capable of restricting the kT of the high temperature component and therefore kT of this component has been arbitrarily fixed at 4 keV similar to values previously reported for Cas A and Tycho (Davison et al. 1976). If we require identical elemental abundances for both components, the best fit values so determined for Mg, Si, S, Ar, and Fe are given in Table 1. The best fit column density was $6 \pm 1 \times 10^{21}$ atoms/cm² (2σ errors). In these fits, the abundances of He, C, O, N, and Ne are all assumed to be solar (see Becker et al. 1979a). The best fit model has been superimposed upon the data in Figure 1. Based on this model, we infer an X-ray intensity of 7.2×10^{-11} ergs/cm²-s between 1-3 keV.

It is hardly surprising that Kepler's SNR is an X-ray source. Nonetheless these measurements are important because they allow a new channel for comparisons among the young remnants of Cas A, Tycho, and Kepler. It is widely believed that Cas A evolved from the explosion of a very massive star (Chevalier 1977) while Tycho's and Kepler's SNR resulted from Type I SN (Minkowski 1966). The SSS has now observed all three of these remnants, with the results for Cas A and Tycho already reported (Becker et al. 1979a, 1979b).

Direct comparisons among Kepler and the other remnants are complicated by the uncertainty in the distance and column density to Kepler's SNR. Tuohy et al. (1979) suggest a distance between 3.5 and 6 kpc based on the brightness of the historical SN. However, Bunner (1979) estimated a hydrogen column density of $<10^{21}$ atoms/cm² from observations of the X-ray spectrum, suggesting Kepler's SNR is closer than previously accepted values. Our

data do not indicate such a low column density. At the other extreme, van den Bergh and Kamper (1977) suggest Kepler's SNR is within the galactic nuclear bulge, ~ 9 kpc distant.

If Kepler's SN is indeed similar to Tycho's SN, we would conclude the correct distance to Kepler's SNR is greater than ~ 5 kpc (assuming Tycho's SNR is ≈ 1.5 kpc distant (Hughes, Thompson, and Calvin 1971)). The angular extent of Kepler's SNR is only $\sim 35\%$ that of Tycho's SNR, while its X-ray intensity is only $\sim 8\%$ of Tycho's SNR (based on HEAO-1 observations of Tycho by Pravdo et al. 1979), thus suggesting Kepler's SNR to be ~ 3 times more distant than Tycho's SNR. At a distance of 5 kpc, the X-ray luminosity of Kepler's SNR between 1-3 keV is 3.3×10^{35} ergs after correcting for the effects of interstellar absorption.

Qualitatively, the spectra of Cas A, Tycho, and Kepler are very similar. All show strong line emission from helium-like species of Si, S, and Ar, but relatively little emission below 4 keV from ions of Mg and Fe. In all three sources, the inferred abundance (relative to solar values) of S is 2-3 times that of Si, while Ar is 4-8 times that of Si. Furthermore, results from observations of MSH11-54 (Clark Tuohy, and Becker 1980) show the same relative excess of S. When taken together, the spectra of SNR's suggest that the galactic abundance ratio of S to Si is 2-3 times that usually estimated for the solar system, with an even greater relative excess existing for Ar. However, these conclusions will remain questionable until a complete non-equilibrium model for X-ray emission is developed. The relative underabundance of Mg in all three remnants may be indicative of the uncertainties in our models. Since even in these young SNR, a substantial amount of interstellar material (presumably containing Mg) has been swept-up, a complete lack of Mg in the SN ejecta still could not explain such low Mg abundances in the X-ray emitting plasma. The model also fails to account for an apparent line feature at 2.9 keV. This energy corresponds to the $1s^2-1s3p$ transition of helium-like S. In the Tycho SNR spectrum, an excess is present at 2.2

keV, the corresponding transition for helium-like Si. Enhancement of these transitions might be indicative of a non-equilibrium situation in which the electron temperature is higher than that indicated by the line spectrum as is probably the case in young SNR (Itoh 1977).

In summary, we can not account in detail for the various line strengths observed in the spectra of young SNR. However, it is clear that qualitatively similar conditions exist in Cas A, Tycho, and Kepler in spite of the vastly different origins between Cas A and the two Type I remnants. Future work will extend these comparisons to older, more evolved, remnants.

We wish to acknowledge B. Smith and J. Raymond for helpful discussions.

REFERENCES

- Becker, R.H., Holt, S.S., Smith, B.W., White, N.E., Boldt, E.A., Mushotzky, R.F., and Serlemitsos, P.J. 1979a, Ap. J. (Letters), 234, L73.
- Becker, R.H., Holt, S.S., Smith, B.W., White, N.E., Boldt, E.A., Mushotzky, R.F., and Serlemitsos, P.J. 1979b, submitted to Ap. J. (Letters).
- Bunner, A.N. 1979, B.A.A.S. 11, 461.
- Chevalier, R.A. 1977, Ann. N.Y. Acad. Sci. 302, 106.
- Clark, D.H., Tuohy, I.R., and Becker, R.H. 1980, in preparation.
- Davison, P.J.N., Culhane, J.L., and Mitchell, R.J. 1976, Ap. J. (Letters) 206, L37.
- Hermann, B.R. and Dickel, J.R. 1973, Ap. J. 78, 879.
- Holt, S.S., White, N.E., Becker, R.H., Boldt, E.A., Mushotzky, R.F., Serlemitsos, P.J., and Smith, B.W. 1979, Ap. J. (Letters), 234, L65.
- Itoh, H. 1977, Publ. Astron. Soc. Japan 29, 813.
- Joyce, R.M., Becker, R.H., Birsa, F.B., Holt, S.S., and Noordzy, M.P. 1978, IEEE Trans. Nuc. Sci. 25, 453.
- Meyer, J.P. 1978, 22nd Liege International Astrophysical Symp.
- Minkowski, R. 1966, A.J. 71, 371.
- Pravdo, S.H., Smith, B.W., Charles, P.A., and Tuohy, I.R. 1979, Ap. J. (Letters), in press.
- Raymond, J.C. and Smith, B.W. 1977, Ap. J. Suppl. 35, 419.
- Raymond, J.C. and Smith, B.W. 1979, in preparation.
- Tuohy, I.R., Nugent, J.J., Garmire, G.P., and Clark, D.H. 1979, Nature 279, 139.
- van den Bergh, S., and Kamper, K.W. 1977, Ap. J. 218, 617.

FIGURE CAPTIONS

Figure 1 - X-ray spectrum of Kepler's SNR as observed by the SSS on HEAO 2. Superimposed upon the data is the expected response of the SSS to a two-component thermal model. The lower second trace is the estimated contribution to the spectrum from emission by H, He, C, O, N, and Ne. The three peaks in the spectrum at 1.85, 2.45, and 3.1 keV are attributable to line emission from Si XIII, S XV, and Ar XVII respectively.

KEPLER X-RAY SPECTRUM HEAO-B5




TABLE 1: ELEMENTAL ABUNDANCES INFERRED FOR KEPLER'S SNR

<u>ELEMENT</u>	<u>ABUNDANCE⁺</u>	<u>RATIO TO SOLAR⁺⁺</u> <u>ABUNDANCE</u>	<u>EQUIVALENT</u> <u>WIDTH (eV)</u>	<u>PHOTONS</u> <u>($\text{cm}^{-2}\text{s}^{-1}$)</u>
Mg	0.8×10^{-5}	$0.2 \pm .2$	---	---
Si	1.3×10^{-4}	$3.7 \pm .2$	1300	.0035
S	1.4×10^{-4}	$8.2 \pm .5$	1300	.0012
Ar	5×10^{-5}	19 ± 5	400	.00013
Ca	1.6×10^{-5}	7 ± 6	---	---
Fe	1×10^{-4}	3 ± 1	---	---

⁺ Strongly dependent on assumed abundances of C, N, O, and Ne. See Becker et al. (1979a).

⁺⁺ Based on estimates of solar abundances by Meyer (1978). Error estimates are 1 σ .

BIBLIOGRAPHIC DATA SHEET

1. Report No. TM 80637	2. Government Accession No.	3. Recipient's Catalog No.	
4. Title and Subtitle X-Ray Spectrum of Kepler's SNR		5. Report Date January 1980	6. Performing Organization Code 661
		8. Performing Organization Report No.	
7. Author(s) R.H. BECKER, E.A. BOLDT, S.S. HOLT, P.J. SERLEMITSOS, N.E. WHITE		9. Performing Organization Name and Address Code 661 Laboratory for High Energy Astrophysics NASA/Goddard Space Flight Center Greenbelt, MD 20771	
12. Sponsoring Agency Name and Address		10. Work Unit No.	11. Contract or Grant No.
		13. Type of Report and Period Covered	
		14. Sponsoring Agency Code	
15. Supplementary Notes <p style="text-align: center;">Accepted for publication in Ap. J. (Letters)</p>			
16. Abstract <p>Observations made with the Solid State Spectrometer (SSS) aboard the Einstein Observatory confirm Kepler's SNR as an X-ray source with an intensity between 1-3 keV of 7.2×10^{-11} ergs/cm²-s. The X-ray spectrum is similar to those of Cas A and Tycho, with strong line emission from the helium-like species of Si, S and Ar. Direct comparisons to Tycho's SNR suggest a distance of Kepler's SNR of ≥ 5 kpc.</p>			
17. Key Words (Selected by Author(s))		18. Distribution Statement	
19. Security Classif. (of this report) U	20. Security Classif. (of this page) U	21. No. of Pages 9	22. Price*