

General Disclaimer

One or more of the Following Statements may affect this Document

- This document has been reproduced from the best copy furnished by the organizational source. It is being released in the interest of making available as much information as possible.
- This document may contain data, which exceeds the sheet parameters. It was furnished in this condition by the organizational source and is the best copy available.
- This document may contain tone-on-tone or color graphs, charts and/or pictures, which have been reproduced in black and white.
- This document is paginated as submitted by the original source.
- Portions of this document are not fully legible due to the historical nature of some of the material. However, it is the best reproduction available from the original submission.

Lockheed Electronics Company, Inc.

A SUBSIDIARY OF
LOCKHEED CORPORATION

1830 NASA Road 1, Houston, Texas 77058
Tel. 713-333-5411

JSC-14910

7.9-10271
CR-160293

"Made available under NASA sponsorship
in the interest of early and wide dis-
semination of Earth Resources Survey
Program information and without liability
for any use made thereof."

Ref: 642-7650
Contract NAS 9-15200

TECHNICAL MEMORANDUM

THE ECOLOGICAL VARIATIONS IN THERMAL INFRARED EMISSIVITY OF VEGETATION

By

G. K. Arp
D. E. Phinney

(E79-10271) THE ECOLOGICAL VARIATIONS IN
THERMAL INFRARED EMISSIVITY OF VEGETATION
(Lockheed Electronics Co.) 27 p
HC A03/MF A01

N79-32605

CSCL 08F

Unclas

G3/43 00271

Approved By:

B. L. Carroll
B. L. Carroll, Manager
EO Development and
Evaluation Department

Distribution
LEC/Technical Library (5)

This document is a revision of and supersedes THE ECOLOGICAL SIGNIFICANCE
OF EMISSIVITY (LEC-11021 Revision B).

April 1979



| | | | |
|---|--|--|------------|
| 1. Report No. JSC-14910 | 2. Government Accession No. | 3. Recipient's Catalog No. | |
| 4. Title and Subtitle The Ecological Variations in Thermal Infrared Emissivity of Vegetation | | 5. Report Date April 1979 | |
| | | 6. Performing Organization Code LEC-13390 | |
| 7. Author(s) G. K. Arp and D. E. Phinney | | 8. Performing Organization Report No. LEC-13390 | |
| | | 10. Work Unit No. | |
| 9. Performing Organization Name and Address Lockheed Electronics Company, Inc. 1830 NASA Road 1 Houston, Texas 77058 | | 11. Contract or Grant No. NAS 9-15200 | |
| | | 13. Type of Report and Period Covered | |
| 12. Sponsoring Agency Name and Address National Aeronautics and Space Administration Lyndon B. Johnson Space Center Houston, Texas 77058 Technical Monitor: J. D. Erickson/SF3 | | 14. Sponsoring Agency Code | |
| | | 16. Supplementary Notes | |
| 16. Abstract Thermal emissivity measurements in the 10.5 μm to 12.5 μm spectral region were taken for a variety of native common and dominant plants of southern U.S. and Mexico. Results of these measurements are reported with a statistical analysis that suggests there is a significant difference between the emissivity and hence, the thermal properties of plants from desert, tropical, and temperate regions. A discussion of the significance and interpretation of these results is presented. | | | |
| 17. Key Words (Suggested by Author(s)) Emissivity Desert and mesic plants Radiation | | 18. Distribution Statement | |
| 19. Security Classif. (of this report) Unclassified | 20. Security Classif. (of this page) Unclassified | 21. No. of Pages 27 | 22. Price* |

*For sale by the National Technical Information Service, Springfield, Virginia 22161

CONTENTS

| Section | Page |
|--|------|
| 1. INTRODUCTION | 1-1 |
| 1.1 <u>DEFINITION OF EMISSIVITY</u> | 1-1 |
| 1.2 <u>THEORY OF MEASUREMENT OF SURFACE EMISSIVITY</u> | 1-4 |
| 2. BACKGROUND | 2-1 |
| 3. METHODS AND MATERIALS | 3-1 |
| 4. DISCUSSION | 4-1 |
| 4.1 <u>ANALYSIS OF DATA</u> | 4-1 |
| 4.2 <u>INTERPRETATION OF RESULTS.</u> | 4-1 |
| 4.3 <u>CONCLUSIONS.</u> | 4-3 |
| 5. REFERENCES | 5-1 |

PROCEEDINGS SHOULD BE MARKED NOT FILED

TABLES

| Table | | Page |
|-------|---|------|
| 3-1 | EMISSIVITY VALUES FOR VARIOUS MEXICAN AND SOUTHWESTERN U.S. PLANTS | 3-4 |
| 4-1 | MAJOR ECOLOGICAL GROUPS EXAMINED FOR VARIATION IN EMISSIVITY | 4-2 |
| 4-2 | PRINCIPAL CONTRASTS | 4-2 |
| 4-3 | MEANS AND STANDARD DEVIATIONS OF EMISSIVITY FOR THREE VEGETATIVE GROUPS | 4-2 |

FIGURES

| Figure | | Page |
|--------|---|------|
| 1-1 | Temperature error ($^{\circ}\text{C}$) associated with an incorrect assumption of emissivity at 300°K | 1-3 |
| 1-2 | Simplified longwave radiative balance at the earth's surface | 1-5 |

1. INTRODUCTION

The representative thermal emissivity measurements of common and conspicuous plants of the southwestern U.S. and Mexico were part of a project sponsored by the National Aeronautics and Space Administration (NASA) Health Applications Office from 1973 to 1975 for the screwworm eradication project. These data, soil data, and ancillary information were used to compute factors for use in the correction of thermal infrared temperature measurements made by the National Oceanic and Atmospheric Administration (NOAA) satellites. The results of this work were used to obtain accurate ground air temperature estimates twice daily over Mexico and the southwestern U.S. These estimates were applied to the screwworm eradication base and were used in the prediction of the screwworm fly infestation sites. Refer to Barnes and Forsberg⁽¹⁾ for details of this project.

A review of the emissivity values of this project suggests that there are significant differences between values obtained for desert plants and other ecological species. This significance supports preliminary work by Gates⁽¹⁰⁾, who suggested that in very dry areas, plants might alleviate some of their potential heat absorption by efficiently emitting energy in the thermal infrared regions. This feature is especially important in desert regions where large amounts of heat and light are present but mechanisms for heat dispersal are limited due to restricted availability of water needed for cooling in evapotranspiration. To determine the difference in emissivity values between desert plants and other ecological plants, a series of statistical tests was performed on the collected data. In this report, a discussion of emissivity including background is presented prior to documenting the procedures and significant findings of these tests.

1.1 DEFINITION OF EMISSIVITY

The spectral emissivity, ϵ , of a homogeneous surface is defined by Huschke⁽¹³⁾ as the ratio of the radiance of the surface at a specified wavelength and emitting temperature to the radiance of an ideal blackbody at the same wavelength and temperature. The values for emissivity may range from zero to unity.

Planck's law gives the spectral distribution of the radiance from a perfect radiator (blackbody) at temperature T as:

$$B_{\lambda} = C_1 \lambda^{-5} [\text{EXP}(C_2/\lambda T) - 1]^{-1} \quad (1)$$

where

$$C_1 = 3.75 \times 10^{-16} \text{ Wm}^2$$

$$C_2 = 1.44 \times 10^{-2} \text{ m}^{\circ}\text{K}$$

λ = wavelength in meters

T = absolute temperature in degrees Kelvin

The spectral radiance emitted by an opaque gray-body may then be written:

$$L_{\lambda}(T) = \epsilon(\lambda) B_{\lambda}(T) \quad (2)$$

Thus, if the actual emissivity of a surface is not considered, the temperature calculated from radiometric data will be lower than the true surface temperature.

For naturally occurring surfaces, emissivity values in the thermal infrared wavelengths have been reported ranging from 0.82 for granite to near 1.0 for water, Buettner and Kern⁽⁴⁾. Most surfaces seem to fall within this range. Generally, rock ranges from 0.86 to 0.93, Buettner and Kern⁽⁴⁾; soil ranges from 0.90 to 0.97, varying with type and moisture content, Fuchs and Tanner⁽⁸⁾. Most vegetative surfaces lie between 0.96 and 0.98.

Equations (1) and (2) may be used to evaluate the magnitude of the error associated with using an incorrect value for emissivity. Figure 1-1 presents this error for the 10.5 μm to 12.5 μm spectral band which corresponds to the spectral sensitivity of the radiometers carried by the NOAA satellites. The data for this figure was developed for a 300° K surface. Estimated surface temperatures were calculated by numerically inverting equation (1) to satisfy the following relationship:

$$\frac{\hat{\epsilon}}{\epsilon} B_{\lambda}(T) = B_{\lambda}(\hat{T}) \quad (3)$$




Figure 1-1.— Temperature error ($^{\circ}\text{C}$) associated with an incorrect assumption of emissivity at 300°K .

#

where the true emissivity ϵ and the estimated emissivity $\hat{\epsilon}$ were varied between 0.9 and 1.0. For simplicity, the emissivity was assumed constant over the spectral region. While this assumption is not strictly valid, particularly for siliceous minerals, an average emissivity can generally be used without serious error in the thermal infrared region.

As can be seen from Figure 1-1, a 0.01 error in emissivity will result in an approximately 0.7° C temperature error. The increasingly sophisticated uses being made of radiometric data can no longer allow errors of several degrees simply due to lack of adequate information on surface emissivity.

1.2 THEORY OF MEASUREMENT OF SURFACE EMISSIVITY

In this report, the radiation terminology proposed by the World Meteorological Organization⁽¹⁶⁾ is used and all radiances are for the entire infrared spectrum.

Consider the longwave radiative balance at the earth's surface which is shown schematically in Figure 1-2. The outgoing spectral radiance, L_{\uparrow} , consists of two parts. The largest part, $\epsilon_s L_b$, is emitted by the surface; the remainder is the portion of the incoming longwave radiation, L_{\downarrow} , that is reflected by the surface. Thus, the radiative balance at the surface may be written:

$$L_{\uparrow} = \epsilon_s L_b + r_s L_{\downarrow} \quad (4)$$

where r_s , the longwave reflectivity, equals $1 - \epsilon_s$. Solving equation (4) for emissivity yields the following equation:

$$\epsilon_s = \frac{L_{\uparrow} - L_{\downarrow}}{L_b - L_{\downarrow}} \quad (5)$$

Thus, to calculate the infrared emissivity of a surface L_{\uparrow} , L_{\downarrow} and L_b must be measured.

In practice, only a portion of the longwave radiance is measured as determined by the spectral sensitivity of the radiometer used. Therefore, care is necessary when comparing emissivities measured with instruments of differing spectral sensitivities. An analysis of the sensitivity of the calculated




Figure 1-2.— Simplified longwave radiative balance at the earth's surface.

emissivity to measurement errors in the component radiances has been carried out by Davies *et al.*⁽⁶⁾ It was shown that for typical conditions, the sensitivity of ϵ_s to errors in $L\uparrow$ was 0.0001 per °K. (The radiance is expressed in terms of equivalent blackbody temperature.) Equivalent sensitivities for L_b and $L\uparrow$ were -0.026 and -0.028 per °K, respectively. However, under conditions of extremely warm sky, the values for $L\uparrow$ become significant. Thus, equal care should be taken with all measurements.

2. BACKGROUND

Infrared emissivities have been measured experimentally by a number of researchers using a variety of field and laboratory instrumentation. Before the field trip for plant specimens and emissivity measurements was made, literature on the measurement of infrared emissivities was examined with attention directed toward techniques, instrumentation, and results.

In 1923, Falckenberg⁽⁷⁾ measured emissivities at 10µm using a single beam spectrophotometer which ranged from 0.89 for sand to 0.955 for snow.

The first evidence of a systematic variation in components of the radiative heat balance with a change in ecological communities was reported by Billings and Morris⁽³⁾. The authors present visible reflectance values for five Great Basin communities ranging from hot desert to cool moist subalpine forests. Their data suggest that communities with hotter, dryer conditions have visible reflectance values higher than values obtained from communities with cooler, wetter conditions. Their results involve the need for corresponding information about components of the infrared energy balance.

Gates and Tantraporn⁽⁹⁾ measured the reflectivity of numerous deciduous trees and shrubs using a double-beam Baird spectrophotometer. A systematic variation in emissivity for some species was noted with the upper surface of the leaf higher than the lower, the shade leaf more than the sun, and old leaves more than new. It can be noted in their data that many plants from dry areas had a relatively higher emissivity. This phenomenon was attributed to the presence of a layer of waxy cuticle on the leaf surface.

The work of Buettner and Kern⁽⁴⁾ represents a milestone in the measurement of surface emissivities. This and most subsequent work made use of portable infrared radiometers developed by Barnes Engineering Company of Stamford, Connecticut. The technique developed by Buettner and Kern is fairly cumbersome and is more suited to the laboratory than the field. However, the results

from their numerous measurements are of high quality and represent a basic reference for the emissivity of a number of minerals.

Buettner and Kern's approach was to create a controlled environment through the use of an emissivity "box". A box with highly reflective sides was constructed such that the top could alternately be a high reflective surface or a temperature controlled pseudo-blackbody. When the highly reflective top was in place, a blackbody cavity (hohlraum) was simulated, and the spectral radiance emitted by the surface was measured through a hole in the top. The high emittance top was maintained at a temperature well below ambient. When the radiance was measured with this top in place, the resultant was the sum of the surface emittance and the reflected portion of the downwelling radiance from the top. If the temperature, the emissivity of the high emittance top, and the radiance of the surface are known when in the hohlraum, the emissivity were easily calculated.

Buettner and Kern used an IT-2 infrared radiometer with a spectral sensitivity from 8 to 12 μ . A number of their measurements compared favorably with the integrated readings from a Beckman IR-8 spectrophotometer.

Lorenz⁽¹⁵⁾ studied several surfaces yielding generally good results. However, his results were somewhat erratic probably due to his method of measuring sky radiation. Using an IT-1 infrared radiometer (8 to 14 μ), Lorenz measured the surface emittance using an aluminum lined box for a hohlraum. He then measured the combined surface emittance and reflected sky radiation directly by placing the surface under an open sky. The sky radiation was then estimated by integrating several direct readings of sky temperature made at different zenith angles.

Fuchs and Tanner⁽⁸⁾ developed their own method of measurement and report experimental data for a few agricultural crops as well as for bare soils. This technique involves using a reference target of known temperature and emissivity to estimate downwelling radiation from the sky. Fuchs and Tanner used an IT-2 and an IT-3 radiometer sensitive to the 8 to 13 μ spectral band.

Fuchs and Tanner⁽⁹⁾ presented measurements on sand and illustrated the dependence of emissivity on moisture content. Using a sandy soil, Fuchs and Tanner observed variations from 0.90 with 0.7 percent water to 0.94 with 8.4 percent water. At that time they also raised a question as to the relative validity of measurements made with the techniques of Buettner and Kern. Idso and Jackson⁽¹⁴⁾ experimentally examined the rival methods and found them to be equivalent in accuracy with root mean square errors ranging from 0.003 to 0.008.

In contrast with Fuchs and Tanner's work, Hovis⁽¹²⁾ reported emissivities of clay and loam soils close to 0.96 with no apparent variation due to soil moisture.

Conaway and Van Bavel⁽⁵⁾ reported additional measurements on bare soil using the Buettner and Kern technique. This study examined the use of radiometrically determined surface temperature in calculating evaporation from bare soils.

Davies *et al.*⁽⁶⁾ conducted additional measurements of the emissivity of water using a Barnes PRT-5 (8 - 14 μ) infrared radiometer. They report a value of 0.972 with no detectable variation due to turbidity. This compares poorly to Buettner and Kern's value of 0.993, perhaps due to the differing spectral sensitivities of the instruments used in the two studies.

Bartholic *et al.*⁽²⁾ measured the emissivity of cotton and bare soil in the course of a study to determine the use of thermal infrared in delineating moisture stress and soil moisture conditions.

In general, all of the workers who have developed the techniques for measuring emissivity seem to have reported on a fairly random selection of whatever material was on hand. As a result, persons working on applications which require a knowledge of surface emissivities have been forced to take their own measurements. In addition, with the exception of work by Billings and Morris⁽³⁾ and Gates *et al.*^(10 and 11), little effort has been made to study systematically the collective emissivities of species which occur together in a given ecological situation.

3. METHODS AND MATERIALS

To gather the desired emissivity data, a series of trips were made to eastern and northern Mexico, Texas, New Mexico, and Arizona. Field measurements were made of the important dominant species of each area. The choice of species included only those that formed the exposed overstory in each community as only their radiational surfaces would contribute significantly to the scene emissivity as perceived by the NOAA satellite.

When first entering a study area, the scientist determined the kind and number of dominant plant communities. The use of botanical literature and available aerial photographs greatly simplified the problems associated with determining the distribution of the key Mexican and U.S. communities. After a general survey, representative communities were selected for detailed analyses. Quadrats were used to determine plant cover and dominance. At each site, representative localities were chosen, quadrats 50 m on a side were marked off, and the vegetation measured and mapped. Based upon a plant's relative occurrence within a community, the conspicuously dominant, common, and occasional plants were listed for each community.

After an area had been surveyed and the candidates for measurements were known, the instruments were set up in a clear area with no overhead trees or other radiational obstructions in the immediate area. The measurement site was away from cars and the accessory instruments to ensure that radiation from cars, people, and accessory instruments did not affect the field measurements. While the instruments were being assembled and warmed up, specimens, representing all desired material, were gathered quickly. Time is a critical factor in all phases of emissivity measurements because temperatures and sky radiation can fluctuate rapidly within a few minutes and specimens can wilt, often quickly.

Only leafy branch tips from the exposed upper surfaces of the plant were clipped for emissivity measurements. Branches from the lateral but exposed portions are best because the leaf orientation with respect to the sun will

11

remain approximately the same when measured by the radiometer. Several branches 8 to 10 inches long were selected and laid one upon the other with upper leaf surfaces facing upward and correctly aligned so that the leaf sample orientation was as nearly normal as possible. Careful attention made sure that enough layers of leaves were placed together so that none of the underlying surface showed through. Usually, a dense bundle of leaves 10 inches in diameter was created. If botanical reference specimens were needed, it was useful and sometimes critical to collect samples of fruit, flowers, and/or seeds for use in species identification. These latter portions were not included in the emissivity sample unless they formed a conspicuous portion of the canopy. A bundle for each species was made and laid in order of collection number on the ground at the measurement site. For woody species or those that habitually show dead or bare branches in the canopy, bare twigs were included in the bundle. For such cases, it was frequently difficult to create a representative mass of vegetation and branches.

Each specimen was given a collection number and reference name (or botanical name if known). In the field notebook, the collection number was recorded with data on local distribution and relative dominance. Once the collection numbers were assigned, measurements were made, alleviating the potential for wilting which can be a serious problem in dry or windy areas. Afterwards, further notes were recorded. Upon completion of the critical measurements, portions of each specimen sample were placed in the plant press as needed for later use in specimen identification and verification. If fruits, flowers, or seeds were previously collected, these were included plus enough vegetative material to make two herbarium sheets of voucher material.

The field readings were screened on the spot with complete reduction occurring at a later date in the laboratory. Generally, it is a good practice to evaluate at least a part of the data in the field to eliminate spurious readings.

The calibration curves for the Barnes PRT-5 were used to convert the readings from the digital voltmeter to temperature. The temperatures were then converted to radiances.

The magnitude of the downwelling sky radiation was calculated from the measurements made upon the reference target. Using equation (4) and solving for L_{\downarrow} yields:

$$L_{\downarrow} = \frac{L_{\uparrow} - \epsilon_r L_b}{1 - \epsilon_r} \quad (6)$$

Thus, by measuring L directly over the reference's target and measuring L_b using the emissivity box, L_{\downarrow} can be calculated when the emissivity of the reference target, ϵ_r , is known.

The emissivity of the surface (over the spectral range of the radiometer) may then be calculated. After examining equation (5), it is seen that the thermal infrared emissivity can be calculated directly from L_{\downarrow} and the measurements of L_b and L_{\uparrow} taken over the unknown. The emissivity values for various Mexican and southwestern U.S. plants that were measured as part of this study are presented in Table 3-1.

The equipment used for conducting these measurements consisted of a modified Barnes PRT-5 with spectral sensitivity from 10.5 μ m to 12.5 μ m and a digital voltmeter for the radiometric measurements, an aluminum lined emissivity box for measurement of surface radiance, and a brass reference target used to calculate downwelling sky radiation. All radiance values used hereafter are for the 10.5 μ m to 12.5 μ m spectral region.

The measurement sequence for each surface is conducted as follows:

1. Measure L_{\uparrow} of the reference target
2. Measure L_s of the reference target using the emissivity box
3. Measure L_{\uparrow} of the unknown
4. Measure L_s of the unknown

The ideal conditions for measurement are low winds with a cold clear sky. Often, early morning and late afternoon produce the best results as the changes

TABLE 3-1.— EMISSIVITY VALUES FOR VARIOUS MEXICAN AND SOUTHWESTERN U.S. PLANTS

| Habitat/niche | Measurement site | Botanical name | Common name | Collection number | Emissivity | Native or cultivated | Date | Number of replications |
|--------------------------------|--------------------------------|--|-------------|-------------------|--------------|----------------------|-------|------------------------|
| Chaparral | | | | | | | | |
| Chaparral component | Marathon, TX | <i>Acacia constricta</i> Gray | Acacia | 4633 | .974 | Native | 10/75 | 3 |
| Chaparral component | Chinati Mountains, TX | <i>Acacia neovevnicosa</i> Jasty | Acacia | 4618 | .982 | Native | 10/75 | 2 |
| Ground cover; invader | Laredo, TX | <i>Boraginaceae</i> family | | 4283 | .991 | Native | 11/74 | 1 |
| Introduced range grass | Laredo/Del Rio, TX | <i>Cenchrus ciliaris</i> L. | Buffelgrass | 4286 | .976 | Cultivated | 11/74 | 2 |
| Chaparral | Chinati Mountains, TX | <i>Condalia viridis</i> L. M. Johnston | | 4619 | .963 | Native | 10/75 | 3 |
| Chaparral component | Starr County, TX | <i>Hollettia parvifolia</i> (Gray) Benth. | Barreta | 4260 | .987 | Native | 11/74 | 1 |
| Chaparral component | Starr County, TX | <i>Karwinskia humboldtiana</i> (R.S.) Zucc. | Coyotillo | 4261 | .945 | Native | 11/74 | 1 |
| Chaparral component | Starr County, TX | <i>Leucophyllum frutescens</i> (Berl.) I.M. Johnston | Chenizo | 4258 | .958 | Native | 11/74 | 1 |
| Chaparral component | Laredo, TX | <i>Leucophyllum frutescens</i> (Berg.) I.M. Johnston | Chenizo | 4287 | .989 .984 | Native | 11/74 | 2 |
| Chaparral component | Laredo, TX | <i>Portulaca angustifolia</i> (Engelm.) Gray | Guayacán | 4288 | .950 | Native | 11/74 | 2 |
| Dominant shrub | Laredo, TX | <i>Prosopis glandulosa</i> Torr. | Mesquite | 4284 | .987 | Native | 11/74 | 2 |
| Dominant tall brush | Lower Valley, TX | <i>Prosopis glandulosa</i> Torr. | Mesquite | 5K | .988 | Native | 11/74 | 1 |
| Cloud forest | | | | | | | | |
| High elevation shrub | Mirador near Esperanza, Puebla | <i>Baccharis confusa</i> H.B.K. | Encino | 4528 | .978 | Native | 1/75 | 2 |
| High elevation mesophytic pine | Mirador near Esperanza, Puebla | <i>Pinus latophylla</i> Schlecht. and Cham. | Pine | 4529 | .958 | Native | 1/75 | 2 |
| Temperate tree | Coscomatepec, Veracruz | <i>Platanus lindeniana</i> Mart. and Gal. | Sycamore | 4548 | .966 | Native | 1/75 | 3 |
| High elevation tree | Mirador near Esperanza, Puebla | <i>Quercus condiciana</i> Nee | Encino | 4531 | .969 | Native | 1/75 | 3 |
| High elevation tree | Mirador near Esperanza, Puebla | <i>Quercus crassifolia</i> Humb. and Bonpl. | Encino | 4530 | .973 | Native | 1/75 | 2 |
| High elevation shrub | Mirador near Esperanza, Puebla | <i>Solanum corvanticum</i> Lag. | Nightshade | 4532 | .958 | Native | 1/75 | 4 |
| Desert | | | | | | | | |
| Desert shrub | Laredo, TX | <i>Acacia farnesiana</i> (L.) Willd. | Huisache | 4282 | .989 | Native | 11/74 | 1 |
| Common in rosette form deserts | Hot Springs, TX | <i>Agave lecheguilla</i> Torr. | Lecheguilla | 4606 | .997 | Native | 10/75 | 3 |
| Desert grassland | Marathon, TX | <i>Agrostis</i> sp. | | 4626 | .961 | Native | 10/75 | 3 |
| Common in washes | Ft. Stockton, TX | <i>Aloynia graveolens</i> (Gill. & Hook.) Troncoso. | White brush | 4311 | .988 | Native | 11/74 | 2 |

REPRODUCED FROM THE
 NATIONAL ARCHIVES
 COLLEGE PARK, MARYLAND
 Page 13
 OF 13

TABLE 3-1.— Continued.

| Habitat/niche | Measurement site | Botanical name | Common name | Collection number | Emissivity | Native or cultivated | Date | Number of replications |
|--------------------------|----------------------|--|---------------------|-------------------|---------------|----------------------|-------|------------------------|
| Desert | | | | | | | | |
| Desert grass | Laredo, TX | <i>Arctostida glauca</i> (Nees.) Walp. | Three-awn | 4285 | 0.983 .982 | Native | 11/74 | 2 |
| Limestone hills | Sanderson Canyon, TX | <i>Arctostida</i> sp. | Three-awn | 4655 | .972 | Native | 10/75 | 4 |
| Winter annual understory | Ft. Stockton, TX | <i>Antropogon</i> sp. | Hoik-vetch | 4323 | .993 | Native | 11/74 | 3 |
| Alkali desert shrub | Marathon, TX | <i>Atriplex canescens</i> (Pursh) Nutt. | Four-wing salt bush | 4628 | .966 | Native | 10/75 | 3 |
| Desert grass-land | Marathon, TX | <i>Bouteloua curtipendula</i> (Michx.) Torr. | Side-oats grama | 4630 | .987 | Native | 10/75 | 4 |
| Desert grass | Van Horn, TX | <i>Bouteloua eriopoda</i> (Torr.) Torr. | Black grama | 4327 | .990 | Native | 11/74 | 2 |
| Desert grass-land | Marathon, TX | <i>Bouteloua hirsuta</i> Lag. Hairy Grama | Gramma | 4627 | .969 | Native | 10/75 | 3 |
| Ground cover | Ft. Stockton, TX | <i>Boraginaceae</i> family | | 4259 | .988 | Native | 11/74 | 1 |
| Rosette form desert | Hot Springs, TX | <i>Boraginaceae</i> family | | 4601 | .983 | Native | 10/75 | 2 |
| Desert grass | Hot Springs, TX | <i>Budloe distyloides</i> (Nutt.) Engelm. | Buffalo grass | 4604 | .978 | Native | 10/75 | 3 |
| Creosote bush hills | Plata, TX | <i>Chrysothamnus</i> sp. | Rabbit-brush | 4610 | .985 | Native | 10/75 | 4 |
| Rosette form desert | Hot Springs, TX | <i>Compositae</i> | | 4603 | .975 | Native | 10/75 | 3 |
| Desert shrub | Van Horn, TX | <i>Condalia arborescens</i> (Gray) M.C. Johnson | Havelina bush | 4326 | .988 | Native | 11/74 | 2 |
| Roadside weed | Chiatl Mountains, TX | <i>Croton Fattali</i> (Kt.) Muell. Arg. | Leather weed | 4620 | .955 | Native | 10/75 | 3 |
| Desert grass | Van Horn, TX | <i>Friesea pulchellum</i> (H.B.K.) Tateoka | Fluffgrass | 4331 | .978 | Native | 11/74 | 2 |
| Desert cactus | Tehuacan, Puebla | <i>Echinocactus chiotilla</i> (Web.) Rose | Chiotilla | 4534 | .960 | Native | 1/75 | 2 |
| Desert shrub | Ft. Stockton, TX | <i>Flourensia cernua</i> D.C. | Tarbush | 4312 | .993 | Native | 11/74 | 2 |
| Desert shrub | Van Horn, TX | <i>Flourensia cernua</i> D.C. | Tarbush | 4328 | .993 | Native | 11/74 | 2 |
| Creosote bush | Plata, TX | <i>Flourensia cernua</i> D.C. | Tarbush | 4609 | .993 | Native | 10/75 | 3 |
| Desert grass | Ft. Stockton, TX | <i>Hilaria mutica</i> (Buckl.) Benth. | Tobosa | 4310 | .982 | Native | 11/74 | 3 |
| Desert mountain | Ft. Stockton, TX | <i>Juniperus ashei</i> Buchh. | Rock cedar | 4304 | .993 | Native | 11/74 | 2 |
| Limestone desert | Sanderson Canyon, TX | <i>Juniperus deppeana</i> Steud. | Alligator juniper | 4658 | .994 | Native | 10/75 | 5 |
| Desert shrub | Ft. Stockton, TX | <i>Koeberlinia spinosa</i> Zucc. | Allthorn | 4317 | .982 | Native | 11/74 | 2 |
| Desert grass-land | Marathon, TX | <i>Labiatae</i> | | 4632 | .986 | Native | 10/75 | 2 |
| Desert shrub | Ft. Stockton, TX | <i>Larrea tridentata</i> (D.C.) Cov. | Creosote bush | 4313 | .981 | Native | 11/74 | 3 |
| Desert shrub | Van Horn, TX | <i>Larrea tridentata</i> (D.C.) Cov. | Creosote bush | 4325 | .981 | Native | 11/74 | 3 |

TABLE 3-1.-- Continued.

| Habitat/niche | Measurement site | Botanical name | Common name | Collection number | Emissivity | Native or cultivated | Date | Number of replications |
|---|---------------------------------|--|---------------------|-------------------|------------|----------------------|-------|------------------------|
| Desert | | | | | | | | |
| Desert shrub | Hot Springs, TX | <i>Larrea tridentata</i> (D.C.) Cov. | Creosote bush | 4602 | 0.986 | Native | 10/75 | 2 |
| Desert shrub | Plata, TX | <i>Larrea tridentata</i> (D.C.) Cov. | Creosote bush | 4608 | .995 | Native | 10/75 | 3 |
| Hot desert cactus | Tehuacan Valley, Puebla | <i>Leontodon setosus</i> (Pfeiffer) Br. & Rose | Organ pipe | 4533 | .969 | Native | 1/75 | 3 |
| Hot desert cactus | Tehuacan Valley, Puebla | <i>Leontodon setosus</i> (Cault.) Br. & Rose | Cardon | 4536 | .996 | Native | 1/75 | 2 |
| Limestone hills | Sanderson Canyon, TX | <i>Leucophyllum candidum</i> I.M. Johnston | Chenizo | 4659 | .977 | Native | 10/75 | 4 |
| Desert shrub, xerophytic | Ft. Stockton, TX | <i>Lycium Torreyi</i> Gray | Wolf berry | 4316 | .991 | Native | 11/74 | 2 |
| Limestone hills | Sanderson Canyon, TX | <i>Mahonia trifoliata</i> (Nutt.) Fedde | Agarito | 4656 | .984 | Native | 10/75 | 3 |
| Desert grass | Ft. Stockton, TX | <i>Muhlenbergia Porteri</i> Scribn. | Bush muhly | 4322 | .979 | Native | 11/74 | 2 |
| Desert grass | Van Horn, TX | <i>Muhlenbergia Porteri</i> Scribn. | Bush muhly | 4329 | .979 | Native | 11/74 | 3 |
| Limestone hills | Sanderson, TX | <i>Nolina erumpens</i> (Turr.) | Bear-grass | 4654 | .979 | Native | 10/75 | 5 |
| Desert grass-land | Marathon, TX | <i>Nolina texana</i> Wats. | Bunch-grass | 4631 | .985 | Native | 10/75 | 5 |
| Prominent exposed cactus | Ft. Stockton, TX | <i>Opuntia phaeacantha</i> Engelm. | Prickly pear | 4318 | .977 | Native | 11/74 | 3 |
| Desert grass-land | Chinati Mountains, TX | <i>Opuntia phaeacantha</i> Engelm. | Prickly pear | 4616 | .953 | Native | 10/75 | 3 |
| Creosote hills | Plata, TX | <i>Opuntia violacea</i> Engelm. | Purple prickly pear | 4612 | .964 | Native | 10/75 | 2 |
| Desert | Tehuacan Valley, Puebla | <i>Opuntia</i> sp. | Prickly pear | 4535 | .982 | Native | 1/75 | 2 |
| Desert shrub often near water courses | Ft. Stockton, TX | <i>Prosopis glandulosa</i> Torr. | Mesquite | 4315 | .989 | Native | 11/74 | 3 |
| Creosote hills | Plata, TX | <i>Prosopis glandulosa</i> Torr. | Mesquite | 4611 | .981 | Native | 10/75 | 3 |
| Desert washes | Marathon, TX | <i>Prosopis glandulosa</i> Torr. | Mesquite | 4629 | .987 | Native | 10/75 | 5 |
| High desert shrub | Canada Morelos, Puebla | <i>Quercus a.f. depressipes</i> Tre) | Encino | 4539 | .982 | Native | 1/75 | 2 |
| Hot, seasonally dry low hills of Veracruz | Hirador near Huatusco, Veracruz | <i>Quercus olacoides</i> Schlect. & Chan. | Encino tesmole | 4519 | .979 | Native | 1/75 | 3 |
| Hot, seasonally dry low hills of Veracruz | Hirador near Huatusco, Veracruz | <i>Quercus puberularis</i> Nees | Encino | 4527 | .989 | Native | 1/75 | 2 |
| Limestone hills | Sanderson Canyon, TX | <i>Rhus virana</i> Gray | Evergreen sumac | 4660 | .988 | Native | 10/75 | 4 |
| Aggressive weed | Marathon, TX | <i>Salsola Kali</i> L. | Russian-thistle | 4634 | .995 | Introduced | 10/75 | 3 |
| High desert tree | Canada Morelos, Puebla | <i>Schinus molle</i> L. | Piru | 4538 | .965 | Introduced | 1/75 | 2 |

TABLE 3-1.— Continued.

| Habitat/niche | Measurement site | Botanical name | Common name | Collection number | Emissivity | Native or cultivated | Date | Number of replications |
|---|---------------------------------|--|------------------------|-------------------|------------|----------------------|-------|------------------------|
| Desert | | | | | | | | |
| Desert grass | Ft. Stockton, TX | <i>Scleropogon brevifolius</i> Phil. | Burro grass | 4314 | 0.977 | Native | 11/74 | 3 |
| Desert grass-land | Marathon, TX | <i>Setaria</i> sp. | | 4625 | .980 | Native | 10/75 | 4 |
| Agressive waddy herb | Fortin, Veracruz | <i>Sida rhombifolia</i> L. | | 4513 | .988 | Native | 1/75 | 2 |
| Roadside weed | Chiati Mountains, TX | <i>Sida</i> sp. | | 4617 | .985 | Native | 10/75 | 3 |
| Roadside weed | Chiati Mountains, TX | <i>Verbenaceae</i> family | | 4621 | .971 | Native | 10/75 | 3 |
| Agressive weed | Fortin, Veracruz | <i>Verbena turbaenensis</i> H.B.K. | | 4512 | .989 | Native | 1/75 | 2 |
| Limestone hills | Sanderson Canyon, TX | <i>Yucca Thompsoniana</i> Trel. | Yucca | 4657 | .958 | Native | 10/75 | 3 |
| Rosette form desert | Hot Springs, TX | <i>Yucca Torreyi</i> Shafer | Yucca | 4605 | .988 | Native | 10/75 | 3 |
| Pinon-Juniper | | | | | | | | |
| Pinon-Juniper belt | Davis Mountains, TX | <i>Juniperus scopulorum</i> Sarg. | Rocky mountain Juniper | 4673 | .991 | Native | 10/75 | 5 |
| Pinon-Juniper belt | Davis Mountains, TX | <i>Pinus oambroidee</i> Zull. | Hexicon Pinon | 4674 | .986 | Native | 10/75 | 5 |
| Pinon-Juniper belt | Davis Mountains, TX | <i>Pinus ponderosa</i> Laws. | Ponderosa pine | 4675 | .978 | Native | 10/75 | 5 |
| Pinon-Juniper belt | Davis Mountains, TX | <i>Quercus arizonica</i> Sarg. | Arizona oak | 4672 | .977 | Native | 10/75 | 4 |
| Pinon-Juniper belt | Davis Mountains, TX | <i>Quercus turbinella</i> Greene | Scrub oak | 4676 | .982 | Native | 1/75 | 2 |
| Mangrove | | | | | | | | |
| Coastal estuaries | Coast of Veracruz | <i>Laguncularia racemosa</i> (L.) Gaertn. f. | Black-mangrove | 4553 | .962 | Native | 1/75 | 3 |
| Coastal estuaries | Coast of Veracruz | <i>Rhizophora Mangle</i> L. | Red-mangrove | 4552 | .960 | Native | 1/75 | 2 |
| Montane rain forest | | | | | | | | |
| Secondary succession in disturbed areas | Fortin, Veracruz | <i>Acacia</i> sp. | acacia | 4506 | .952 | Native | 1/75 | 3 |
| Secondary succession in disturbed areas | Fortin, Veracruz | <i>Cecropia obtusifolia</i> | Cecropia | 4509 | .955 | Native | 1/75 | 2 |
| Coffee cover crop | Mirador near Huatusco, Veracruz | <i>Inga</i> sp. | Inga | 4507 | .970 | Cultivated | 1/75 | 3 |
| Coffee cover crop | Mirador near Huatusco, Veracruz | <i>Inga</i> sp. | Inga | 4516 | .943 | Cultivated | 1/75 | 2 |
| Montane rain forest tree | Fortin, Veracruz | <i>Pereoa schiedana</i> Nees | | 4511 | .901 | Native | 1/75 | 3 |
| Secondary growth shrubby herb | Coscomatepec, Veracruz | <i>Pluchea odorata</i> (L.) Cass. | | 4547 | .990 | Native | 1/75 | 2 |
| Montane rain forest | Fortin, Veracruz | <i>Pothomorphe umbellata</i> (L.) Hig. | | 4510 | .943 | Native | 1/75 | 2 |

TABLE 3-1.— Concluded.

| Habitat/niche | Measurement site | Botanical name. | Common name | Collection number | Emissivity | Native or cultivated | Date | Number of replications |
|---|---------------------------------|-------------------------------------|--|-------------------|---------------|----------------------|-------|------------------------|
| Temperate forest | | | | | | | | |
| Mesophyte | Galveston County, TX | <i>Ilex vomitoria</i> Ait. | Yaupon | 4335 | 0.981 .982 | Native | 12/74 | 2 1 |
| Old field invador | League City, TX | <i>Juniperus virginiana</i> L. | Eastern red cedar | 4336 | .996 | Native | 12/74 | 4 |
| Agressive understory | League City, TX | <i>Lonicera Japonica</i> Thunb. | Japanese honeysuckle | 4334 | .981 | Introduced | 12/74 | 2 |
| Wetland tree | Galveston County, TX | <i>Quercus nigra</i> L. | Water oak | 4338 | .987 .993 | Native | 12/74 | 2 1 |
| Mesophytic tree, central and coastal TX | Galveston County, TX | <i>Quercus virginiana</i> Mill. | Live oak, upper area; Live oak, lower area | 4333 | .988 | Native | 12/74 | 3 |
| Epiphyte | Galveston County, TX | <i>Tillandsia usneoides</i> (L.) L. | Spanish-moss | 4337 | .985 | Native | 12/74 | 2 |
| Tropical deciduous forest | | | | | | | | |
| Agressive weedy herb | Hirador near Hautusco, Veracruz | <i>Mangifera indica</i> L. | Mango | 4518 | .960 | Cultivated | 1/75 | 2 |
| Fruit tree | Playa Carino, Veracruz | <i>Mangifera indica</i> L. | Mango | 4557 | .960 | Cultivated | 1/75 | 2 |
| Palmar | Piedras Negras, Veracruz | <i>Sabal maritima</i> Mart. | Sabal palm | 4550 | .962 | Native | 1/75 | 3 |
| Woodland savanna | | | | | | | | |
| Woodland savanna | Playa Carino, Veracruz | <i>Acacia</i> sp. | acacia | 4551 | .952 | Native | 1/75 | 2 |
| Widespread in woodland savanna | Playa Carino, Veracruz | <i>Catba pantandra</i> | Kapok | 4556 | .966 | Native | 1/75 | 2 |
| Hot, low woodland savanna | Playa Carino, Veracruz | <i>Tabebuia rosea</i> (Bert.) D.C. | Tabebuia | 4555 | .942 | Native | 1/75 | 2 |

18

in temperature produced by shading the surface from the sun are smallest at these times. A high overcast also produces favorable conditions. Broken or low warm cloud conditions should be avoided whenever possible. However, rapid measurements and frequent replications generally produce usable results even under difficult conditions.

4. DISCUSSION

4.1 ANALYSIS OF DATA

The variability of the measured emissivity values was examined using analysis of variance techniques. Through a series of contrasts, the statistical significance of differences in emissivity between broad ecological groups was determined. The group studied and the number of observations available are given in Table 4-1. The difference between desert vegetation and all other types was clear. The hypothesis stating that the means of each ecological group were equal was strongly rejected. No significant differences were found between the two types of desert vegetation or among the four types of non-desert vegetation. However, it was found that the rain forest vegetation was significantly different from that of the temperate region. These comparisons may be seen in Table 4-2 along with a comparison of desert, rain forest, and temperate regions. This comparison showed significant differences among the group means. The means and standard deviations of each group may be seen in Table 4-3.

4.2 INTERPRETATION OF RESULTS

The results of the statistical analysis suggest the following ecologically important ideas.

As a means of avoiding excessive and possibly fatal absorption and retention of heat in the desert, desert plants reemit virtually all incoming radiation. This aids in keeping plant temperature at a viable level without benefit of the common evapotranspiration mechanisms available to more mesic plants.

Temperate region plants face less of a heat stress problem than desert plants, yet their leaf temperatures must be kept within a range consistent with their metabolic requirements. In the temperate areas of the U.S. where these plants were studied, a moisture stress develops in the late summer when temperatures are highest but soil moisture levels are low. An adaptive advantage can be speculated for plants that can increase their heat reduction during warm dry periods without increasing their evapotranspirational losses.

TABLE 4-1.— MAJOR ECOLOGICAL GROUPS EXAMINED FOR
VARIATION IN EMISSIVITY

| Group | Number of observations |
|-----------------------|------------------------|
| Dry desert | 61 |
| Humid desert | 15 |
| Montane rain forest | 11 |
| Salt water aquatic | 2 |
| Deciduous rain forest | 10 |
| Temperate region | 11 |

TABLE 4-2.— PRINCIPAL CONTRASTS OF THE ECOLOGICAL GROUPS

| Contrast | F-test | Degrees of freedom | Significance |
|---|--------|--------------------|------------------------------------|
| Desert versus all others | 21.7 | 1,108 | Highly significant |
| Dry versus humid desert | .4 | 1,74 | Not significant |
| Montane rain forest versus aquatic versus deciduous rain forest versus temperate region | 1.7 | 3,30 | Not significant |
| Deciduous rain forest versus temperate region | 5.3 | 1,30 | Significant at the 5-percent level |
| Desert versus rain forest versus temperate region | 16.1 | 2,105 | Significant at the 1-percent level |

TABLE 4-3.— MEANS AND STANDARD DEVIATIONS OF EMISSIVITY
FOR THREE VEGETATIONAL GROUPS

| Group | Emissivity | Standard deviation |
|------------------|------------|--------------------|
| Desert | 0.981 | 0.011 |
| Rain forest | .962 | .020 |
| Temperate region | .977 | .012 |

In the two tropical groups studied, the montane rain forest and tropical deciduous forest, abundant moisture occurs during the growing season. There is no shortage of moisture needed in cooling. In the cooler dry season, the deciduous forest is dormant and leafless while the montane rain forest has a lesser but still sufficient amount of moisture to meet its needs.

4.3 CONCLUSIONS

It appears from this work that there is some physiological adaptation in plants to their radiational environment. The data and analyses presented suggest that on a community-wide level, plants of the desert, tropics, and temperate regions have each adapted to deal with specific and characteristic radiation levels found in each area.

5. REFERENCES

1. BARNES C. M. and FORSBERG F. C. (June 1975) An Overview of Development of Remote Sensing Techniques for the Screwworm Eradication Program. Proceedings of the NASA Earth Resources Survey Symposium, vol. 1-A, Houston, 263-287.
2. BARTHOLIC J. F., NAMKEN L. N., and WIEGAND C. L. (Sept.-Oct. 1972) Aerial Thermal Scanner to Determine Temperatures of Soils and of Crop Canopies Differing in Water Stress. Agronomy Journal 64, 603-608.
3. BILLINGS W. D. and MORRIS R. J. (1951) Reflection of Visible and Infrared Radiation from Leaves of Different Ecological Groups. American Journal Bot. 38, 327-331.
4. BUETTNER K. J. K. and KERN C. D. (March 1965) The Determination of Infrared Emissivities of Terrestrial Surfaces. Journal of Geophysical Research 70, no. 6, 1329-1337.
5. CONAWAY J. and VAN BAVEL C. H. M. (Aug. 1967) Evaporation from a Wet Soil Surface Calculated from Radiometrically Determined Surface Temperatures. Journal of Applied Meteorology 10, no. 4, 650-655.
6. DAVIES J. A., ROBINSON P. J., and NUNEZ M. (Aug. 1971) Field Determinations of Surface Emissivity and Temperature for Lake Ontario. Journal of Applied Meteorology 10, no. 4, 811-819.
7. FALCKENBERG G. (1928) :Absorptionskonstanten einiger meteorologisch wichtiger Korper fur infrarote Wellen. Meteorol. Z. 45, 334-337.
8. FUCHS M. and TANNER C. B. (Nov.-Dec. 1966) Infrared Thermometry of Vegetation. Agronomy Journal 58, 597-601.
9. FUCHS M. and TANNER C. B. (April 1968) Surface Temperature Measurements of Bare Soils. Journal of Applied Meteorology 7, no. 2, 303-305.
10. GATES D. M. and TANTRAPORN W. (June 6, 1952) The Reflectivity of Deciduous Trees and Herbaceous Plants in the Infrared to 25 Microns. Science 115, no. 2997, 613-616.
11. GATES M., KEEGAN H. J., SCHLETER J. C. and WEIDNER V. R. (Jan. 1965) Spectral Properties of Plants. Applied Optics 4, no. 1, 11-20.
12. HOVIS W. A. (1966) Optimum Wavelength Intervals for Surface Temperature Radiometry. Applied Optics 5, 815-818.

13. HUSCHKE R. E., ed. (1959) Glossary of Meteorology, American Meteorological Society, Boston, 200 pp.
14. IDSO S. B. and JACKSON R. D. (Feb. 1969) Comparison of Two Methods for Determining Infrared Emittances of Bare Soils. Journal of Applied Meteorology 8, no. 1, 168-169.
15. LORENZ D. (Aug. 1966) The Effect of the Long-wave Reflectivity of Natural Surface Temperature Measurements Using Radiometers. Journal of Applied Meteorology 5, no. 4, 421-430.
16. World Meteorological Organization (1969) Guide to Meteorological Instrument and Observing Practices, WMO-8, T.P. 3.