

NASA Technical Memorandum 78556

Viscosity and Thermal Conductivity of Model Jupiter Atmospheres

C. Frederick Hansen
Ames Research Center
Moffett Field, California

NASA

National Aeronautics
and Space Administration

**Scientific and Technical
Information Office**

1979

VISCOSITY AND THERMAL CONDUCTIVITY
OF MODEL JUPITER ATMOSPHERES

C. Frederick Hansen

Ames Research Center

SUMMARY

The viscosity and thermal conductivity coefficients are estimated for three models of the atmosphere of Jupiter: a Heavy Model consisting of 22% helium and 78% hydrogen, a Nominal Model consisting of 11% helium and 89% hydrogen, and a Light Model consisting of pure hydrogen. The effect of trace elements is neglected. Linearized approximations are used for the transport coefficients of the mixtures; these are found to be in almost constant ratio to the values for pure hydrogen, independent of temperature. Short basic language programs for computing the coefficients are listed.

INTRODUCTION

The Jovian atmosphere is presently thought to consist mainly of molecular hydrogen, H_2 , perhaps mixed with helium, He, and with traces of methane, CH_4 , and ammonia, NH_3 . The amount of He seems to be the most uncertain; estimates range from about 11% by volume for the Nominal Jovian Atmosphere Model, to 22% for the Heavy Models, and to 0.0% for the Light Models. The traces of methane are thought to be about 0.1% and of ammonia about 0.02%; these are significant amounts as far as radiation absorption and emissivity are concerned, but are trivial amounts as far as thermodynamic and transport properties of the atmosphere are concerned. The temperature range of interest is from 100 to 1000 K. The thermodynamic properties needed for Jupiter probe design have been previously calculated and tabulated for this range (J. Sperans, Ames Research Center, NASA, private communication). Recently, a need has appeared for estimates of viscosity and thermal conductivity of the Jovian atmosphere, in connection with the subsonic motion of bodies through that atmosphere. The present report gives estimates of these quantities based on linearized formulas for the transport properties of mixtures and neglecting the effect of traces of CH_4 and NH_3 .

VISCOSITY OF MODEL JUPITER ATMOSPHERES

Viscosity of Mixtures

According to rigorous kinetic theory, the coefficient of viscosity of a multicomponent gas mixture should be expressed as a ratio of determinants.

However, Hirschfelder et al. (ref. 1) show that the off-diagonal terms are small compared to the trace elements of these matrices and that the major contribution to the viscosity of the mixture can be expressed as

$$\eta = \sum_i \frac{x_i^2}{H_{ii}} + \dots \quad (1)$$

where x_i is the mol fraction of the component i and the matrix trace elements H_{ii} are

$$H_{ii} = \frac{x_i^2}{\eta_{ii}} + \sum_{k \neq i} \frac{2x_i x_k}{\eta_{ik}} \frac{m_i m_k}{(m_i + m_k)^2} \left(\frac{5}{3A_{ik}} + \frac{m_k}{m_i} \right) \quad (2)$$

The coefficients η_{ik} are defined as

$$\eta_{ik} = \frac{5}{16Q_{ik}} \left(2\pi kT \frac{m_i m_k}{m_i + m_k} \right)^{1/2} \quad (3)$$

where m_i is the molecular weight of component i and Q_{ik} is the Chapman-Enskog cross section for momentum transfer in collisions between particles i and k . The coefficient A_{ik} is the ratio of Q_{ik} and the cross sections appropriate for mass diffusion; it is somewhat greater than unity because deeper penetration of the interparticle potential is required to influence particle flux than for momentum transfer. In fact, if A_{ik} is assumed equal to 5/3, the off-diagonal elements of the determinants vanish exactly (ref. 1); although this is somewhat too large a value, the effect on H_{ii} from using this approximation is not serious, because terms with $m_k \ll m_i$ contribute little to the sum in equation (2). With this simplification, the elements H_{ii} become

$$H_{ii} = 2x_i \sum_k \frac{x_k}{\eta_{ik}} \left(1 + \frac{m_i}{m_k} \right)^{-1} \quad (4)$$

and equation (1) can be expressed in the simple linear form

$$\eta = \frac{5\pi}{32} \sum_i \rho_i \bar{u}_i \lambda_i \quad (5)$$

if the effective mean free path λ_i is defined

$$\lambda_i^{-1} = \sqrt{2}n \sum_k x_k Q_{ik} \left(\frac{1 + m_i/m_k}{2} \right)^{-1/2} \quad (6)$$

Hansen (ref. 2) shows that this value of λ_i can be given a simple physical interpretation as the mean distance traveled by a molecule of type i

before it loses its excess momentum relative to the surrounding gas medium in a series of head-on elastic conditions, since the fraction of excess momentum transferred in a single collision between particles i and k is

$$f_{ik} = \frac{2}{1 + m_i/m_k} \quad (7)$$

and the fraction of excess momentum transferred per unit time is the product of f_{ik} and the collision rate θ_{ik} summed over all k :

$$\sum_k f_{ik} \theta_{ik} = \sqrt{2} n \bar{u}_i \sum_k \frac{x_k Q_{ik}}{\left(\frac{1 + m_i/m_k}{2}\right)} \quad (8)$$

Multiplying the reciprocal of equation (8) by the mean velocity \bar{u}_i of particle i leads to the expression of equation (6). Equation (5) can thus be expressed

$$\eta = \sum_i \eta_i \left(\frac{\lambda_i}{\lambda_i^0} \right) \quad (9)$$

where the η_i values are the viscosities of the pure components of the mixture, and λ_i^0 is the mean free path in a pure gas of species i at the density ρ_i . Thus,

$$\frac{\lambda_i^0}{\lambda_i} = 1 + \sum_{k \neq i} \frac{x_k Q_{ik}}{x_i Q_{ii}} \left(\frac{1 + m_i/m_k}{2} \right)^{-1/2} \quad (10)$$

Buddenberg and Wilke (ref. 3) empirically find that agreement with experimental measurements of viscosities of many gas mixtures is improved if each term in the summation of equation (10) is multiplied by 0.69; presumably, this correction accounts for the deviation of A_{ik} from the 5/3 value assumed. Accordingly, the formula used for the calculations of the viscosity of the Jupiter atmosphere is

$$\frac{\lambda_i^0}{\lambda_i} = 1 + 0.69 \sum_{k \neq i} \frac{x_k Q_{ik}}{x_i Q_{ii}} \left(\frac{1 + m_i/m_k}{2} \right)^{-1/2} \quad (10a)$$

The collision cross sections Q_{11} are most expeditiously determined by fitting the known viscosities of the pure species with the theoretical formula:

$$\eta_1 = \frac{5\pi}{32} \rho_1 u_1 \lambda_1^0 = \frac{5\pi}{32} \left(\frac{2m_1 kT}{\pi} \right)^{1/2} Q_{11}^{-1} \quad (11)$$

A convenient functional form used for Q_{11} in making this fit is

$$Q_{11}(T) = Q_0 \left(\frac{T}{T_0} \right)^n \left(\frac{1 + C/T_0}{1 + C/T} \right) \quad (12)$$

In equations (5) and (6) it is assumed that there is no dependence of viscosity on density (usually a good approximation except at very high densities where the collisions can no longer be considered as simple two-body collisions), so that the entire temperature dependence of the coefficient of viscosity is due to the $T^{1/2}$ dependence of the mean molecular velocity and to the decrease in cross-section size as the collision energy increases.

The cross sections Q_{ik} are determined from the pure species cross sections, assuming that the relation between them is the same as for hard spheres,

$$Q_{ik} = \left(\frac{Q_{11}^{1/2} + Q_{kk}^{1/2}}{2} \right)^2 \quad (13)$$

namely, the collision diameter for the unlike collision pair is the average of the collision diameters for the pure specie collisions. This hard-sphere type approximation can be expected to be quite good when the colliding species both have closed-shell electronic structures, which is the case for both H_2 and He, the species of concern for the Jovian atmosphere.

Viscosity of Pure Molecular Hydrogen

The coefficient of viscosity of pure H_2 is reproduced reasonably well over the temperature range of 100 to 1000 K by the formula

$$\eta = \frac{5\pi}{32} \left(\frac{2mkT}{\pi} \right)^{1/2} Q^{-1} = 90.6 \left(\frac{T}{300} \right)^{0.6658} (1 + 4.0/T)^{-1} \mu P \quad (14)$$

where m is the mass of the H_2 molecule. Accordingly, the momentum transfer cross section Q for $H_2 - H_2$ collisions is taken to be

$$Q = 32.2 \times 10^{-16} \left(\frac{300}{T} \right)^{0.1658} \left(1 + \frac{4.0}{T} \right) \text{cm}^2 \quad (15)$$

The deviations of equation (14) from the H_2 coefficients of viscosity given by Watson (ref. 4), by Kestin (ref. 5), by Hilsenrath et al. (ref. 6),

and by Keyes (ref. 7) are shown in figure 1. The parameters of equation (14) were chosen to give the best fit to Hilsenrath's values from 100 to 300 K and to Watson's values from 300 to 1000 K. The latter are expected to be the most reliable in that range since they have been carefully fitted to the most recent and most carefully calibrated experimental data. Hilsenrath's values are based on collision integrals calculated using a Lennard-Jones 6-12 potential, with the force parameters of the potential fitted to older viscosity measurements; these are expected to be the best values in the low-temperature range. It can be seen that the deviations of equation (14) are generally well within 1% of the best values. Even the old formula given by Keyes for the range of 100 to 500 K agrees within 3%,




Figure 1.- Deviations of equation (14) from published viscosities of pure H₂.

$$\eta = \frac{6.23T^{1/2}}{1 + \frac{70.8}{T} e^{-39.8/T}} \mu P \quad (16)$$

and the value documented by Kestin at 293 K falls within the 1% range. In view of uncertainties in the mixture formulas, equation (14) is considered adequately precise.

Viscosity of Pure Helium

The coefficient of viscosity of pure He is reproduced reasonably well over the temperature range of 100 to 1000 K by the formula

$$\eta = \frac{5\pi}{32} \left(\frac{2mkT}{\pi} \right)^{1/2} Q^{-1} = 191.6 \left(\frac{T}{300} \right)^{0.7176} \left(1 - \frac{11.4}{T} \right)^{-1} \mu P \quad (17)$$

where m is the mass of the He atom. Accordingly, the momentum transfer cross section Q for He-He collisions is taken to be

$$Q = 21.5 \times 10^{-16} \left(\frac{300}{T} \right)^{0.2176} \left(1 - \frac{11.4}{T} \right) \text{cm}^2 \quad (18)$$




Figure 2.- Deviations of equation (17) from published viscosities of pure He.

The deviations of equation (17) from the He coefficients of viscosity given by Watson (ref. 4), by Kestin (ref. 5), by Kestin and Leidenfrost (ref. 8), and by Keyes (ref. 7) are shown in figure 2. Again the values of Watson are considered to be the best in the range from 300 + 1000 K. The formula has been fitted to these values and to the 100 K value of Kestin and Leidenfrost. The deviations from Watson's values are generally less than 0.5%; in the low-temperature range, where the formula bisects the values of Kestin and Leidenfrost and of the Keyes equation designed for the range of 104 to 373 K,

$$\eta = \frac{18.05T^{1/2}}{1 + \frac{253}{T} e^{-115.1/T}} \mu\text{P} \quad (19)$$

deviations are generally less than 2%. Again, the uncertainties are consistent with the uncertainties in the mixture formulas.

Viscosity of Jovian Atmospheres

The viscosity of model Jovian atmosphere mixtures is shown as a function of temperature in table 1 for the temperature range from 100 to 1000 K. The first column is the temperature; the second column is the viscosity of pure H_2 (which corresponds to both the Cool Light Model and Warm Light Model of the Jovian Atmosphere); the third column is the viscosity of an 89% H_2 - 11% He mixture (which corresponds to the Nominal Jovian Atmosphere Model); and the final column is the viscosity of a 78% H_2 - 22% He mixture (which corresponds to both the Cool Heavy Model and the Warm Heavy Model of the Jovian Atmosphere). The Cool and Warm Models have only a different altitude dependence

of temperature. The viscosity of the Nominal Model Atmosphere (with 11% He) is almost exactly 25% greater than the viscosity of pure H₂. The ratio of the two is almost independent of temperature. Similarly, the viscosity of the Heavy Atmosphere Models (with 22% He) is almost exactly 47% greater than the viscosity of pure H₂.

TABLE 1.- VISCOSITY OF JUPITER MODEL ATMOSPHERES

| Temperature, K | Light atm., ^a μP | Nominal atm., ^b μP | Heavy atm., ^c μP |
|----------------|-----------------------------|-------------------------------|-----------------------------|
| 100 | 41.92 | 52.98 | 62.77 |
| 150 | 55.62 | 69.92 | 82.46 |
| 200 | 67.8 | 85.08 | 100.17 |
| 250 | 78.97 | 99.02 | 116.52 |
| 300 | 89.4 | 112.08 | 131.87 |
| 350 | 99.25 | 124.44 | 146.42 |
| 400 | 108.64 | 136.23 | 160.32 |
| 450 | 117.63 | 147.54 | 173.68 |
| 500 | 126.29 | 158.46 | 186.57 |
| 550 | 134.66 | 169.02 | 199.06 |
| 600 | 142.78 | 179.27 | 211.19 |
| 650 | 150.67 | 189.24 | 223.01 |
| 700 | 158.36 | 198.97 | 234.54 |
| 750 | 165.86 | 208.47 | 245.81 |
| 800 | 173.2 | 217.77 | 256.84 |
| 850 | 180.39 | 226.88 | 267.65 |
| 900 | 187.44 | 235.81 | 278.27 |
| 950 | 194.35 | 244.59 | 288.7 |
| 1000 | 201.15 | 253.22 | 298.95 |

^aViscosity of pure H₂.

^bViscosity of 89% H₂ - 11% He mixture.

^cViscosity of 78% H₂ - 22% He mixture.

Table 2 shows the viscosity of the Jupiter Nominal Atmosphere as a function of altitude and the ratio of the mixture viscosity to the pure H₂ viscosity.

TABLE 2.- VISCOSITY OF JUPITER NOMINAL ATMOSPHERE

| Altitude, km | Temperature, K | Viscosity, μP | Viscosity ratio, mix/H ₂ |
|--------------|----------------|--------------------------|-------------------------------------|
| -270.79 | 681.77 | 195.5 | 1.256 |
| -246.93 | 637.33 | 186.79 | 1.256 |
| -224.62 | 595.74 | 178.45 | 1.255 |
| -203.74 | 556.83 | 170.48 | 1.255 |
| -184.23 | 520.44 | 162.85 | 1.255 |
| -165.98 | 486.39 | 155.56 | 1.254 |
| -148.91 | 454.55 | 148.59 | 1.254 |
| -132.96 | 424.78 | 141.93 | 1.254 |
| -118.03 | 396.93 | 135.55 | 1.254 |
| -104.09 | 370.89 | 129.46 | 1.254 |
| -91.06 | 346.51 | 123.62 | 1.254 |
| -78.67 | 323.66 | 118.03 | 1.253 |
| -67.49 | 302.25 | 112.68 | 1.253 |
| -56.86 | 282.17 | 107.54 | 1.253 |
| -46.94 | 263.31 | 102.6 | 1.254 |
| -37.66 | 245.59 | 97.86 | 1.254 |
| -29.03 | 228.92 | 93.29 | 1.254 |
| -20.97 | 213.21 | 88.88 | 1.254 |
| -13.48 | 198.4 | 84.63 | 1.255 |
| -6.49 | 184.43 | 80.53 | 1.255 |
| 0 | 171.23 | 76.55 | 1.256 |
| 6.02 | 158.77 | 72.71 | 1.256 |
| 11.6 | 147 | 68.98 | 1.257 |
| 16.75 | 134.3 | 64.85 | 1.258 |
| 21.43 | 121.6 | 60.59 | 1.26 |
| 25.79 | 116.65 | 58.9 | 1.261 |
| 29.97 | 111.7 | 57.17 | 1.261 |
| 34.04 | 110.9 | 56.89 | 1.262 |
| 38.09 | 110.1 | 56.61 | 1.262 |
| 42.16 | 112.35 | 57.4 | 1.261 |
| 46.32 | 114.6 | 58.18 | 1.261 |
| 50.57 | 117.29 | 59.12 | 1.261 |
| 54.91 | 119.98 | 60.04 | 1.26 |
| 59.36 | 122.66 | 60.96 | 1.26 |
| 63.9 | 125.35 | 61.87 | 1.259 |

TABLE 2.- CONTINUED

| Altitude, km | Temperature, K | Viscosity, μP | Viscosity ratio, mix/H ₂ |
|--------------|----------------|--------------------------|--|
| 68.54 | 128.04 | 62.77 | 1.259 |
| 73.29 | 130.73 | 63.67 | 1.259 |
| 78.13 | 133.41 | 64.56 | 1.258 |
| 83.07 | 136.1 | 65.45 | 1.258 |
| 88.1 | 138.1 | 66.1 | 1.258 |
| 93.2 | 140.09 | 66.75 | 1.258 |
| 98.37 | 142.09 | 67.4 | 1.258 |
| 103.62 | 144.09 | 68.05 | 1.257 |
| 108.94 | 146.08 | 68.69 | 1.257 |
| 114.34 | 148.08 | 69.33 | 1.257 |
| 119.81 | 150.08 | 69.97 | 1.257 |
| 125.36 | 152.08 | 70.6 | 1.257 |
| 130.98 | 154.07 | 71.23 | 1.257 |
| 136.67 | 156.07 | 71.86 | 1.257 |
| 142.44 | 158.07 | 72.49 | 1.256 |
| 148.28 | 160.06 | 73.11 | 1.256 |
| 154.2 | 162.06 | 73.73 | 1.256 |
| 160.19 | 164.06 | 74.35 | 1.256 |
| 166.25 | 166.05 | 74.97 | 1.256 |
| 172.39 | 168.05 | 75.58 | 1.256 |
| 178.61 | 170.05 | 76.19 | 1.256 |
| 184.9 | 172.04 | 76.8 | 1.256 |
| 191.26 | 174.04 | 77.41 | 1.256 |
| 197.7 | 176.04 | 78.01 | 1.255 |
| 204.21 | 178.03 | 78.61 | 1.255 |
| 210.8 | 180.03 | 79.21 | 1.255 |
| 217.46 | 182.03 | 79.81 | 1.255 |
| 224.2 | 184.03 | 80.41 | 1.255 |
| 231.01 | 186.02 | 81 | 1.255 |
| 237.89 | 188.02 | 81.59 | 1.255 |
| 244.85 | 190.02 | 82.18 | 1.255 |
| 251.89 | 192.01 | 82.77 | 1.255 |
| 259 | 194.01 | 83.35 | 1.255 |
| 266.19 | 196.01 | 83.94 | 1.255 |
| 273.45 | 198 | 84.52 | 1.255 |

TABLE 2.-- CONTINUED

| Altitude, km | Temperature, K | Viscosity, μP | Viscosity ratio, mix/H_2 |
|--------------|----------------|--------------------------|--|
| 280.78 | 200 | 85.1 | |
| 288.16 | 200 | 85.1 | 1.255 |
| 295.53 | 200 | 85.1 | 1.255 |
| 302.91 | 200 | 85.1 | 1.255 |
| 310.29 | 200 | 85.1 | 1.255 |
| 317.67 | 200 | 85.1 | 1.255 |
| 325.05 | 200 | 85.1 | 1.255 |
| 332.44 | 200 | 85.1 | 1.255 |
| 339.82 | 200 | 85.1 | 1.255 |
| 347.21 | 200 | 85.1 | 1.255 |
| 354.6 | 200 | 85.1 | 1.255 |
| 362.39 | 221.67 | 91.27 | 1.254 |
| 370.98 | 243.33 | 97.24 | 1.254 |
| 380.37 | 265 | 103.05 | 1.254 |
| 390.57 | 286.67 | 108.7 | 1.253 |
| 401.58 | 308.33 | 114.21 | 1.253 |
| 413.39 | 330 | 119.6 | 1.253 |
| 426 | 351.67 | 124.87 | 1.254 |
| 439.42 | 373.33 | 130.03 | 1.254 |
| 453.65 | 395 | 135.1 | 1.254 |
| 468.69 | 416.67 | 140.08 | 1.254 |
| 484.54 | 438.33 | 144.98 | 1.254 |
| 501.19 | 460 | 149.8 | 1.254 |
| 518.66 | 481.67 | 154.54 | 1.254 |
| 536.95 | 503.33 | 159.21 | 1.255 |
| 556.04 | 525 | 163.82 | 1.255 |
| 575.96 | 546.67 | 168.37 | 1.255 |
| 596.69 | 568.33 | 172.85 | 1.255 |
| 618.24 | 590 | 177.29 | 1.255 |
| 640.6 | 611.67 | 181.67 | 1.256 |
| 663.79 | 633.33 | 185.99 | 1.256 |
| 687.8 | 655 | 190.27 | 1.256 |
| 712.64 | 676.67 | 194.51 | 1.256 |
| 738.3 | 698.33 | 198.7 | 1.256 |
| 764.79 | 720 | 202.85 | 1.256 |

TABLE 2.- CONCLUDED

| Altitude, km | Temperature, K | Viscosity, μP | Viscosity ratio, mix/H ₂ |
|--------------|----------------|--------------------------|-------------------------------------|
| 792.11 | 741.67 | 206.96 | 1.257 |
| 820.27 | 763.33 | 211.02 | 1.257 |
| 849.25 | 785 | 215.06 | 1.257 |
| 879.07 | 806.67 | 219.05 | 1.257 |
| 909.73 | 828.33 | 223.01 | 1.257 |
| 941.23 | 850 | 226.94 | 1.258 |

THERMAL CONDUCTIVITY OF JUPITER ATMOSPHERES

Thermal Conductivity of Mixtures

Theoretical expressions for thermal conductivity of gases and of gas mixtures have not been verified with the same accuracy as for viscosities. This occurs partly because a number of intricate corrections are involved in making measurements, and also because the heat transfer associated with internal modes of molecular energy is difficult to treat with the same rigor as that associated with the kinetic modes of energy. In the simple Eucken approximation, the internal degrees of freedom are assumed to have a Maxwell distribution that is totally independent of the molecule's velocity. This is certainly a good approximation for the stiff vibrational modes of the H₂ molecule at the temperatures of interest, where there is very little coupling between vibrational and kinetic modes during a collisional perturbation; however, the same may not be exactly true for the rotational modes of the H₂ molecule. Nevertheless, the Eucken approximation seems to agree just as well with experimental data as the more elaborate theoretical models that have been devised; consequently, it will be used here in view of its simplicity and the fact that the comparisons between experiment and the theoretical model seem adequate to the order of accuracy being considered.

The coefficient of thermal conductivity of a pure gas can be expressed with the Eucken approximation

$$\kappa = \frac{5\pi}{32} \frac{\rho u \lambda}{M} \left(\frac{5}{2} C_v + C_v^* \right) \quad (20)$$

where M is the gram molecular weight, C_v is the specific heat per mole of the gas associated with its kinetic energy, and C_v^* is the specific heat associated with internal molecular energy. For a Maxwell-Boltzmann distribution of velocities, the kinetic specific heat is just $3/2$ times the gas constant R

$$C_v = \frac{3R}{2} \quad (21)$$

The internal specific heat is zero for He and for H₂ it is the sum of the rotational and vibrational contributions:

$$C_v^* (\text{He}) = 0 \quad (22a)$$

$$C_v^* (\text{H}_2) = R \left[1 + \left(\frac{\theta_v/2T}{\sinh \theta_v/2T} \right)^2 \right] \quad (22b)$$

In equations (22a) and (22b) it has been assumed that no electronic energy is excited for either He or H₂ (which is a good assumption below 1000 K) and that the rotational states of H₂ are fully excited while the vibrational modes are at the equilibrium excitation for a harmonic oscillator. Since the characteristic vibrational temperature for H₂ is large ($\theta_v = 6159$ K) the latter assumption is very good below 1000 K; however, the characteristic rotational temperature for H₂ is rather large ($\theta_r = 85.42$ K), so the approximation of equation (22b) can be expected to hold some error at the low-temperature end of the range near 100 K.

The thermal conductivity of the gas mixture is now expressed

$$\kappa = \frac{5\pi}{32} \sum_i \frac{\rho_i \bar{u}_i \lambda_i^0}{m_i} \left(\frac{\lambda_i}{\lambda_i^0} \right) \left(\frac{5}{2} C_i + C_i^* \right) \quad (23)$$

Strictly speaking, the ratio of the effective mean free path for energy transfer (λ_i) to the mean free path in the pure *i*th component with density ρ_i (λ_i^0) should be somewhat different for the translational and internal degrees of freedom. The difference is not large, however, if the molecular masses are about equal, which is the case for the present gases of interest. The fraction of excess kinetic energy, associated with the motion between the line of centers at impact, transferred between particle *i* and *k* in elastic collision is

$$f_{ik} = \frac{4m_i/m_k}{1 + m_i/m_k} \quad (24)$$

Accordingly, the excess kinetic energy transferred per unit time is the product of the collision rate θ_{ik} and the fraction f_{ik} summed over all *k*

$$\sum_k f_{ik} \theta_{ik} = \sqrt{2} n \bar{u}_i \sum_k \frac{x_k \frac{m_i}{m_k} Q_{ik}}{\left(\frac{1 + m_i/m_k}{2} \right)^2} \quad (25)$$

and the ratio λ_1^0/λ_1 can be expressed

$$\frac{\lambda_1^0}{\lambda_1} = 1 + \sum_{k \neq 1} \frac{x_k}{x_1} \frac{Q_{1k}}{Q_{11}} \frac{m_1/m_k}{\left(\frac{1 + m_1/m_k}{2}\right)^{3/2}} \quad (26)$$

In this approximation, the thermal conductivity of the mixture is given by a simple linear expression analogous to the linearized formula for the viscosities of mixtures

$$\kappa = \sum_i \kappa_i \left(\frac{\lambda_1}{\lambda_1^0}\right) \quad (27)$$

Undoubtedly, an empirical correction term, such as that provided by Buddenberg and Wilke for the viscosity of mixtures, could account for some difference in the cross sections for diffusion and for energy transfer and thus improve the theoretical model somewhat; however, such a correction has not been established at present. Since the molecular masses are nearly equal in this case, the correction could only be a modest one, of the order of a few percent at best. Such a correction takes on major importance only in ionized gases where the very lightweight electrons carry a large fraction of the energy transfer.

Thermal Conductivity of Pure Molecular Hydrogen

The coefficient of thermal conductivity of pure H_2 is reproduced reasonably well over the temperature range of 100 to 1000 K by the formula

$$\begin{aligned} \kappa &= \frac{5\pi}{32} \left(\frac{2mkT}{\pi}\right)^{1/2} \left[\frac{19}{4} + \left(\frac{\theta_v/2T}{\sinh \theta_v/2T}\right)^2 \right] \frac{R}{mQ} \\ &= 1.10 \times 10^{-4} \left(\frac{T/300}{1 + \frac{49.4}{T}}\right)^{0.6983} \left[\frac{19}{4} + \left(\frac{3080/T}{\sinh \frac{3080}{T}}\right)^2 \right] \text{cal/cm deg sec} \quad (28) \end{aligned}$$

where again m is the mass of the H_2 molecule. Accordingly, the energy transfer cross section Q , for $H_2 - H_2$ collisions, is taken to be

$$Q = 26.1 \times 10^{-16} \left(\frac{300}{T}\right)^{0.1983} \left(1 + \frac{49.4}{T}\right) \text{cm}^2 \quad (29)$$

The deviation of equation (28) from the H_2 coefficients of thermal conductivity given by Hilsenrath et al. (ref. 6) is shown in figure 3. The deviations, generally less than 1%, are adequate for present purposes. No




Figure 3.- Deviations of equation (28) from published thermal conductivities of pure H₂

doubt, some of the deviation near 100 K is due to the approximate treatment of the equilibrium rotational excitation.

Thermal Conductivity of Pure Helium

The coefficient of thermal conductivity of pure He is reproduced reasonably well over the temperature range of 100 to 1000 K by the formula

$$\kappa = \frac{5\pi}{32} \left(\frac{2mkT}{\pi} \right)^{1/2} \frac{15}{4} \frac{R}{mQ} = 3.42 \times 10^{-4} \frac{(T/300)^{0.7412}}{(1 - 13.7/T)} \text{ cal/cm deg sec} \quad (30)$$

where now m is the mass of the He atom. Accordingly, the energy transfer cross section Q , for He - He collisions, is taken to be

$$Q = 5.96 \times 10^{-16} \left(\frac{300}{T} \right)^{0.7412} \times \left(1 - \frac{13.7}{T} \right) \text{ cm}^2 \quad (31)$$

The deviations of equation (30) from the He coefficients of thermal conductivity given by Powell, Ho, and Liley (ref. 9) are shown in figure 4. The deviations are generally within 2%. Some deviations from coefficients given by Powell (ref. 10) are also shown, with about the same result. These deviations are about double the deviations previously encountered for the simple cross-section expressions, but are still well within the uncertainties of the mixture formulas used for thermal conductivities.




Figure 4.- Deviations of equation (30) from published thermal conductivities of pure He

Thermal Conductivity of Jovian Atmospheres

The thermal conductivity of model Jovian atmosphere mixtures is shown as a function of temperature in table 3 for the temperature range from 100 to 1000 K. Again, the first column is the temperature; the second column is the thermal conductivity of pure H₂ (which corresponds to both the Cool Light Model and the Warm Light Model of the Jovian Atmosphere), the third column is the conductivity of an 89% H₂ - 11% He mixture (which corresponds to the Nominal Jovian Atmosphere Model); and the final column is the conductivity of a 78% H₂ - 22% He mixture (which corresponds to both the Cool Heavy Model and Warm Heavy Model of the Jovian Atmosphere). The conductivity of the Nominal Model Atmosphere (with 11% He) is almost constantly 2% less than for pure H₂, independent of temperature. Similarly, the conductivity of the Heavy Atmosphere Models (with 22% He) is almost constantly 4% less than for pure H₂.

TABLE 3.- THERMAL CONDUCTIVITY OF JUPITER MODEL ATMOSPHERES

| Temperature, K | Thermal conductivity, mcal/cm-deg-sec | | |
|----------------|---------------------------------------|---------------------------|-------------------------|
| | Light atm. ^a | Nominal atm. ^b | Heavy atm. ^c |
| 100 | 0.1623 | 0.161 | 0.1597 |
| 150 | .2422 | .2395 | .2365 |
| 200 | .3156 | .3115 | .307 |
| 250 | .3841 | .3787 | .3727 |
| 300 | .4486 | .4419 | .4345 |
| 350 | .5099 | .5021 | .4933 |
| 400 | .5685 | .5596 | .5496 |
| 450 | .6249 | .6149 | .6038 |
| 500 | .6794 | .6684 | .6561 |
| 550 | .7323 | .7203 | .7069 |
| 600 | .7839 | .771 | .7565 |
| 650 | .8344 | .8205 | .805 |
| 700 | .884 | .8693 | .8527 |
| 750 | .9331 | .9174 | .8998 |
| 800 | .9816 | .965 | .9464 |
| 850 | 1.0299 | 1.0123 | .9926 |
| 900 | 1.0779 | 1.0594 | 1.0387 |
| 950 | 1.1258 | 1.1064 | 1.0845 |
| 1000 | 1.1737 | 1.1533 | 1.1303 |

^aThermal conductivity of pure H₂.

^bThermal conductivity of 89% H₂ - 11% He mixture.

^cThermal conductivity of 78% H₂ - 28% He mixture.

Table 4 shows the thermal conductivity of the Jupiter Nominal Atmosphere as a function of altitude and the ratio of the mixture conductivity to the pure H₂ conductivity.

TABLE 4.- THERMAL CONDUCTIVITY OF JUPITER NOMINAL ATMOSPHERE

| Altitude, km | Temperature, K | Conductivity, mcal/cm-deg sec | Conductivity ratio, mix/H ₂ |
|--------------|----------------|----------------------------------|---|
| -270.79 | 681.77 | 0.8516 | 0.9833 |
| -246.93 | 637.33 | .8081 | .9834 |
| -224.62 | 595.74 | .7667 | .9835 |
| -203.74 | 556.83 | .7273 | .9836 |
| -184.23 | 520.44 | .6898 | .9837 |
| -165.98 | 486.39 | .654 | .9839 |
| -148.91 | 454.55 | .6199 | .984 |
| -132.96 | 424.78 | .5873 | .9842 |
| -118.03 | 396.93 | .5561 | .9843 |
| -104.09 | 370.89 | .5264 | .9845 |
| - 91.06 | 346.51 | .498 | .9847 |
| - 78.87 | 323.66 | .4707 | .9849 |
| - 67.49 | 302.25 | .4447 | .9851 |
| - 56.86 | 282.17 | .4198 | .9854 |
| - 46.94 | 263.31 | .3959 | .9856 |
| - 37.66 | 245.59 | .3729 | .9859 |
| - 29.03 | 228.92 | .3509 | .9862 |
| - 20.97 | 213.21 | .3297 | .9866 |
| - 13.48 | 198.4 | .3093 | .987 |
| - 6.49 | 184.43 | .2897 | .9874 |
| 0 | 171.23 | .2707 | .9878 |
| 6.02 | 158.77 | .2525 | .9883 |
| 11.6 | 147 | .2349 | .9888 |
| 16.75 | 134.3 | .2156 | .9895 |
| 21.43 | 121.6 | .1958 | .9903 |
| 25.79 | 116.65 | .188 | .9906 |
| 29.97 | 111.7 | .18 | .991 |
| 34.04 | 110.9 | .1788 | .991 |
| 38.09 | 110.1 | .1775 | .9911 |
| 42.16 | 112.35 | .1811 | .9909 |
| 46.32 | 114.6 | .1847 | .9908 |
| 50.57 | 117.29 | .189 | .9906 |
| 54.91 | 119.98 | .1932 | .9904 |
| 59.36 | 122.66 | .1975 | .9902 |
| 63.9 | 125.35 | .2017 | .99 |

TABLE 4.- CONTINUED

| Altitude, km | Temperature, K | Conductivity, mcal/cm-deg sec | Conductivity ratio, mix/H ₂ |
|--------------|----------------|----------------------------------|---|
| 68.54 | 128.04 | 0.2059 | 0.9898 |
| 73.29 | 130.73 | .2101 | .9897 |
| 78.13 | 133.41 | .2142 | .9895 |
| 83.07 | 136.1 | .2184 | .9894 |
| 88.1 | 138.1 | .2214 | .9893 |
| 93.2 | 140.09 | .2245 | .9892 |
| 98.37 | 142.09 | .2275 | .9891 |
| 103.62 | 144.09 | .2306 | .989 |
| 108.94 | 146.08 | .2336 | .9889 |
| 114.34 | 148.08 | .2366 | .9888 |
| 119.81 | 150.08 | .2396 | .9887 |
| 125.36 | 152.08 | .2426 | .9886 |
| 130.98 | 154.07 | .2455 | .9885 |
| 136.67 | 156.07 | .2485 | .9884 |
| 142.44 | 158.07 | .2515 | .9883 |
| 148.28 | 160.06 | .2544 | .9882 |
| 154.2 | 162.06 | .2574 | .9882 |
| 160.19 | 164.06 | .2603 | .9881 |
| 166.25 | 166.05 | .2632 | .988 |
| 172.39 | 168.05 | .2661 | .9879 |
| 178.61 | 170.05 | .269 | .9879 |
| 184.9 | 172.04 | .2719 | .9878 |
| 191.26 | 174.04 | .2748 | .9877 |
| 197.7 | 176.04 | .2777 | .9876 |
| 204.21 | 178.03 | .2805 | .9876 |
| 210.8 | 180.03 | .2834 | .9875 |
| 217.46 | 182.03 | .2863 | .9874 |
| 224.2 | 184.03 | .2891 | .9874 |
| 231.01 | 186.02 | .2919 | .9873 |
| 237.89 | 188.02 | .2948 | .9873 |
| 244.85 | 190.02 | .2976 | .9872 |
| 251.89 | 192.01 | .3004 | .9871 |
| 259 | 194.01 | .3032 | .9871 |
| 266.19 | 196.01 | .306 | .987 |
| 273.45 | 198 | .3087 | .987 |

TABLE 4.- CONTINUED

| Altitude, km | Temperature, K | Conductivity, mcal/cm-deg sec | Conductivity ratio, mix/H ₂ |
|--------------|----------------|----------------------------------|---|
| 280.78 | 200 | 0.3115 | 0.9869 |
| 288.16 | 200 | .3115 | .9869 |
| 295.53 | 200 | .3115 | .9869 |
| 302.91 | 200 | .3115 | .9869 |
| 310.29 | 200 | .3115 | .9869 |
| 317.67 | 200 | .3115 | .9869 |
| 325.05 | 200 | .3115 | .9869 |
| 332.44 | 200 | .3115 | .9869 |
| 339.82 | 200 | .3115 | .9869 |
| 347.21 | 200 | .3115 | .9869 |
| 354.6 | 200 | .3115 | .9869 |
| 362.39 | 221.67 | .3412 | .9864 |
| 370.98 | 243.33 | .37 | .986 |
| 380.37 | 265 | .398 | .9856 |
| 390.57 | 286.67 | .4254 | .9853 |
| 401.58 | 308.33 | .4522 | .9851 |
| 413.39 | 330 | .4784 | .9848 |
| 426 | 351.67 | .504 | .9846 |
| 439.42 | 373.33 | .5292 | .9845 |
| 453.65 | 395 | .5539 | .9843 |
| 468.69 | 416.67 | .5783 | .9842 |
| 484.54 | 438.33 | .6022 | .9841 |
| 501.19 | 460 | .6258 | .984 |
| 518.66 | 481.67 | .649 | .9839 |
| 536.95 | 503.33 | .6719 | .9838 |
| 556.04 | 525 | .6946 | .9837 |
| 575.96 | 546.67 | .7169 | .9837 |
| 596.69 | 568.33 | .739 | .9836 |
| 618.24 | 590 | .7609 | .9835 |
| 640.6 | 611.67 | .7826 | .9835 |
| 663.79 | 633.33 | .8041 | .9834 |
| 687.8 | 655 | .8255 | .9834 |
| 712.64 | 676.67 | .8466 | .9833 |
| 738.3 | 698.33 | .8677 | .9833 |
| 764.79 | 720 | .8886 | .9832 |

TABLE 4.- CONCLUDED

| Altitude, km | Temperature, K | Conductivity, mcal/cm-deg sec | Conductivity ratio, mix/H ₂ |
|--------------|----------------|----------------------------------|---|
| 792.11 | 741.67 | 0.9094 | 0.9832 |
| 820.27 | 763.33 | .9301 | .9831 |
| 849.25 | 785 | .9508 | .9831 |
| 879.07 | 806.67 | .9714 | .983 |
| 909.73 | 828.33 | .9919 | .983 |
| 941.23 | 850 | 1.0123 | .9829 |

CONCLUDING REMARKS

The approximations used in estimating the viscosity of Jovian atmosphere mixtures are believed reliable within a few percent. The approximations used in estimating the thermal conductivities of Jovian atmosphere mixtures hold uncertainties of the order of 10 to 15%. In both cases, the mixture transport coefficients are almost a constant factor times the value for pure H₂, independent of temperature. This permits a quick, easy method of evaluating the appropriate values for the mixtures.

The transport coefficients included here are appropriate only for subsonic flow fields in the Jovian atmosphere. At high supersonic speeds, the H₂ dissociates and both the He and H atoms are electronically excited and ionized in flow following strong shock waves and in boundary-layer flow. In this case, the dissociation and ionization reactions promote very large peaks in the thermal conductivity over the range of pressures and temperatures where the reaction rates are significant. The ionization of the species also decreases viscosity by an order of magnitude or more, because the long-range coulomb forces prevent long mean-free-path momentum transfer in the gas. The characteristic dissociation temperature of H₂ (that is, the ratio of dissociation energy Q to the Boltzmann constant k) is 51,965 K, and the characteristic electronic excitation and ionization temperatures are several times larger. These reactions will not become important until temperatures are of the order of 0.1 the characteristic temperature. In other words, the mixture formulas used here could be applied to temperatures of several thousand degrees. However, in such a case, the simple formulas used for the viscosities and thermal conductivities of the pure H₂ and He would have to be modified; they were designed only to apply in the range from 100 to 1000 K.

Tables 5 and 6 are printouts of the BASIC language programs that have been used to calculate the coefficients of viscosity and thermal conductivity of H₂ - He mixtures. In the future, we may well find that other mixtures are appropriate for the Jovian atmosphere, in which case the appropriate values of the mole fractions x_1 (of H₂) and x_2 (of He) can be introduced as input to

TABLE 5.- BASIC PROGRAM FOR VISCOSITY OF JOVIAN MODEL ATMOSPHERES

```

10 PRINT "TABLE I - VISCOSITY OF JUPITER MODEL ATMOSPHERES, MICROPOISE"
20 PRINT
30 PRINT "T,DEG K","LIGHT ATM","NOMINAL ATM","HEAVY ATM"
40 PRINT
50 X1=.89      ! Mol fraction of H2 in nominal atmosphere model
60 X2=.11      ! Mol fraction of He in nominal atmosphere model
70 Y1=.78      ! Mol fraction of H2 in heavy atmosphere model
80 Y2=.22      ! Mol fraction of He in heavy atmosphere model
90 T=1E2       ! Initial temperature, deg K
100 FOR I=1 TO 19
140 N1=90.6*(T/300)^.6658
142 N1=N1/(1+4/T)
150 N2=191.6*(T/300)^.7176      ! Viscosity of H2, micropoise
152 N2=N2/(1-11.4/T)          ! Viscosity of He, micropoise
160 Q1=32.3*(1+4/T)*(300/T)^.1658      ! Cross section H2-H2, Ang2
170 Q2=21.5*(1-11.4/T)*(300/T)^.2176  ! Cross section He-He, Ang2
180 Q3=(Q1^.5+Q2^.5)/2
190 Q3=Q3*Q3                    ! Cross section H2-He, Ang2
200 L1=1+.7967*(X2/X1)*(Q3/Q1)      ! Ratio L(pure)/L(mix) for H2
210 L2=1+.5634*(X1/X2)*(Q3/Q2)      ! Ratio L(pure)/L(mix) for He
220 N=N1/L1+N2/L2      ! Viscosity of nominal atmosphere, micropoise
230 L3=1+.7967*(Y2/Y1)*(Q3/Q1)      ! Ratio L(pure)/L(mix) for H2
240 L4=1+.5634*(Y1/Y2)*(Q3/Q2)      ! Ratio L(pure)/L(mix) for He
250 N3=N1/L3+N2/L4      ! Viscosity of heavy atmosphere, micropoise
260 PRINT T,1E-2*INT(1E2*N1),1E-2*INT(1E2*N),1E-2*INT(1E2*N3)
270 T=T+50
280 NEXT I
999 END
! Next value of temperature, deg K

```

TABLE 6.- BASIC PROGRAM FOR CONDUCTIVITY OF JOVIAN MODEL ATMOSPHERES

```

10 PRINT "TABLE III - THERMAL CONDUCTIVITY OF JUPITER MODEL ATMOSPHERES"
11 PRINT "
12 PRINT "
13 PRINT "
14 PRINT "
15 PRINT "
16 PRINT "
17 PRINT "
18 PRINT "
19 PRINT "
20 PRINT
21 PRINT
22 PRINT
23 PRINT "T, DEG K", "LIGHT ATM", "NOMINAL ATM", "HEAVY ATM"
24 PRINT
25 PRINT
26 PRINT
27 PRINT
28 PRINT
29 PRINT
30 PRINT "T, DEG K", "LIGHT ATM", "NOMINAL ATM", "HEAVY ATM"
31 PRINT
32 PRINT
33 PRINT
34 PRINT
35 PRINT
36 PRINT
37 PRINT
38 PRINT
39 PRINT
40 PRINT
41 PRINT
42 PRINT
43 PRINT
44 PRINT
45 PRINT
46 PRINT
47 PRINT
48 PRINT
49 PRINT
50 X1=.89 ! Mol fraction of H2 in nominal atmosphere model
51 X2=.11 ! Mol fraction of He in nominal atmosphere model
52 Y1=.78 ! Mol fraction of H2 in heavy atmosphere model
53 Y2=.22 ! Mol fraction of He in heavy atmosphere model
54 T=100 ! Initial temperature, deg K
55 FOR I=1 TO 19
56 K1=.11*(T/300)^.6983/(1+49.4/T)
57 V=3079.5/T
58 V=2*V/(EXP(V)-EXP(-V))
59 K1=K1*(4.75+V*V) ! Conductivity of H2, millical/cm-deg-sec
60 K2=.3418*(T/300)^.7412/(1-13.74/T) ! Conductivity of He, mcal/cm-deg-sec
61 Q1=26.1*(1+49.4/T)*(300/T)^.1983 ! Cross section H2-H2, Ang2
62 Q2=5.96*(1-13.74/T)*(300/T)^.2412 ! Cross section He-He, Ang2
63 Q3=(Q1^.5+Q2^.5)/2
64 Q3=Q3*Q3 ! Cross section H2-He, Ang2
65 L1=1+.7698*(X2/X1)*(Q3/Q1) ! Ratio L(pure)/L(mix) for H2
66 L2=1+1.0887*(X1/X2)*(Q3/Q2) ! Ratio L(pure)/L(mix) for He
67 K=K1/L1+K2/L2 ! Thermal conductivity nominal atm, mcal/cm-deg-sec
68 L3=1+.7698*(Y2/Y1)*(Q3/Q1) ! Ratio L(pure)/L(mix) for H2
69 L4=1+1.0887*(Y1/Y2)*(Q3/Q2) ! Ratio L(pure)/L(mix) for He
70 K3=K1/L3+K2/L4 ! Thermal conductivity heavy atm, mcal/cm-deg-sec
71 PRINT T, 1E-4*INT(1E4*K1), 1E-4*INT(1E4*K), 1E-4*INT(1E4*K3)
72 T=T+50 ! Next value of temperature, deg K
73 NEXT I
74 END

```

the program. If sizable fractions of other gases were to be discovered in the Jovian atmosphere, they would have to be included, of course.

Amea Research Center
National Aeronautics and Space Administration
Moffett Field, California 94035, January 18, 1979

REFERENCES

1. Hirschfelder, Joseph O.; Curtiss, Charles F.; and Bird, R. Byron: Molecular Theory of Gases and Liquids. John Wiley and Sons, New York, 1954, p. 532.
2. Hansen, C. Frederick: Interpretation of Linear Approximations for the Viscosity of Gas Mixtures. Phys. Fluids, vol. 4, 1961, pp. 926-927.
3. Buddenberg, J. W.; and Wilke, C. R.: Calculation of Gas-Mixture Viscosities. Ind. & Eng. Chem., vol. 41, 1949, pp. 1345-1347.
4. Watson, J. T. R.: Viscosity of Gases in Metric Units, Dept. of Trade and Industry, National Engineering Laboratory, Her Majesty's Stationery Office, Edinburgh, 1972.
5. Kestin, J.: Viscosity of Gases. Ch. 2u, American Institute of Physics Handbook, Dwight E. Gray ed., Second ed., McGraw Hill Book Co., 1963.
6. Hilsenrath, Joseph; Beckett, C. W.; Benedict, L. F.; Hoge, H. J.; Masi, J. F.; Nuttall, R. L.; Touloukian, Y. S.; and Woolley, H. W.: Tables of Thermodynamic and Transport Properties of Air, Argon, Carbon Dioxide, Carbon Monoxide, Hydrogen, Nitrogen, Oxygen, and Steam. Pergamon Press, 1960. (Also NBS Circular 564, 1955).
7. Keyes, Frederick G.: The Heat Conductivity, Viscosity, Specific Heat, and Prandtl Number for Thirteen Gases. Project SQUID, MIT Tech Rept. 37, 1952.
8. Kestin, J. and Leidenfrost, W.: The Viscosity of Helium. Physica, vol. 25, 1959, pp. 537-555.
9. Powell, R. W.; Ho, C. Y.; and Liley, P. E.: Thermal Conductivity of Selected Materials. NSRDS - NBS Report 8, Nov. 25, 1966.
10. Powell, R. L.: Thermal Conductivity, Ch. 4g. American Institute of Physics Handbook, Dwight E. Gray Ed., Second ed., McGraw Hill Book Co., 1963.