

## General Disclaimer

### One or more of the Following Statements may affect this Document

- This document has been reproduced from the best copy furnished by the organizational source. It is being released in the interest of making available as much information as possible.
- This document may contain data, which exceeds the sheet parameters. It was furnished in this condition by the organizational source and is the best copy available.
- This document may contain tone-on-tone or color graphs, charts and/or pictures, which have been reproduced in black and white.
- This document is paginated as submitted by the original source.
- Portions of this document are not fully legible due to the historical nature of some of the material. However, it is the best reproduction available from the original submission.

LACIE-00504

JSC-11701

# LARGE AREA CROP INVENTORY EXPERIMENT (LACIE)

7.9-10148  
CR-158151

*"Data available under NASA sponsorship  
in the interest of easy and wide dis-  
semination of Earth Resources Survey  
Program information and without liability  
for any use made thereof."*



## NASA NOAA USDA

(E79-10142) LARGE AREA CROP INVENTORY  
EXPERIMENT (LACIE). A WHEAT YIELD MODEL FOR  
BRAZIL (National Oceanic and Atmospheric  
Administration) 16 p HC A02/MF A01 CSCI 02C

N79-18407

Unclas  
G3/43 00142

## A WHEAT YIELD MODEL FOR BRAZIL

# NASA

National Aeronautics and  
Space Administration


Lyndon B. Johnson Space Center  
Houston, Texas 77058

SEPTEMBER 1977

CENTER FOR CLIMATIC AND  
ENVIRONMENTAL ASSESSMENT

Technical Note 76-6

A Wheat Yield Model for Brazil

Authorized by:   
Assistant Deputy Project Manager  
for LACIE, NOAA

Center for Climatic and  
Environmental Assessment  
Federal Building, Room 116  
Columbia, Missouri 65201

December, 1976

CCEA TECHNICAL NOTE 76-6

A Wheat Yield Model for Brazil

C. M. Sakamoto<sup>1</sup>

INTRODUCTION

This particular study is in accord with the previously developed wheat models for other countries developed by CCEA. These models, known as the CCEA first generation type, are based on the following criteria: 1) it is a regression model; 2) it uses monthly temperature and precipitation data; 3) the model utilizes a heat stress term (temperature) and a moisture stress term (precipitation or some other moisture index); and 4) the trend term is described by a dummy variable or time (year) is used as a surrogate. Although this assumption is not totally adequate, this procedure must suffice until better quantitative inputs can be practically accomplished. The trend term consists of several technological factors that have changed yield over time. These include such factors as fertilizer, varietal improvements, use of better and bigger farm machinery, better pesticide management programs, etc. As additional information and techniques are gained by modelers, quantitative inputs by these factors, representing independent variables may be possible.

This study is based on a limited set of data, from 1961-1972. Since 1965, introduction of new varieties into Brazil, combined with increased acreage, has led to a rapid increase in wheat yield as well as in production. Consequently, the trend term needs to be monitored closely in the 1970's. Caution should be exercised by users of the model described herein, not

---

<sup>1</sup>Research Meteorologist, CCEA, Columbia, Missouri, December, 1976.

only because of the rapid potential technological changes in Brazil, but also because of the limited data upon which the model is based. However, as the reader will note, the selected variables are consistent with models developed in Brazil (see References) and do make agronomic sense.

#### DESCRIPTION OF WHEAT GROWING AREA

Wheat is grown primarily in the extreme southern region of Brazil (latitude 24° to 34°S) in the states of Rio Grande do Sul, Santa Catarina and Parana (Figure 1). Rio Grande do Sul, however, produces about 85 percent of the total wheat. Some wheat is also grown in Mato Grosso and Sao Paulo, but the amount is minor compared to that in Rio Grande do Sul.

Southern Brazil is atypical of wheat growing areas throughout the world. The area is relatively humid and warm. Furthermore, daylength at heading is between 12-14 hours. Consequently, locally adapted varieties that require low vernalization, high disease resistance and low photoperiod are desirable. The result is the introduction of varieties that are indifferent to daylength, have no or little response to vernalization, and are responsive to high temperatures from jointing to heading (da Mota and Acosta, 1973).

Depending on the variety, wheat is sown as early as April in Parana so that a double crop, including soybeans, can be planted in October (Beukenkamp, 1976). In Rio Grande do Sul, however, planting is usually in May or June while the major harvesting activity is in November and December.

By United States standards, farm sizes in southern Brazil are relatively small. According to Beukenkamp (1976), 42 percent of the farms average 5-6 hectares, 52 percent of the farms lie in the 10-100 hectares range and only 5 percent are over 300 hectares.

Annual precipitation ranges from a mean of 1300 mm to approximately 1700 mm in the wheat belt with three peaks in the year: June, October, and January (see Figure 2). Temperature ranges from about 11 to 13°C in July to about 20°C in November and December.

Frost at the critical heading and flowering period and summer hail are weather events that can reduce yield overnight in Brazil. Rust disease is a perennial problem but can be controlled by use of fungicide. It is estimated that yields can be boosted from an average of 1200 kilograms per hectare to 1600 kilograms per hectare by a program of fungicide application (Foreign Agriculture, 1976). However, the high cost of chemicals and equipment may limit its use.

In Parana, winter droughts have hampered yields. According to da Mota and Acosta (1973), initially low precipitation occurs 30 percent of the time in northern Parana. Parana is also where acreage as well as production of wheat is escalating. In 1975, it was estimated at 1.1 million hectares (Dalrymple, 1976). Introduction of Mexican semi-dwarf varieties will accelerate in 1976. Because of this variety's susceptibility to aluminum toxicity, it is estimated that the introduction of semi-dwarf varieties will be limited to certain areas of Brazil (Dalrymple, 1976).

#### SOURCE OF DATA

Meteorological data for Rio Grande do Sul were obtained from the publication Observacoes Meteorologicas (1974). Eight selected stations (Table 1) were used to find average temperature and precipitation for Rio Grande do Sul from 1962-1972. Yield data are shown in Table 2. As indicated earlier, data from 1962-1972 show that Rio Grande do Sul produces approximately 85 percent of Brazil's wheat, although this may change in the

mid-1970's. However, the relationship between Rio Grande do Sul and Brazil (country) yield suggests that the model applied to the state may also be used for the country (Figure 3).

#### WHEAT MODEL

The basic form of the model is:

$$\hat{Y} = \text{Trend} + f(\text{weather})$$

where:

$\hat{Y}$  = estimated yield (kgm/ha)

Trend = time trend where 1962=1, 1963=1, ..., 1965=1, 1966=2, ..., 1972=8, ...

$f(\text{weather})$  = function of weather elements in July through October.

After several iterations of model testing, it was determined that the following model with final truncation in October provided the best fit as well as being consistent with known climatic hazards in Brazil.

$$\begin{aligned} \hat{Y} = & \alpha + \beta_1 \text{Trend} + \beta_2 \text{ (July Temperature DFN)} \\ & + \beta_3 \text{ (September-October Weighted P-PET)} \\ & + \beta_4 \text{ (September-October Weighted P-PET)}^2 \end{aligned} \quad (1)$$

where:

P = precipitation (mm)

PET = potential evapotranspiration by the Thornthwaite procedure

$\alpha$  = a constant while  $\beta_1, \dots, \beta_4$  are coefficients of the selected variables.

Other truncated models, including their coefficients are shown in Table 3.

Weighted September-October precipitation minus potential evapotranspiration is determined as follows:

$$\frac{(P-PET)_{\text{Sep}} \times .50 + (P-PET)_{\text{Oct}}}{1.50} = \text{Weighted } (P-PET)_{\text{Sep-Oct}}$$

The variables selected in equation (1) are consistent with that reported by da Mota and Acosta (1973), da Mota and Wendt (1975), and da Mota (private communication, 1975). In the 1973 paper, July daily mean temperature and October rainfall were used in a regression model. However, as indicated by the authors, the introduction of improved varieties since 1965 may alter the correlation between temperature and yield. In the 1975 model, da Mota and Wendt (1975) used sunshine and relative humidity. Their equation is as follows:

$$Y = 5.99X_1 - 3.45X_2 - 20.72X_3 + 1871 \quad (2)$$

where:

Y = average yield (kgm/ha) for Rio Grande do Sul

X<sub>1</sub> = September sunshine

X<sub>2</sub> = August sunshine

X<sub>3</sub> = October relative humidity.

A simple test referred to as "jackknife" was performed on the model. In this procedure, one year was left out of the model and the remaining years were used to develop the coefficients. Yield for the year left out was estimated from the developed model. This was accomplished ten times so that the year left out proceeds sequentially from 1963 through 1972 and the model recalculated each time. The results of the test showing its value with the observed yield are shown in Figure 4.



SUMMARY

In spite of only 11 years of data, the wheat model developed for Rio Grande do Sul, Brazil appears to be reasonable. Greater variability in the data series can be expected with additional years. Revision of the model is essential with added data, particularly since 1972.

#### REFERENCES

- Beukenkamp, R. L. "Field Crops Take Lead in South Brazil's Rich Farmland," Foreign Agriculture, June 7, 1976, page 10.
- "Brazil Hopes for Rebound in Wheat," Foreign Agriculture, April 5, 1976, page 9.
- Dalrymple, D. G. "Development and Spread of High-Yielding Varieties of Wheat and Rice in the Less Developed Nations," Foreign Agricultural Economic Report No. 95, Economic Research Service, U.S. Department of Agriculture in Cooperation with the U.S. Agency for International Development.
- da Mota, F. S. Private Communication, October 16, 1975.
- da Mota, F. S. and M. J. C. Acosta. "Agrometeorology of the Wheat Crop in Brazil," paper presented at the Symposium on Agrometeorology of the Wheat Crop, Braunschweig, Federal Republic of Germany, October 22-27, 1973.
- da Mota, F. S. and W. Wendt. "Agrometeorological Wheat Yield Prediction in the State of Rio Grande do Sul, Brazil," Departamento Nacional de Meteorologia, 1975. (In Press)
- Equipe de Ecologia Agricola-IPACRGO, Observacoes Meteorologicas, Porto Alegre-RS-Brasil Janeiro de 1974, 179 pages.

Figure 1. Wheat Growing State of Brazil: Rio Grande do Sul, Santa Catarina and Parana







Figure 2. Water Balance of Passo Fundo, Typical of Wheat Zones in Southern Brazil (Source: da Mota and Acosta, 1973).

RELATIONSHIP OF RIO GRANDE DO SUL YIELD  
TO BRAZIL (COUNTRY) YIELD  
1962 - 1972



Source: Comissão Central de Levantamento Fiscalização das Safras Triticolas

Figure 3

"JACK-KNIFE" TEST  
RIO GRANDE DO SUL, BRAZIL  
1962 - 1972




Figure 4

Table 1. List of Stations and Locations Used for Meteorological Data  
in Rio Grande do Sul Wheat Yield Model

Station	Altitude (M)	Latitude	Longitude
Bage'	214	31°21'13'	54°06'21"
Encruzilhado do Sul	420	30°32'35"	52°31'20"
Julio de Castilhos	514	29°13'26"	53°40'45"
Rio Grande	5	32°01'44"	52°15'37"
Sao Borja	99	28°39'44"	56°00'15"
Tramandai	3	29°56'22"	50°30'12"
Veranopolis	705	28°56'14"	51°33'11"

Table 2 . Acreage, Production and Yield in Rio Grande do Sul, Brazil and Brazil (Country), 1962-1972\*

Harvest Year	Rio Grande do Sul			Brazil (Country)		
	Area (Ha)	Production (Qu)	Yield (Qu/Ha)	Area (Ha)	Production (Qu)	Yield (Qu/Ha)
1962	226,611.52	274,148.702	1.210	258,221.19	303,396.042	1.175
1963	218,877.30	103,591.763	.371	302,122.20	115,716.576	.383
1964	271,918.20	225,661.588	.830	300,542.60	250,452.634	.833
1965	325,390.60	236,951.433	.728	354,680.15	256,746.601	.724
1966	343,470.80	295,999.453	.862	384,960.26	333,516.981	.866
1967	487,687.66	339,628.440	.696	561,987.16	405,748.371	.721
1968	689,139.17	618,712.086	.897	845,693.69	765,076.711	.904
1969	1,044,731.06	1,090,107.567	1.043	1,299,518.52	1,303,426.759	1.004
1970	1,584,414.78	1,706,756.933	1.077	1,861,204.59	1,946,044.839	1.045
1971+	1,800,000.00	1,900,000.000	1.055	2,126,570.00	2,214,400.000	1.041
1972+	2,100,000.00	2,200,000.000	1.047	2,538,000.00	2,622,500.000	1.033

\*Source: Comissao Central Levantamento e Fiscalizacao das Safras Triticolas, Brazil.

+Estimated



