

General Disclaimer

One or more of the Following Statements may affect this Document

- This document has been reproduced from the best copy furnished by the organizational source. It is being released in the interest of making available as much information as possible.
- This document may contain data, which exceeds the sheet parameters. It was furnished in this condition by the organizational source and is the best copy available.
- This document may contain tone-on-tone or color graphs, charts and/or pictures, which have been reproduced in black and white.
- This document is paginated as submitted by the original source.
- Portions of this document are not fully legible due to the historical nature of some of the material. However, it is the best reproduction available from the original submission.

EFFECT OF 6-DAY HYPOKINESIA ON OXYGEN METABOLISM
INDICES IN ELDERLY AND SENILE SUBJECTS

L. A. Ivanov and P. A. Orlov

Translation of: "Vliyaniye 6-dnevnoy gipokinezii
na pokazateli kislorodnogo obmena v pozhilom i
starcheskom vozraste," Byulleten Eksperimental'noy
Biologii i Meditsiny, Vol. 76, No. 7, July 1973,
pp. 35-37.



(NASA-TM-75521) EFFECT OF 6-DAY HYPOKINESIA
ON OXYGEN METABOLISM INDICES IN ELDERLY AND
SENILE SUBJECTS (National Aeronautics and
Space Administration) 6 p HC A02/MF A01

N78-27753

CSCL 06S G3/52

Unclas
25725

NATIONAL AERONAUTICS AND SPACE ADMINISTRATION
WASHINGTON, D. C. 20546 JULY 1978

1. Report No. NASA TM-75521	2. Government Accession No.	3. Recipient's Catalog No.	
4. Title and Subtitle EFFECT OF 6-DAY HYPOKINESIA ON OXYGEN METABOLISM INDICES IN ELDERLY AND SENILE SUBJECTS		5. Report Date JULY 1978	6. Performing Organization Code
		8. Performing Organization Report No.	
7. Author(s) L. A. Ivanov and P. A. Orlov		10. Work Unit No.	
		11. Contract or Grant No. NASw-2791	
9. Performing Organization Name and Address SCITRAN Box 5456 Santa Barbara, CA 93108		13. Type of Report and Period Covered Translation	
		14. Sponsoring Agency Code	
12. Sponsoring Agency Name and Address National Aeronautics and Space Administration Washington, D.C. 20546			
15. Supplementary Notes Translation of: "Vliyaniye 6-dnevnoy gipokinezii na pokazateli kislородnogo obmena v pozhilom i starcheskom vozraste," Byulleten Eksperimental'noy Biologii i Meditsiny, Vol. 76, No. 7, July 1973, pp. 35-37.			
16. Abstract After a strict 6-day confinement to bed of elderly and senile subjects the oxygen supply of the subcutaneous cellular tissue was impaired, and the intensity of its tissue respiration was somewhat reduced. The vacat-oxygen of the blood and urine, the coefficient of incomplete oxidation, and the oxygen deficiency in the organism were increased. <p style="text-align: right;">ORIGINAL PAGE IS OF POOR QUALITY</p>			
17. Key Words (Selected by Author(s))		18. Distribution Statement Unclassified - Unlimited	
19. Security Classif. (of this report) Unclassified	20. Security Classif. (of this page) Unclassified	21. No. of Pages 6	22.

EFFECT OF 6-DAY HYPOKINESIA ON OXYGEN METABOLISM INDICES
IN ELDERLY AND SENILE SUBJECTS

L. A. Ivanov and P. A. Orlov

After a strict 6-day confinement to bed of elderly and senile subjects the oxygen supply of the subcutaneous cellular tissue was impaired, and the intensity of its tissue respiration was somewhat reduced. The vacat-oxygen of the blood and urine, the coefficient of incomplete oxidation, and the oxygen deficiency in the organism were increased. /35*

Hypokinesia is a significant factor in premature aging [5, 7, 9, 10]. Due to the role of oxygen deficiency in the development of senile shifts [4, 6, 8] of especial importance is the study of the hypodynamic effects on the oxygen metabolism indices and the oxidation-reduction processes in elderly and senile people.

This work studied the effect of hypokinesia on the oxygen absorption, its stress in the tissue, and the content of incompletely oxidized products in the organism of elderly and senile people. /36

Technique

Oxygen stress (PO_2) was determined in the subcutaneous cellular tissue of the left forearm by the polarographic method. The active cathode was an exposed platinum needle electrode 0.4 mm in diameter, while the comparison anode--silver chloride electrode superimposed on the distal third of the right shin. Measurements were made with a supply of constant voltage 0.7 v to the

*Numbers in the margin indicate pagination in the orginial foreign text.

electrodes. Graphic recording was made with the help of an automatic electron potentiometer EPP-09MZ through a photocompensation amplifier F 116/1. The results were expressed in millimeters of a mercury column by conducting electrode calibration after each study in physiological solutions with a certain oxygen pressure.

As functional tests permitting an evaluation of the oxygen supply to the subcutaneous cellular tissue and the intensity of its oxygen consumption 10-minute oxygen inhalation and pinching of the vessels in the limb for 10 minutes were used. A spiograph SG-1m was employed to measure out and record the oxygen load. Spiographic study also made it possible to determine oxygen absorption and its deficiency in the organism [3].

The content of incompletely oxidized products in the blood was determined according to Roman's method with correction for sugar according to Hagedorn-Iensen, and with correction for chlorides according to Levinson. The vacat-oxygen of the urine was determined by the Brin micromethod with correction for chlorides according to More, and the protein-free nitrogen of the urine--K'yel'dal's semimicromethod.

Twenty-one essentially healthy men were examined of elderly (60-74) and senile (75-89) age before and after a strict 6-day confinement to bed.

Results

After a course of strict confinement to bed for elderly and senile people the increase in PO_2 in the subcutaneous cellular tissue during oxygen inhalation was reduced from 38.2 ± 4.69 to 24.4 ± 3.21 mm Hg ($D < 0.02$), which reflects the exacerbation of the oxygen supply to the subcutaneous cellular tissue. There was a certain tendency for the latent period in the reduction of PO_2 in the subcutaneous cellular tissue to be lengthened in the transition from oxygen inhalation to air respiration (41.6 ± 4.24 sec. as compared to 34.4 ± 4.12 sec. before hypokinesia; $0.2 < D < 0.3$), as well as in the test with vessel constriction (11.9 ± 1.55 sec. as compared to 8.6 ± 0.92 sec.;

$0.05 \leq D \leq 0.1$.

The indicated parameters reflect the intensity of tissue respiration, and they increase with its reduction [2]. The reality of the noted trend towards reduction in oxygen consumption of the subcutaneous cellular tissue after hypokinesia is confirmed by the fact that despite the significant drop in PO_2 increase during oxygen inhalation under these conditions, the length of stabilization of the PO_2 level during the transition from oxygen respiration to air respiration essentially was not altered (respectively 483.6 ± 31.92 and 473.2 ± 27.39 sec.). In addition, despite the exacerbation in the oxygen supply to the subcutaneous cellular tissue, the amount of PO_2 in this tissue in the elderly and senile subjects after hypokinesia was practically preserved at the initial level (48.6 ± 2.62 mm Hg).

After the hypokinesia course oxygen absorption in the elderly and senile subjects did not change (214.2 ± 20.12 ml/min with initial 228.9 ± 8.54 ml/min).

EFFECT OF HYPOKINESIA ON OXIDATION-REDUCTION PROCESSES INDICES
IN ELDERLY AND SENILE SUBJECTS ($M \pm m$)

Index	Before hypokinesia	After hypokinesia	D
Vacat-oxygen of diurnal urine (in g)	9.08 ± 0.91	11.5 ± 0.52	< 0.05
Coefficient of incomplete oxidation	1.14 ± 0.034	1.36 ± 0.059	< 0.001
Vacat-oxygen of blood (in mg%)	233 ± 9.08	270 ± 9.71	0.01

In individuals of elderly and senile age after a 6-day bed confinement the level of vacat-oxygen increased in the blood and diurnal urine, and the coefficient of incomplete oxidation was augmented (see the table). The reliable increase in concentration of incompletely oxidized products in the blood and urine reflects the exacerbation in the oxidation-reduction processes under conditions of hypokinesia, which results in a less complete oxidation of the metabolites. An increase in the coefficient of incomplete oxidation indicates that the oxidation reactions are less complete.

As is known, excess oxygen absorption from hyperoxic mixtures as compared to oxygen absorption during air respiration ("oxygen deficit") is

137

governed by an accumulation of metabolites in the organism whose oxidation consumes an additional quantity of oxygen [1, 11]. This is why an increase in the "oxygen deficit" in the organism that is detected in elderly and senile people after a 6-day hypokinesia ($24.8 \pm 3.76\%$ versus $12.2 \pm 1.33\%$; $D < 0.003$) can be explained by the increase in the level of incompletely oxidized products detected under such conditions.

References

1. Dembo, A. G., Nedostatochnost' funktsii vneshnego dyzhaniya ["Deficiency in the Function of External Respiration"], Leningrad, 1957.
2. Korkushko, O. V. and L. A. Ivanov, Klin. med., No. 12 (1969), p. 54.
3. Korkushko, O. V. and L. A. Ivanov, Fiziol. zh. SSSR, No. 12 (1970), p. 1813.
4. Mil'man, M. S., in Starost' ["Old Age"], Kiev, 1939, p. 41.
5. Mogendovich, M. R., in Dvigatel'naya aktivnost' i starenie [Motor Activity and Aging], Kiev, 1969, p. 227.
6. Sirotinin, N. N., Klin. med., No. 8 (1960), p. 72.
7. Frol'kis, V. V. and I. V. Muravov, in Fizicheskaya kul'tura--istochnik dolgoletiya ["Athletics -- the Source of Longevity"], Moscow, 1965, p. 24.
8. Chebotarev, D. F., O. V. Korkushko, and L. A. Ivanov, in Starenie i fiziologicheskiye sistemy organizma ["Aging and the Physiological System of the Organism"], Kiev, 1969, p. 221.
9. Kraus, H. and W. Raab, Hypokinetic Disease, Springfield, 1961.
10. Le Compte, H. Z., Alternsforsch, No. 18 (1965), p. 79.
11. Uhlenbruck, P., Z. ges. exp. Med., No. 74 (1930), p. 1.