

General Disclaimer

One or more of the Following Statements may affect this Document


- This document has been reproduced from the best copy furnished by the organizational source. It is being released in the interest of making available as much information as possible.
- This document may contain data, which exceeds the sheet parameters. It was furnished in this condition by the organizational source and is the best copy available.
- This document may contain tone-on-tone or color graphs, charts and/or pictures, which have been reproduced in black and white.
- This document is paginated as submitted by the original source.
- Portions of this document are not fully legible due to the historical nature of some of the material. However, it is the best reproduction available from the original submission.



DEPARTMENT OF MATHEMATICS
UNIVERSITY OF HOUSTON

CR 151698
HOUSTON, TEXAS

A FIXED POINT THEOREM FOR
CERTAIN OPERATOR VALUED MAPS
BY
D.R. BROWN & M.J. O'MALLEY
REPORT #69 FEBRUARY 1978



(NASA-CR-151698) A FIXED POINT THEOREM FOR
CERTAIN OPERATOR VALUED MAPS (Houston Univ.)
9 p HC A02/MF A01 CSCL 12A

N78-22849

Unclas
G3/67 14087

PREPARED FOR
EARTH OBSERVATION DIVISION, JSC
UNDER
CONTRACT NAS-9-15000

HOUSTON, TEXAS 77004

A FIXED POINT THEOREM FOR CERTAIN OPERATOR VALUED MAPS

D.R. Brown & M.J. O'Malley
Department of Mathematics
University of Houston
Houston, Texas

Report #69

March 1978

A FIXED POINT THEOREM FOR CERTAIN OPERATOR VALUED MAPS

by D.R. Brown and M.J. O'Malley¹

1. Introduction. Let H be a real Hilbert space, and let $B_1(H)$ denote the space of symmetric, bounded operators on H which have numerical range in $[0,1]$, topologized by the strong operator topology (that is, the topology of point-wise convergence). It is well known [3], that if $T \in B_1(H)$, then there exists a unique $S \in B_1(H)$ such that $S^2 = T$. We represent S by $T^{\frac{1}{2}}$. The following theorem is due to John Neuberger [2].

Theorem A: Suppose $w \in H$, P is an orthogonal projection on H , and L is a (strongly) continuous function from H into $B_1(H)$. Let $Q_0 = P$, and set $Q_{n+1} = Q_n^{\frac{1}{2}} L(Q_n^{\frac{1}{2}} w) Q_n^{\frac{1}{2}}$, $n = 0, 1, 2, \dots$. Then $\{Q_n\}_{n=0}^{\infty}$ converges to an element $Q \in B_1(H)$ for which $z = Q^{\frac{1}{2}} w$ is a fixed point of P and a fixed point of L in the sense that $L(z)z = z$.

In this paper, under the same hypotheses as Theorem A, we develop a family of Neuberger-like results to find points $z \in H$ satisfying $L(z)z = z$ and $P(z) = z$. This family includes Neuberger's theorem and has the additional property that "most" of the sequences $\{Q_n\}$ converge to idempotent elements of $B_1(H)$. The limit operator of Theorem A need not be idempotent.

Such theorems as those above not only play a valuable role in the search for numerical solutions of partial differential equations, but are also useful, in the finite-dimensional case, in attacking the problem of determining the nonzero

¹Both authors received partial support under NASA contract NAS-9-15000.

fixed points of a function $\phi: \mathbb{R}^n \rightarrow \mathbb{R}^n$. In particular, if $x \in \mathbb{R}^n - \{0\}$, then x is a fixed point of ϕ if and only if $A(x)x = x$, where A is the matrix valued function defined by $A(x) = (||x||^{-2}) \cdot \phi(x) \cdot (x^T)$. In fact, it follows that this can occur if and only if $A(x)$ is a nonzero symmetric idempotent.

It is a pleasure to record our indebtedness to H.P. Decell for the remark immediately above, and to several other members of the University of Houston Mathematics Department, particularly Phillip Walker, for helpful conversations regarding the preparation of this paper.

2. Fixed Points of $L(z)$. Recall that an operator is positive if $\langle Ax, x \rangle \geq 0$ for all $x \in H$, where $\langle \cdot, \cdot \rangle$ is the inner product of H . We presume familiarity with the standard properties of positive operators as set forth, for example, in [3]. By invocation of the Spectral Theorem, or, alternately, by a sequential construction, it is possible to provide, for any $T \in B_1(H)$ and any positive integer n , a unique operator $T^{1/n} \in B_1(H)$ such that $(T^{1/n})^n = T$. This notion extends immediately to arbitrary positive rational powers of T by defining $T^{r/s} = (T^{1/s})^r$. Moreover, by again appealing to the Spectral Theorem, it follows that if $\{Q_j\}$ is a sequence in $B_1(H)$ converging strongly to Q , and t is an arbitrary positive rational number, then $\{Q_j^t\}$ converges strongly to Q^t . Finally, recall that the usual quasi-order defined for positive operators by $A \leq B$ if and only if $B - A$ is positive satisfies an additional anti-symmetry condition, to wit: if A and B are positive and commute, then $A \leq B$ and $B \leq A$ forces $A = B$.

Lemma 1. Let $Q \in B_1(H)$ and let α be a positive rational number other than 1. If $Q^\alpha = Q$, then $Q = Q^2$; that is, Q is an idempotent.

Proof: Let $\alpha = r/s$; the presumed equality is equivalent to $Q^r = Q^s$. Without loss of generality, assume $r < s$ and that r is the minimal positive power of Q which reoccurs in the sequence $\{Q^n\}$. From the fact that powers of an operator descend in the quasi-order mentioned above, together with the limited anti-symmetry of this relation, it follows that $Q^t = Q^r$ for all integral t between r and s . From $Q^r = Q^{r+1}$, it follows that $Q^t = Q^r$ for all $t \geq r$. If r is odd, then $(Q^{(r+1)/2})^2 = Q^{r+1} = Q^{2r} = (Q^r)^2$. By uniqueness of square roots, $Q^r = Q^{(r+1)/2}$, whence $r = (r+1)/2$ and $r = 1$. If r is even, then $(Q^{r/2})^2 = Q^r = (Q^r)^2$, whence $r = r/2$, which is impossible for positive r . Thus $r = 1$ and $Q = Q^2$.

We are now ready to prove our

Theorem 2. Let $w \in H$, let P be an orthogonal projection on H , and let $L: H \rightarrow B_1(H)$ be strongly continuous. Let α, β be positive rational numbers with $\alpha \in [\frac{1}{2}, \infty)$. Set $Q_0 = P$, and let $Q_{n+1}^\alpha = Q_n^\alpha L(Q_n^\beta w) Q_n^\alpha$, $n = 0, 1, 2, \dots$. Then $\{Q_n\}_{n=0}^\infty$ is a decreasing sequence of elements of $B_1(H)$ which converge to an element $Q \in B_1(H)$ such that

(1) if $\alpha > \frac{1}{2}$, then Q is idempotent and $z = Qw$ satisfies

$L(z)z = z$, and $Pz = z$, and

(2) if $\alpha = \frac{1}{2}$ and $\beta \geq \frac{1}{2}$, then $z = Q^\beta w$ satisfies $L(z)z = z$ and

$Pz = z$.

Proof: Fix $\alpha \geq \frac{1}{2}$ and $\beta > 0$. Since $Q_0 = P \in B_1(H)$ and the range of L

is in $B_1(H)$, it follows inductively that $Q_n \in B_1(H)$ for all n . Since $2\alpha \geq 1$, $Q_n^{2\alpha} \leq Q_n$; moreover, $\langle (Q_n^{2\alpha} - Q_{n+1})x, x \rangle = \langle (Q_n^{2\alpha} - Q_n^\alpha L(Q_n^\beta w) Q_n^\alpha)x, x \rangle = \langle Q_n^\alpha (I - L(Q_n^\beta w)) Q_n^\alpha x, x \rangle = \langle (I - L(Q_n^\beta w)) Q_n^\alpha x, Q_n^\alpha x \rangle$. Thus, since $I - L(Q_n^\beta w) \geq 0$, it follows that $Q_{n+1} \leq Q_n^{2\alpha}$. Hence we have

$$(*) \quad Q_{n+1} \leq Q_n^{2\alpha} \leq Q_n, \quad n = 0, 1, 2, \dots$$

In particular, the sequence $\{Q_n\}$ is monotonically decreasing in the (operator) interval from 0 to I . Thus we have by [3, p.318] that the sequence $\{Q_n\}$ converges strongly to an element $Q \in B_1(H)$, whence $\{Q_n^\alpha\}$ converges to Q^α and $\{Q_n^\beta\}$ converges to Q^β . Since L is continuous and operator multiplication is jointly continuous in the strong topology on $B_1(H)$, we have by uniqueness of limits that $Q = Q^\alpha L(Q^\beta w) Q^\alpha$. Also, from (*) and the closed graph of the relation \leq , we have $Q \leq Q^{2\alpha} \leq Q$. Thus, since Q and $Q^{2\alpha}$ commute, we have that $Q = Q^{2\alpha}$. Moreover, since $P = Q_0$, we have $PQ_n = Q_n$, whence $PQ_n^\gamma = Q_n^\gamma$ for all positive rational γ .

(i) Suppose $\alpha > \frac{1}{2}$. By lemma 1, $Q = Q^2$, from which it follows that $Q = Q^\gamma$ for all positive rational γ , and, in particular, $Q = QL(Qw)Q$.

Let $z = Qw$, and fix $x \in H$. Then $\langle Qx, x \rangle = \langle QL(z)Qx, x \rangle = \langle L(z)Qx, Qx \rangle$, and since $Q^2 = Q$, it follows that $0 = \langle Qx, Qx \rangle - \langle L(z)Qx, Qx \rangle = \langle (I - L(z))Qx, Qx \rangle$. Therefore, since $I - L(z)$ and hence $(I - L(z))^{\frac{1}{2}}$ belong to $B_1(H)$, we have that $Q = L(z)Q$. In particular, $z = Qw = L(z)Qw = L(z)z$.

(ii) Suppose $\alpha = \frac{1}{2}$, $\beta \geq \frac{1}{2}$. Let $z = Q^\beta w$; then $Q = Q^{\frac{1}{2}} L(z) Q^{\frac{1}{2}}$ from which $\langle Qx, x \rangle = \langle Q^{\frac{1}{2}} L(z) Q^{\frac{1}{2}} x, x \rangle = \langle L(z) Q^{\frac{1}{2}} x, Q^{\frac{1}{2}} x \rangle$. Since $\langle Qx, x \rangle = \langle Q^{\frac{1}{2}} x, Q^{\frac{1}{2}} x \rangle$ also, we have $0 = \langle Q^{\frac{1}{2}} x - L(z) Q^{\frac{1}{2}} x, Q^{\frac{1}{2}} x \rangle = \langle (I - L(z)) Q^{\frac{1}{2}} x, Q^{\frac{1}{2}} x \rangle$. Now, as in (i), it follows

that $Q^{\frac{1}{2}} = L(z)Q^{\frac{1}{2}}$. In particular, $z = Q^{\beta}w = Q^{\frac{1}{2}}Q^{\beta-\frac{1}{2}}w = L(z)Q^{\frac{1}{2}}Q^{\beta-\frac{1}{2}}w = L(z)Q^{\beta}w = L(z)z$. That $Pz = z$ in both cases is obvious from the fact that $PQ^{\gamma} = Q^{\gamma}$ for all positive rational γ . This completes the proof.

Given a nonzero element $z \in H$ such that $L(z)z = z$, it is reasonable to ask if our sequences are able to produce z . We note now that, by proper selection of w and P , z is attainable from each of our sequences. Specifically, if α and β are fixed as in the theorem, then let $w = z$ and let P be the orthogonal projection of H onto the line through z . From the construction of the sequence $\{Q_n\}$, $Q_1 = PL(z)P$, whence $Q_1 = P$. It follows immediately that $Q_n = P$ for all n and thus $Q = P$. Hence $z = Qw = Pw$ (or $z = Q^{\beta}w = P^{\beta}w = Pw$) is the fixed point yielded by our theorem.

While it is not reasonable to expect the practitioner to guess P so accurately, these remarks do attach the virtue of theoretical completeness to these processes.

3. Examples. (1) Suppose that $\alpha = \frac{1}{2}$ and that $\gamma, \delta \in [\frac{1}{2}, \infty)$ such that neither of γ, δ is an integral multiple of the other. We show that for fixed $w \in H$ and P , the Q and z obtained by using γ for β need not be the same as those obtained by using δ for β . Moreover, the limit operator Q in this case need not be an idempotent, although it can be one. Assume $\delta < \gamma$. Let k be the least positive integer such that $\gamma < k\delta$. Note $2 \leq k$ and $(k-1)\delta < \gamma$. Let a be any number in the interval $(0,1)$. Then

$$a^{k\delta} < a^{\gamma} < a^{(k-1)\delta} \leq a^{\delta}.$$

Define $L:R \rightarrow [0,1]$ by

$$L(x) = \begin{cases} 1, & x \leq a^\gamma \\ [(1-a)/(a^\gamma - a^{(k-1)\delta})] \cdot (x - a^\gamma) + 1, & a^\gamma \leq x \leq a^{(k-1)\delta} \\ a, & a^{(k-1)\delta} \leq x. \end{cases}$$

Set $P = 1$, $w = 1$. Using γ for β in the theorem yields $Q_0 = 1$ and $Q_1 = a$. Inductively, $Q_n = a$, so that $Q = a$. Hence $z = Q^\gamma w = a^\gamma \cdot 1 = a^\gamma$ in this case. On the other hand, using δ for β gives $Q_0 = 1$, $Q_1 = a$, but $Q_2 = a^2, \dots, Q_k = a^k$. Moreover, $Q_n = a^k$ for $n \geq k$, hence $Q = a^k$ and $z = Q^\delta w = a^{k\delta} \cdot 1 = a^{k\delta}$. By the choices of a and k , the exponents γ and δ yield distinct operators and distinct fixed points. Moreover, neither of the limit operators determined by γ and δ is idempotent.

(2) Suppose that $\alpha > \frac{1}{2}$, so that any limiting Q obtained through the theorem is idempotent. We show for fixed $w \in H$ and P , that the resulting limit idempotents may vary with the choice of β , as may the fixed points determined in this manner. To this end, let $\alpha = 1$ in the theorem. Let $L:R^3 \rightarrow B_1(R^3)$ be as follows: all image matrices are diagonal, where $\begin{pmatrix} x & 0 & 0 \\ 0 & y & 0 \\ 0 & 0 & z \end{pmatrix}$ will be represented as $\text{diag}(x,y,z)$. We require $L(1,1,1) = \text{diag}(1, \frac{1}{2}, 1)$, $L(1, \frac{1}{2}, 1) = \text{diag}(1, \frac{1}{2}, \frac{1}{2})$, $L(1, \frac{1}{2}, \frac{1}{2}) = \text{diag}(\frac{1}{2}, \frac{1}{2}, 1)$, $L(1,y,z) = \text{diag}(1,y,z)$ for $(y,z) \in [0, \frac{1}{2}] \times [0, \frac{1}{2}]$, and $L(x,y,1) = \text{diag}(x,y,1)$ for $(x,y) \in [0, \frac{1}{2}] \times [0, \frac{1}{2}]$. The extension theorem of Tietze (c.f. [1]) permits a continuous extension of L to all of R^3 into the diagonal matrices whose entries are in the interval $[0,1]$. Let $P = I_3$, the identity operator, and let w be the vector $(1,1,1)$. If $\beta = \frac{1}{2}$, a brief examination of the defining sequence of Q_n 's in Theorem 2

shows that the limit idempotent $Q = \text{diag}(1,0,0)$, and $z = Qw = (1,0,0)$. On the other hand, if $\beta = 1$, then limit $Q = \text{diag}(0,0,1)$, and $z = (0,0,1)$.

(3) With notation as in (2), suppose $\beta = 1$ is fixed. We show for fixed $w \in H$ and P , that the resulting limit idempotents may vary with α , as may the fixed points determined in this manner. Letting $P = I_3$ and $w = (1,1,1)$ as in (2), we require this time that $L(1,1,1) = L(1, \frac{1}{2}, 1) = \text{diag}(1, \frac{1}{2}, 1)$, $L(1, 1/8, 1) = L(1, 0, 0) = \text{diag}(1, 0, 0)$, and $L(1, 1/32, 1) = L(0, 0, 1) = \text{diag}(0, 0, 1)$. Extending as before, we have a continuous L defined on R^3 into the diagonal matrices with entries in $[0,1]$. For any choice of α , $Q_1 = \text{diag}(1, \frac{1}{2}, 1)$. If $\alpha = 1$, $Q_2 = \text{diag}(1, 1/8, 1)$, $Q_3 = Q_n = Q = \text{diag}(1, 0, 0)$, $z = (1, 0, 0)$. On the other hand, if $\alpha = 2$, then $Q_2 = \text{diag}(1, 1/32, 1)$, $Q_3 = Q_n = Q = \text{diag}(0, 0, 1)$, $z = (0, 0, 1)$.

It is easy to see that a slightly more complicated definition of L would yield a single example incorporating the features of all three prior illustrations.

References

1. Kelley, John L., General Topology, Van Nostrand, New York, 1955.
2. Neuberger, John, Projection Methods for Linear and Nonlinear Systems of Partial Differential Equations. Springer-Verlag Lecture Notes (to appear).
3. Schechter, Martin, Principles of Functional Analysis, Academic Press, New York, 1971.