

## General Disclaimer

### One or more of the Following Statements may affect this Document

- This document has been reproduced from the best copy furnished by the organizational source. It is being released in the interest of making available as much information as possible.
- This document may contain data, which exceeds the sheet parameters. It was furnished in this condition by the organizational source and is the best copy available.
- This document may contain tone-on-tone or color graphs, charts and/or pictures, which have been reproduced in black and white.
- This document is paginated as submitted by the original source.
- Portions of this document are not fully legible due to the historical nature of some of the material. However, it is the best reproduction available from the original submission.

INORGANIC CHEMISTRY--DIRECT SYNTHESSES FROM  
PURE LIQUID  $\text{SO}_2$  AND FROM TRIVALENT AND PENTAVALENT  
NITROGEN DERIVATIVES

B. Vandorpe and J. Heubel

(NASA-TM-75171) INORGANIC CHEMISTRY: N78-11214  
DIRECT SYNTHESSES FROM PURE LIQUID  $\text{SO}_2$  AND  
FROM TRIVALENT AND PENTAVALENT NITROGEN  
DERIVATIVES (National Aeronautics and Space Administration) 7 p HC A02/MF A01 CSCL 07D G3/25 Unclas  
49799

Translation of "Chimie Minerale--Synthèses directes à partir  
de  $\text{SO}_2$  pur liquide et de dérivés azotes tri et pentavalents,"  
Comptes Rendues Hbdomaires des Séances de l'Academie des  
Sciences, Paris, Vol. 260 (June 21, 1968) Group  
8, pp. 6619-6621



1. Report No. NASA TM-75171	2. Government Accession No.	3. Recipient's Catalog No.	
4. Title and Subtitle INORGANIC CHEMISTRY--DIRECT SYNTHESES FROM PURE LIQUID SO <sub>3</sub> AND FROM TRIVALENT AND PENTAVALENT NITROGEN DERIVATIVES		5. Report Date September 1977	6. Performing Organization Code
		8. Performing Organization Report No.	
7. Author(s) Bernard Vandorpe and Joseph Heubel, Laboratory of Inorganic Chemistry, Faculty of Sciences, 103, rue B.- Delespaul, Lille, Nord		10. Work Unit No.	
		11. Contract or Grant No. NASW-2790	
9. Performing Organization Name and Address Leo Kanner Associates Redwood City, California 94063		13. Type of Report and Period Covered Translation	
		14. Sponsoring Agency Code	
12. Sponsoring Agency Name and Address National Aeronautics and Space Adminis- tration, Washington, D.C. 20546			
15. Supplementary Notes Translation of "Chimie Minerale--Synthèses directes à partir de SO <sub>3</sub> pur liquide et de dérivés azotes tri et pentavalents," Comptes Rendues Hebdomaires des Seances de l'Academie des Sciences, Paris, Vol. 260 (June 21, 1968) Group 8, pp. 6619-6621			
16. Abstract From pure liquid SO <sub>3</sub> the authors carried out direct synthesis reactions with N <sub>2</sub> O <sub>5</sub> , NO <sub>2</sub> Cl, NOCl which yielded N <sub>2</sub> O <sub>5</sub> 4SO <sub>3</sub> , 3SO <sub>3</sub> , 2SO <sub>3</sub> -NO <sub>2</sub> Cl2SO <sub>3</sub> -NOCl2SO <sub>3</sub> and NOClSO <sub>3</sub> , the latter being obtained for the first time in the pure state. In all cases the crystallized product was obtained first by separating the constituents of the mixture and then going through a single viscous liquid phase.			
17. Key Words (Selected by Author(s))		18. Distribution Statement  Unclassified-Unlimited	
19. Security Classif. (of this report) Unclassified	20. Security Classif. (of this page) Unclassified	21. No. of Pages 5	22. Price

INORGANIC CHEMISTRY--DIRECT SYNTHESSES FROM  
PURE LIQUID  $\text{SO}_3$  AND FROM TRIVALENT AND PENTAVALENT  
NITROGEN DERIVATIVES\*

B. Vandorpe and J. Heubel  
Laboratory of Inorganic Chemistry, Faculty of Sciences,  
103, rue B.-Delespaul, Lille, Nord

Nitryl polysulfates are usually obtained at low /6619\*\*  
temperature from  $\text{SO}_3$  and  $\text{N}_2\text{O}_5$  in solution in various solvents:  
nitromethane [1],  $\text{POCl}_3$  [2],  $\text{CCl}_4$  [3], liquid  $\text{SO}_2$  [4]. To  
our knowledge, synthesis in the absence of solvent has never  
been attempted, or was attempted under such conditions that  
the formation of polysulfates was impossible [3]. One might  
think that the authors relied on solvents because of the  
difficulty of obtaining stable liquid  $\text{SO}_3$  and also because the  
reaction is exothermic.

The fact that we had at our disposal a sure method of  
preparation and a proven apparatus encouraged us to attempt  
this synthesis and then generalize it to other nitrogen  
derivatives.

$\text{SO}_3$  is obtained according to a method specified by J.  
Bernard [6]: reaction of oleum on  $\text{P}_2\text{O}_5$ , distillation of the  
raw product and passing over  $\text{P}_2\text{O}_5$  with the formation of  
 $\text{P}_4\text{O}_{10}\text{SO}_3$ . The thermal dissociation of the latter yields pure  
 $\text{SO}_3$ , the head fraction of which serves to wash the reactor.  
This fraction is passed off into an ampule which is removed.  
The reactor has a ground inlet on the side, connected on the  
one hand to the apparatus which produces the  $\text{N}_2\text{O}_5$  and on the  
other hand to a feed pipe of dry nitrogen. The reactor

\*Paper by Bernard Vandorpe and Joseph Heubel, read by Georges  
Champetier, at the June 14, 1965 meeting of the Academy of  
Sciences, Paris.

\*\*Numbers in the margin indicate pagination in the foreign text.

remains in place under a current of dry nitrogen and is constantly shielded by columns of  $P_2O_5$  until the completion of the operations. By means of a by-pass system the apparatus can be purged and the head fractions of  $N_2O_5$  can be removed. The latter is produced according to a method described earlier [7] which has been improved so as to produce an hourly yield of about 20 grams of pure product.

In the reactor containing  $SO_3$ ,  $N_2O_5$  vapor is caused to be given off at ambient temperature which is carried away by the dry nitrogen. A slight increase in temperature is observed which can be compensated for -- without its being absolutely necessary -- by immersing the reactor in cold water. The liquid becomes cloudy and then the constituents of the mixture separate.

The composition of the lower layer remains constant between  $-30^\circ$  and  $+20^\circ$ . It corresponds approximately to the formula  $N_2O_5 \cdot 7SO_3$ . Following disappearance of the upper layer, the viscosity of the liquid increases sharply, then solid particles form which gradually take over the reactor. When the gas no longer passes easily some pure nitrogen is introduced and the temperature is raised to  $60^\circ$  for about 48 hours in order to remove the excess of  $SO_3$ . /662

The solid obtained sometimes corresponds to the formula  $N_2O_5 \cdot 4SO_3$ , sometimes to  $N_2O_5 \cdot 3SO_3$ . This is explained by the large degree of viscosity of the liquid prior to crystallization. This viscosity promotes the supercooling of  $N_2O_5 \cdot 4SO_3$ , thus the passage of the gas and the concentration of  $N_2O_5$ . In fact, one more often obtains  $N_2O_5 \cdot 3SO_3$  than  $N_2O_5 \cdot 4SO_3$ . In order to be sure of obtaining the latter it is necessary to wait for spontaneous crystallization by supercooling or by feeding which requires a special technique to avoid any traces of water.


The method thus described is limited to the preparation of trisulfates and tetrasulfates, but it never enabled us to obtain directly  $N_2O_3 \cdot 2SO_3$ .

$N_2O_5 \cdot 4SO_3$ , ground and subjected to the action of  $N_2O_5$  gas at ambient temperature, fixes the latter and is gradually transformed into  $N_2O_3 \cdot 3SO_3$ .  $N_2O_5 \cdot 3SO_3$  submitted to the same treatment does not undergo any significant increase in weight.

By contrast, by grinding  $N_2O_5 \cdot 3SO_3$  or  $N_2O_5 \cdot 4SO_3$  with solid  $N_2O_5$  in a glove box or on a heated plate under a current of dry nitrogen, one can increase the  $N_2O_5$  concentration of the initial product, ending up with the final limit formula  $N_2O_5 \cdot 2, 2SO_3$ . This mixture of two polysulfates is stable up to  $125^\circ$ . It is not surprising that one reaches a limit formula without being able to obtain  $N_2O_5 \cdot 2SO_3$ , since a solid-solid action is involved without the release of gas. It is possible, on the other hand, to obtain  $N_2O_5 \cdot 2SO_3$  in the pure state by causing  $N_2O_5$  to react on  $NO_2Cl \cdot 2SO_3$  or  $NOClSO_3$ , the direct synthesis of which is mentioned below. The reactions seem to proceed as follows:



and



With cooled  $NO_2Cl$  carried away by an inert gas the reaction on  $SO_3$  occurs with a slight increase in temperature and is analogous to that of  $N_2O_5$ : formation of turbidity, followed by separation of the constituents, the lower layer corresponding approximately to the formula  $NO_2Cl \cdot 5SO_3$ , increase in viscosity and crystallization of a solid which, after removal of the excess  $SO_3$  at  $60^\circ$ , corresponds to the formula  $NO_2Cl \cdot 2SO_3$ . The S/N ratio = 2.01; N/Cl = 1.01 as against S/N = 2.16 and N/Cl = 1.075 respectively in the violent reaction of  $SO_3$  on  $NO_2Cl$  maintained at  $-80^\circ$  [8].

The same type of reaction is found again with NOCl carried away by inert gas: turbidity, separation of the constituents, lower layer corresponding to  $\text{NOCl}_5\text{SO}_3$ , formation of a solid and obtainment of  $\text{NOClSO}_3$  after heating the reactor to  $70^\circ$ . This compound is particularly interesting, since it is the only one among the derivatives described which corresponds to  $S/N = 1$ . Although having the formula  $2\text{SO}_3$ ,  $\text{NO}_2\text{Cl}$  it seems to have been obtained in a very impure form by Weber in 1864 [9]. Since that time attention has no longer been drawn to it, except by Seel [10] who doubts that it exists. /6621

By causing an excess of  $\text{SO}_3$  to react on NOCl cooled to  $-50^\circ$  one can obtain  $\text{NOCl}_2\text{SO}_3$  pointed out by Seel [10] and prepared by Weinreich in solution in liquid  $\text{SO}_2$  [4, 11].

Tests in progress with  $\text{SO}_3$  and  $\text{N}_2\text{O}_3$  and NO show that all of these synthesis reactions (gas + liquid  $\text{SO}_3$ ) proceed according to the same plan: the synthesis by means of separation of the constituents in the mixture which in turn, when one of the liquid phases disappears, is followed by an increase in viscosity and then crystallization of a solid. Quantitative transformation from liquid into solid presents a certain degree of experimental difficulty.

These syntheses have the advantage of doing without any solvent, taking place at ambient temperature and easily lending themselves to the preparation of large quantities of product.

## REFERENCES

1. Goddard and Ingold, J. Chem. Soc., 2559-2575 (1950).
2. Gerding and Eriks, Rec. Trav. Chim. 71, 773-778 (1952).
3. Pictet and Karl, Bull. Soc. chim. Fr. 3(4), 1114 (1908).
4. Weinreich, Bull. Soc. chim. Fr., 2820 (1963).
5. Pascard, Mém. Serv. Chim. Etat (Paris) 38(4), 305-310 (1953).
6. Bernard, Thesis, Paris, 1957.
7. J. Heubel, Comptes rendus 252, 3268 (1961).
8. Batey and Sisler, J. Amer. Chem. Soc. 74, 3408-3410 (1952).
9. Weber, Pogg. Ann. 123, 333-346 (1864).
10. Seel, Angew. Chem. 68, 272-284 (273) (1956).
11. Ultimately we hope to be able to specify the composition of the various compounds pointed out. For the time being we are using dualistic formulas.