

## General Disclaimer

### One or more of the Following Statements may affect this Document

- This document has been reproduced from the best copy furnished by the organizational source. It is being released in the interest of making available as much information as possible.
- This document may contain data, which exceeds the sheet parameters. It was furnished in this condition by the organizational source and is the best copy available.
- This document may contain tone-on-tone or color graphs, charts and/or pictures, which have been reproduced in black and white.
- This document is paginated as submitted by the original source.
- Portions of this document are not fully legible due to the historical nature of some of the material. However, it is the best reproduction available from the original submission.

505

Infrared Spectra of Molecules and Materials  
of Astrophysical Interest

Grant NGL-41-002-003  
Quarterly Progress Report  
Report Number 36  
June 1976

Office of Grants and Research Contracts  
Office of Space Science and Applications  
National Aeronautics and Space Administration  
Washington, D. C. 20546

(NASA-CR-148292) INFRARED SPECTRA OF  
MOLECULES AND MATERIALS OF ASTROPHYSICAL  
INTEREST Quarterly Progress Report, 15 Mar.  
- 15 Jun. 1976 (South Carolina Univ.) 85 p  
HC \$5.00

N76-28132

CSCI 03E G3/90

Unclas  
44482

(Prepared under Grant NGL-41-002-003 by the  
University of South Carolina, Columbia, South Carolina 29208

Principal Investigator: James R. Durig, Professor of Chemistry

Period Covered: 15 March 1976 to 15 June 1976



## SUMMARY OF PROGRESS

We have been studying the vibrational spectra from 4000 to  $33\text{ cm}^{-1}$  of several molecules which may be present in the atmosphere of the Jovian planets or exist in outer space. These studies have been made to provide vibrational frequencies which can be used to: (1) determine the composition of the cloud covers of several of the planets, (2) provide structural information under favorable circumstances, (3) provide necessary data from which accurate thermodynamic data can be calculated, and (4) furnish information as to the nature of the potential energy function of the molecules and forces acting within them.

Some of the molecules which we have studied can be produced photochemically from methane, ammonia, and hydrogen sulfide which are thought to be constituents of the planets with reducing atmospheres. Some of the compounds will polymerize under ultraviolet radiation and drop out of the atmospheres. However, planets with a hot base, like that of Jupiter, may rebuild molecules destroyed photochemically. Therefore, we have used these criteria in selecting the compounds which we have studied.

Gerald P. Kuiper<sup>1</sup> has pointed out that the Jovian atmosphere is expected to contain  $\text{H}_2$ , He,  $\text{N}_2$ ,  $\text{H}_2\text{O}$ ,  $\text{NH}_3$ ,  $\text{CH}_4$ , Ar and possibly  $\text{SiH}_4$ . He has also listed a number of other gases that should be considered because they are composed of fairly abundant atomic species and have boiling points below  $120^\circ\text{C}$  (see Table 8, pg. 349-350 of reference 1). He has also pointed out that until more is known about the atmospheres of the planets, it is useful to keep a fairly large number of possible constituents in mind in planning further spectroscopic work.

In our initial work on the vibrational spectra of molecules of astrophysical interest, we studied hydrazine,<sup>2</sup> methylamine,<sup>3</sup> as well as several substituted hydrazines.<sup>4-8</sup> Recently both ethane and acetylene have been found in the atmosphere of Jupiter.<sup>9</sup> It is expected that substituted ethanes will also be eventually found in some of the planetary atmospheres. In fact, ethanol has been found in the Sagittarius B12 cloud of dust and gas which is near the center of the Milky Way.<sup>10</sup> In addition, molecules such as acetaldehyde ( $\text{CH}_3\text{CHO}$ ), methanol ( $\text{CH}_3\text{OH}$ ), dimethylether ( $\text{CH}_3$ )<sub>2</sub>O, formic acid ( $\text{HCOOH}$ ), have been identified in outer space. As a continuation of our earlier studies on the low frequency torsional motions of dimethylether<sup>11</sup> we have recorded the high resolution microwave spectra of this molecule along with the two isotopic derivatives,  $\text{CH}_3\text{OCD}_3$  and  $(\text{CD}_3)_2\text{O}$  and reported this study in the last progress report. We have now completed the relatively high resolution far infrared study along with a study of the low frequency Raman spectra of the gas.

Several years ago the gas phase far infrared spectra of molecules of the type  $(\text{CH}_3)_2\text{X}$  and  $(\text{CD}_3)_2\text{X}$  were published. Some of these spectra showed a very complex pattern in the region where the methyl torsional transitions are expected. Tuazon and Fateley<sup>12</sup> reported in 1971 the far infrared spectrum of  $\text{CD}_3\text{OCH}_3$  along with the already measured species  $(\text{CH}_3)_2\text{O}$  and  $(\text{CD}_3)_2\text{O}$ . The spectrum of this compound is similarly complex. These authors took the most likely band in the spectra to obtain an estimate of the barrier height, but detailed interpretation of the complex features were not presented. To our knowledge, only one Raman spectrum of gaseous  $\text{CH}_3\text{OCH}_3$  has been reported. In 1957, Taylor and Vidale<sup>13</sup> reported two Raman bands at 480 and 450  $\text{cm}^{-1}$  as overtones of both torsional modes.

In order to get better barrier information on these isotopic species of dimethylether and eventually to explain the complicated structure of their

far infrared spectra, the Raman spectra of these isotopes were measured in the gas phase on a Cary Model 82 instrument with a Spectra Physics 171 argon ion laser. For the same purpose, the far infrared spectra were remeasured with  $0.5 \text{ cm}^{-1}$  resolution with a Digilab FTS 15B interferometer. In Figure 1 are shown the far infrared spectrum of the normal isotope. The band at  $240 \text{ cm}^{-1}$  is clearly asymmetric, and the band at  $223 \text{ cm}^{-1}$  is split into at least three components at  $223.2$ ,  $224.8$  and  $226.0 \text{ cm}^{-1}$ . Minor bands occur around  $200 \text{ cm}^{-1}$ . The  $240 \text{ cm}^{-1}$  band has been interpreted previously as the  $0 \rightarrow 1$  transition of the infrared active torsion. But the  $223 \text{ cm}^{-1}$  band is not the  $1 \rightarrow 2$  transition of the same mode, as one might believe. If this were the case, the overtone band should occur at  $463 \text{ cm}^{-1}$ . The Raman spectrum in Figure 2 clearly shows that there is no band at this frequency. There are bands at  $480$  and  $450 \text{ cm}^{-1}$ , as previously reported, in addition to the bending mode at  $412 \text{ cm}^{-1}$ . Knowing the far infrared spectrum, the  $480 \text{ cm}^{-1}$  band may really be interpreted as the overtone of the infrared active torsion, assuming that the  $240 \text{ cm}^{-1}$  band consists of at least two torsional transitions. The infrared inactive mode was estimated by Fateley and Miller<sup>14</sup> and several normal coordinate calculations to be at about  $200 \text{ cm}^{-1}$ . Therefore, the  $395 \text{ cm}^{-1}$  band could well be the overtone of this torsion. All Raman bands are strongly polarized. In the torsional region, only a very broad, very weak Raman signal with no structure was observed.

In order to check this preliminary assignment and to determine a better potential function, a computer program has been developed. It solves the eigenvalue problem of the internal Hamiltonian of any model with two internal rotors of  $C_{3v}$  symmetry (see Table I). Since the models for dimethylether  $d_0$ ,  $d_3$  and  $d_6$  have higher symmetry than the most general model, some of the terms in this operator vanish or have equal coefficients. The program is set up in the direct product basis of two exponential basis sets and uses a diagonalization subroutine for complex Hermitian matrices. It first solves the eigenvalue

problems of the single top models (first and second line, respectively) and transforms the matrices of the operators  $\hat{P}_i$ ,  $\cos 3\tau_i$  and  $\sin 3\tau_i$  into their eigenbasis. Using direct product calculus, the Hamiltonian matrix is set up for the coupled problem and diagonalized. A standard iteration routine allows the fitting of observed transition frequencies and determination of the potential coefficients. It further calculates the squares of the matrix elements of  $\cos 3\tau_i$  and  $\sin 3\tau_i$  and weights them with the Boltzman population factor. These numbers are used to identify the energy levels with respect to the symmetry of the two top model and to obtain rough estimates of relative intensities of the transitions. With this program, the fit in Table II was obtained. It looks quite reasonable. However, it does not explain the weak Raman band at  $363 \text{ cm}^{-1}$ . Another weak Raman band at  $381 \text{ cm}^{-1}$  may eventually be interpreted as a line belonging to a different torsional substate of the  $385 \text{ cm}^{-1}$  band, although the calculated splitting is not larger than  $2 \text{ cm}^{-1}$ . Another point which is not quite satisfactory is the fact that the intensity ratio of the  $450$  and  $480 \text{ cm}^{-1}$  bands is about 1:2, the calculated ratio, however, is about 1:10. For the present, this fit seems to be the best we can find and the corresponding potential coefficients are given in the lower half of Table II. For comparison, the results of a recent microwave investigation of ground state and two torsionally excited states splittings are given. Instead of using  $V_{30} = V_{03}$ ,  $V_{33}$  and  $V_{33}^i$  to define the potential function, an identical function may be written as

$$2V = V_{30}(1-c3\tau_0)(1-c3\tau_1) + V_{\text{eff}}(1-c3\tau_0c3\tau_1) + V_{33}^i s^3\tau_0 s^3\tau_1.$$

The coefficients in this equation have lower dispersions and are much less correlated to each other. They are also given in Table II.

In Figure 3 is shown the far infrared spectrum of  $\text{CD}_3\text{OCH}_3$ . Due to the higher resolution, it shows a lot more features than the previously published spectrum. Note the band series at  $162.0$ ,  $154.5$ ,  $145.2$  and  $133.2 \text{ cm}^{-1}$ . They

may be assigned to the  $0 \rightarrow 1$ ,  $1 \rightarrow 2$ ,  $2 \rightarrow 3$  and  $3 \rightarrow 4$  transitions of the  $\text{CD}_3$  torsional mode. In the region of the "light" methyl torsion, the interpretation is not straightforward. The main features are at 224.0, 222.0, 218.2, 210.7 and  $204.9 \text{ cm}^{-1}$ , all but the first two are rather broad and probably consisting of more than one torsional transition. The Raman spectrum, shown in Figure 4 is similarly complex in the overtone region. The bending mode at  $373 \text{ cm}^{-1}$  is relatively weak compared with the light isotope. Bands of similar intensities appear at 384 and  $380 \text{ cm}^{-1}$ , the former being assigned to the lowest combination transition of both torsions. The weak band at  $316.5 \text{ cm}^{-1}$  is assigned as the  $0 \rightarrow 2$  transition of the  $\text{CD}_3$  torsion, since its frequency is the sum of 162.0 and  $154.5 \text{ cm}^{-1}$  which corresponds to two infrared transitions. The  $442 \text{ cm}^{-1}$  band is the sum of 224 and  $218 \text{ cm}^{-1}$ ; therefore, this band is assigned to the  $0 \rightarrow 2$  transition of the light methyl torsion. We suspect Fermi resonance type interactions to be responsible for the particular intensity pattern in the  $380 \text{ cm}^{-1}$  region, although a good frequency fit could be obtained without this assumption. The results of the fit are represented in Table III. This fit again is reasonable, although it does not explain everything, for instance, the infrared absorption at  $210 \text{ cm}^{-1}$ . Again, the potential function can be expressed by different coefficients. One choice is to use  $V_s$ ,  $V_d$  and  $V_{\text{seff}}$  in the form

$$2V = V_s(1 - c3\tau_0)(1 - c3\tau_1) + V_d(\cos 3\tau_2 - c3\tau_1) + V_{\text{seff}}(1 - c3\tau_0 c3\tau_1) + V'_{33} s3\tau_0 s3\tau_1.$$

The coefficients are also given in this Table III, the correlations are weaker than in the original set and they are better determined.

The Raman spectrum of  $\text{CD}_3\text{OCD}_3$  in Figure 5 shows, besides the bending mode, at least 15 different polarized lines. At present, no interpretation can be given since the analysis has not been completed yet.

The agreement between the potential coefficients of different isotopes is not very exciting. Similarly, there are certain discrepancies with the results of the microwave splittings analysis. Several reasons may be responsible for

these facts. The present assignment may not be correct since, as already pointed out, several features of the spectra can not be explained, although we tried quite a number of possibilities. The microwave analysis was performed using splittings in the ground state and two excited states. We think that from this information, at most, three and probably only two potential coefficients can be determined. The torsional analysis, however, includes transitions involving energy levels almost to the top of the effective barrier, so that the information about the higher parts of the potential surface should be much better than in the microwave investigation. Despite these doubts, the cos-cos coupling term is definitely a large number and it has definitely no relation at all with the sin-sin coefficient which has been proposed in some earlier work.



Table I

## Hamiltonian and Basis Functions

$$\begin{aligned} \hat{H} = & \frac{1}{2}g^{44}\hat{p}_0^2 + \frac{1}{2}\{V_{30}(1-\cos 3\tau_0) + V_{60}(1-\cos 6\tau_0) + V'_{60}\sin 6\tau_0\} \\ & + \frac{1}{2}g^{55}\hat{p}_1^2 + \frac{1}{2}\{V_{03}(1-\cos 3\tau_1) + V_{06}(1-\cos 6\tau_1) + V'_{06}\sin 6\tau_1\} \\ & + g^{45}\hat{p}_0\hat{p}_1 + \frac{1}{2}\{V_{33}(\cos 3\tau_0\cos 3\tau_1 - 1) + V'_{33}\sin 3\tau_0\sin 3\tau_1 \\ & \quad + V''_{33}\sin 3\tau_0\cos 3\tau_1 + V'''_{33}\cos 3\tau_0\sin 3\tau_1\} \end{aligned}$$

$$\{\phi_{mn}(\tau_0, \tau_1)\} = \{\phi_m(\tau_0)\} \times \{\phi_n(\tau_1)\}$$

$$\phi_\ell(\tau_j) = (2\pi)^{-\frac{1}{2}} \exp(i\ell\tau_j)$$

$$\left. \begin{array}{l} \text{CH}_3\text{OCH}_3 \\ \text{CD}_3\text{OCD}_3 \end{array} \right\} \begin{cases} g^{44} = g^{55}; V_{30} = V_{03}; V_{60} = V_{06} \\ V'_{60} = V'_{06} = V''_{33} = V'''_{33} = 0 \end{cases}$$

$$\text{CD}_3\text{OCH}_3 \quad V'_{60} = V'_{06} = V''_{33} = V'''_{33} = 0$$

PRECEDING PAGE BLANK NOT FILMED

Table II

Fit of Torsional Frequencies for  $\text{CH}_3\text{OCH}_3$ 

$$g^{44} = g^{55} = 13.5644 \text{ cm}^{-1} \quad g^{45} = -2.7381 \text{ cm}^{-1}$$

Transition	Obs( $\text{cm}^{-1}$ )	Calc( $\text{cm}^{-1}$ )	Obs-Calc	Transition	Obs( $\text{cm}^{-1}$ )	Calc( $\text{cm}^{-1}$ )	Obs-Calc
1 → 3	241.0	243.8	-2.8	1 → 6	481.3	484.2	-2.9
3 → 6	239.9	240.4	-0.5	2 → 9	450.2	446.9	3.3
2 → 5	223.2	222.7	0.5	1 → 4	395.5	397.3	-1.8
5 → 9	226.0	224.2	1.8	2 → 7	385.2	383.8	1.4
4 → 8	200.7	199.7	1.0				
5 → 9	224.8	223.4	1.4	} These transitions were not fitted. They are assigned to different torsional substates.			
4 → 8	199.0	198.1	0.9				

Standard deviation of frequencies  $2.51 \text{ cm}^{-1}$

Potential Coefficients ( $\text{cm}^{-1}$ ) <sup>a</sup>	Microwave	Derived Potential Coefficients	Microwave <sup>b</sup>
$V_{30}=V_{03}$	$1132.91 \pm 37.52$	$V_{\text{eff}} = V_{30} - V_{33}$	$909.05 \pm 0.25$
$V_{33}$	$214.25 \pm 47.09$		$918.66 \pm 10.95$
$V'_{33}$	$20.92 \pm 8.05$		$909.05 \pm 0.25$
$V_{60}=V_{06}$	0(not varied)		

<sup>a</sup> Error limit = dispersion of parameter

<sup>b</sup> J. R. Durig, Y. S. Li and P. Groner, accepted for publication (J. Mol. Spectry).

Table III

Fit of Torsional Frequencies for  $\text{CD}_3\text{OCH}_3$ 

$$g^{44} = 13.0973 \text{ cm}^{-1} \quad g^{55} = 8.1872 \text{ cm}^{-1} \quad g^{45} = -2.4764 \text{ cm}^{-1}$$


Transition	Obs( $\text{cm}^{-1}$ )	Calc( $\text{cm}^{-1}$ )	Obs-Calc	Transition	Obs( $\text{cm}^{-1}$ )	Calc( $\text{cm}^{-1}$ )	Obs-Calc
1 $\rightarrow$ 2	162.0	161.8	0.2	1 $\rightarrow$ 3	224.0	225.9	-1.9
2 $\rightarrow$ 4	154.5	155.3	-0.8	3 $\rightarrow$ 6	218.2	218.3	-0.1
4 $\rightarrow$ 7	145.2	145.6	-0.4	1 $\rightarrow$ 6	442.0	444.2	-2.2
7 $\rightarrow$ 10	133.0	131.8	1.2	3 $\rightarrow$ 11	435.0	433.6	1.4
1 $\rightarrow$ 4	316.5	317.2	-0.7	2 $\rightarrow$ 5	222.0	221.9	0.1
1 $\rightarrow$ 5	384.0	383.7	0.3	5 $\rightarrow$ 9	204.9	202.2	2.7
2 $\rightarrow$ 8	390.0	381.1	-1.1	6 $\rightarrow$ 11*	215.8 sh	215.3	0.5
3 $\rightarrow$ 9	361.5	360.0	1.5				


\* not fitted


Standard deviation of frequencies  $1.55 \text{ cm}^{-1}$ 

Potential Coefficients ( $\text{cm}^{-1}$ ) <sup>a</sup>	Microwave <sup>b</sup>	Derived Potential Coefficients	Microwave <sup>b</sup>
$V_{30}$	$1313.52 \pm 30.31$	$V_s = \frac{1}{2}(V_{30} + V_{03})$	$1285.97 \pm 30.08$
$V_{03}$	$1258.41 \pm 30.45$	$V_d = \frac{1}{2}(V_{30} - V_{03})$	$27.55 \pm 4.24$
$V_{33}$	$417.11 \pm 35.54$	$V_{\text{seff}} = V_s - V_{33}$	$868.85 \pm 6.14$
$V_{33}^{\dagger}$	$34.22 \pm 7.99$	$V_{30\text{eff}} = V_{30} - V_{33}$	$896.41 \pm 6.99$
$V_{06} = V_{60}$	0(not varied)	$V_{03\text{eff}} = V_{03} - V_{33}$	$841.30 \pm 7.92$
			$903.81 \pm 0.65$
			$6.64 \pm 0.56$
			$903.81 \pm 0.65$
			$910.45 \pm 0.17$
			$897.18 \pm 1.21$

<sup>a</sup> Error limit = dispersion of parameter<sup>b</sup> J. R. Durig, Y. S. Li and P. Groner, accepted for publication (J. Mol. Spectry).











Fig. 4





Although this study is not yet complete, we should be able to provide a complete manuscript by the next progress report time.

For several progress reports we have mentioned our work on methylchloroformate and anticipated that this work would soon be finished. We are pleased to report that a manuscript has been submitted to the Journal of Molecular Spectroscopy on this molecule.

The structure of methylchloroformate has been the subject of considerable interest. Four dipole moment studies have been reported.<sup>15-18</sup> From two of these<sup>15,17</sup> the results were interpreted in terms of a planar s-cis structure. The data from the other dipole moment studies<sup>16,18</sup> were interpreted in terms of a planar s-trans structure.

Three analyses of the vibrational spectra have been reported.<sup>19-21</sup> These analyses were all in terms of a molecular model of  $C_s$  symmetry and were consistent with an s-trans structure. An electron diffraction study of the vapor of methylchloroformate was performed.<sup>22</sup> The data yielded only an s-trans structure that was slightly non-planar with the methyl group approximately  $20^\circ$  out of the plane of the other heavy atoms.

An investigation of the infrared spectrum and of the proton NMR spectrum of methylchloroformate in solution was reported.<sup>23</sup> The authors found evidence for the coexistence of both the s-cis and the s-trans conformations and indicated that the s-cis was more stable. From a study comparing observed and calculated  $^{35}\text{Cl}$  nuclear quadrupole resonance frequencies the authors<sup>24</sup> concluded that the s-trans was the most likely stable structure in the solid.

A microwave spectroscopic investigation<sup>25</sup> of natural abundance  $^{35}\text{Cl}$  and  $^{37}\text{Cl}$  isotopic species of methylchloroformate showed that the planar s-trans molecular structure was most consistent with the data. Small amounts of a second rotamer, however, could not be precluded.

Therefore, in an effort to clear up the ambiguities in the conformational preferences of methylchloroformate and to provide more complete structural data, the investigation of the microwave, Raman and far infrared spectra of the vapor were undertaken. The results of this study can be summarized by the abstract which follows of the paper that has been submitted for publication.

Abstract: The microwave spectra of six isotopic species of methylchloroformate,  $\text{ClCO}_2\text{CH}_3$ , have been recorded from 18.0 to 40.0 GHz. Structural parameters have been determined and it is shown that the only stable conformer at ambient temperature is the s-trans. From the Stark effect, the dipole moment components were determined to be  $|\mu_a| = 1.7 \pm 0.2$ ,  $|\mu_b| = 1.0 \pm 0.2$ , and  $|\mu_c| = 2.0 \pm 0.2\text{D}$ . The Raman and far infrared spectra of the vapor are reported. Four cases of Fermi resonance have been observed in the Raman effect. Both the methyl and methoxy torsions have been observed in the far infrared and the methyl barrier to internal rotation has been determined to be 1.15 kcal/mole (1.19 kcal/mole for the  $\text{CD}_3$  rotor) which is lower than the 1.23 kcal/mole obtained from the microwave splitting method. It is shown from both the  $^{13}\text{C}$  and  $^1\text{H}$  NMR spectra that only one conformer exists which is contrary to what was previously reported. The vibrational spectrum of the solid is also reported and discussed.

For the complete paper, see appendix I of this report.

In addition to our studies of carbonyl containing compounds, we have been pursuing investigations on the vibrational and rotational spectra of a number of nitrogen and phosphorus containing compounds which are likely to be found in the atmospheres of the Jovian planets. In earlier progress reports we reported the vibrational spectra and structure of dimethylphosphine, ethylphosphine and isopropylphosphine. As a continuation of these studies, we have investigated the vibrational spectrum of tertiary-butylphosphine.

Vibrational spectroscopy has been shown to be a valuable tool in the determination of intramolecular potential functions governing internal rotation about single bonds.<sup>26,27</sup> This method has recently been extended to the study of asymmetric tops rotating against a molecular frame, as opposed to the more familiar symmetric tops such as  $-\text{CH}_3$  and  $-\text{SiH}_3$ .<sup>28</sup> If the molecular frame is asymmetric as well, it is possible for there to be spectroscopically distinct rotational isomers or conformers. Both ethylphosphine ( $\text{CH}_3\text{CH}_2\text{PH}_2$ ) and isopropylphosphine [ $(\text{CH}_3)_2\text{CHPH}_2$ ] are examples of this type of system since the  $\text{PH}_2$  group constitutes an asymmetric internal rotor. The internal rotation barrier for the  $\text{PH}_2$  top has been determined for both of these molecules.<sup>29,30</sup> If, however, the molecular frame is symmetric, as is the case in tertiary-butylphosphine, [ $(\text{CH}_3)_3\text{CPH}_2$ ], only one conformation is possible and the potential function governing internal rotation is considerably simplified, becoming three-fold symmetric as in the case of methylphosphine ( $\text{CH}_3\text{PH}_2$ ). Thus, the determination of this potential function for t-butylphosphine is a natural extension of the prior studies of ethyl-<sup>29</sup> and isopropylphosphine.<sup>30</sup> Also, it would be of interest to determine the barriers to internal rotation of the methyl groups in this molecule, as part of continuing studies<sup>31,32</sup> of the top-top interactions in such "three-top" molecules.

Finally, the assignment of the vibrational spectrum of t-butylphosphine would finish the series of singly alkyl-substituted phosphines: methyl, ethyl

and isopropyl. Group frequencies determined for the  $\text{PH}_2$  moiety from these studies could be expected to apply to almost any hydrocarbon derivative. The results of this study have been submitted for publication in the Journal of Molecular Structure and can be summarized by the abstract of the paper (see appendix II for the complete paper).

ABSTRACT: The infrared spectra of gaseous and solid tertiary-butylphosphine,  $[(\text{CH}_3)_3\text{CPH}_2]$ , have been recorded from  $50 \text{ cm}^{-1}$  to  $3500 \text{ cm}^{-1}$ . The Raman spectra of gaseous, liquid and solid  $(\text{CH}_3)_3\text{CPH}_2$  have been recorded from 10 to  $3500 \text{ cm}^{-1}$ . A vibrational assignment of the 42 normal modes has been made. A harmonic approximation of the methyl torsional barrier from observed transitions in the solid state gave a result of 4.22 kcal/mole and 3.91 kcal/mole in the gaseous state. Hot band transitions for the phosphino torsional mode have been observed. The potential function for internal rotation about the C-P bond has been calculated. The two potential constants were determined to be:

$$V_3 = 2.79 \pm 0.01 \text{ kcal/mole and } V_6 = 0.07 \pm 0.01 \text{ kcal/mole.}$$

## FUTURE WORK

Studies are nearly complete on  $(\text{CH}_3)_2\text{NH}$  and the closely related silane,  $(\text{CH}_3)_2\text{NSiH}_3$ . The papers are currently being written on both of these compounds. For the dimethylamine work we have written a new computer program for the interpretation of the torsional data for the two-top case. We have also undertaken a study of the vibrational spectrum of isopropyl amine. Studies have also been initiated on the low frequency spectra of dimethylphosphine. Vibrational work is also being done on propylaldehyde. We shall also continue our current work on several compounds which have potential interest as planetary atmosphere materials.

## References

1. G. P. Kuiper, *Atmospheres of the Earth and Planets*, University of Chicago Press, Chicago, 1952.
2. J. R. Durig, S. F. Bush, and E. E. Mercer, *J. Chem. Phys.*, 44, 4238 (1966).
3. J. R. Durig, S. F. Bush, and F. G. Baglin, *J. Chem. Phys.*, 49, 2106 (1968).
4. J. R. Durig, W. C. Harris, and D. W. Wertz, *J. Chem. Phys.*, 50, 1449 (1969).
5. J. R. Durig and W. C. Harris, *J. Chem. Phys.*, 51, 4457 (1969).
6. J. R. Durig and W. C. Harris, *J. Chem. Phys.*, 55, 1735 (1971).
7. J. R. Durig, R. W. MacNamee, L. B. Knight, and W. C. Harris, *Inorg. Chem.*, 12, 804 (1973).
8. J. R. Durig, J. W. Thompson, and J. D. Witt, *Inorg. Chem.*, 11, 2477 (1972).
9. *Chem. Eng. News*, 4, 4 (1974).
10. *Ind. Res.*, May 1974, page 29, by Nat. Radio Astronomy Observatory at Kitt Peak.
11. J. R. Durig, C. M. Player, Jr., J. Bragin and Y. S. Li, *J. Chem. Phys.*, 55, 2895 (1971).
12. E. C. Tuazon and W. G. Fateley, *J. Chem. Phys.*, 54, 4450 (1971).
13. R. C. Taylor and G. L. Vidale, *J. Chem. Phys.*, 26, 122 (1957).
14. W. G. Fateley and F. A. Miller, *Spectrochim. Acta*, 18, 977 (1962).
15. S. Mizushima and M. Kubo, *Bull. Chem. Soc. Jap.*, 13, 174 (1938).
16. R. J. W. LeFevre and A. Sundaram, *J. Chem. Soc. (Lond.)* 1962, 3904.
17. E. Bock and D. Iwacha, *Can. J. Chem.*, 45, 3177 (1967).
18. M. J. Aroney, R. J. W. LeFevre, R. K. Pierens, and H. L. K. The, *Aust. J. Chem.*, 22, 1599 (1969).
19. B. Collingwood, H. Lee, and J. K. Wilmshurst, *Aust. J. Chem.*, 19, 1637 (1966).
20. R. A. Nyquist, *Spectrochim. Acta*, 28A, 285 (1972).
21. J. E. Katon and M. G. Griffin, *J. Chem. Phys.*, 59, 5868 (1973).
22. J. M. O'Gormm, W. Shand, Jr., and V. Schomaker, *J. Amer. Chem. Soc.*, 72, 4222 (1950).
23. M. Oki and H. Nakanishi, *Bull. Chem. Soc. Jap.*, 45, 1552 (1972).
24. A. J. Cohen and M. A. Whitehead, *J. C. S. Faraday II*, 68, 649 (1972).

25. D. G. Lister and N. L. Owen, *J. C. S. Faraday II*, 69, 1036 (1973).
26. J. R. Durig, S. M. Craven, and W. C. Harris, *Vibrational Spectra and Structure*, Vol. 1, J. R. Durig, ed., Marcel Dekker, New York, N.Y., 73-179 (1972).
27. A. V. Cunliffe, *Internal Rotation in Molecules*, W. J. Orville-Thomas, ed., John Wiley and Sons, New York, N.Y., 217-252 (1974).
28. W. G. Fateley, *Pure App. Chem.*, 36, 119 (1973).
29. J. R. Durig and A. W. Cox, Jr., *J. Chem. Phys.*, 63, 2303 (1975)
30. J. R. Durig and A. W. Cox, Jr., *J. Phys. Chem.*, 80, 000 (1976).
31. J. R. Durig, S. M. Craven and J. Bragin, *J. Chem. Phys.*, 51, 5663 (1969).
32. J. R. Durig, S. M. Craven, J. H. Mulligan and C. W. Hawley, *J. Chem. Phys.*, 58, 1281 (1973).

## APPENDIX I

### THE MICROWAVE, RAMAN, FAR INFRARED AND NMR SPECTRA, STRUCTURE AND DIPOLE MOMENT OF METHYLCHLOROFORMATE

#### Abstract

The microwave spectra of six isotopic species of methylchloroformate,  $\text{ClCO}_2\text{CH}_3$ , have been recorded from 18.0 to 40.0 GHz. Structural parameters have been determined and it is shown that the only stable conformer at ambient temperature is the s-trans. From the Stark effect the dipole moment components were determined to be  $|\mu_a| = 1.7 \pm 0.2$ ,  $|\mu_b| = 1.0 \pm 0.2$ , and  $|\mu_t| = 2.0 \pm 0.2\text{D}$ . The Raman and far infrared spectra of the vapor are reported. Four cases of Fermi resonance have been observed in the Raman effect. Both the methyl and methoxy torsions have been observed in the far infrared and the methyl barrier to internal rotation has been determined to be 1.15 kcal/mole (1.19 kcal/mole for the  $\text{CD}_3$  rotor) which is lower than the 1.23 kcal/mole obtained from the microwave splitting method. It is shown from both the  $^{13}\text{C}$  and  $^1\text{H}$  NMR spectra that only one conformer exists which is contrary to what was previously reported. The vibrational spectrum of the solid is also reported and discussed.



## INTRODUCTION

The structure of methylchloroformate has been the subject of considerable interest. Four dipole moment studies have been reported.<sup>1-4</sup> From two of these<sup>1,3</sup> the results were interpreted in terms of a planar s-cis structure. The data from the other dipole moment studies<sup>2,4</sup> were interpreted in terms of a planar s-trans structure (Fig. 1).

Three analyses of the vibrational spectra have been reported.<sup>5-7</sup> These analyses were all in terms of a molecular model of  $C_s$  symmetry and were consistent with an s-trans structure. An electron diffraction study of the vapor of methylchloroformate was performed.<sup>8</sup> The data yielded only an s-trans structure that was slightly non-planar with the methyl group approximately 20° out of the plane of the other heavy atoms.

An investigation of the infrared spectrum and of the proton NMR spectrum of methylchloroformate in solution was reported.<sup>9</sup> The authors found evidence for the coexistence of both the s-cis and the s-trans conformations and indicated that the s-cis was more stable. From a study comparing observed and calculated <sup>35</sup>Cl nuclear quadrupole resonance frequencies the authors<sup>10</sup> concluded that the s-trans was the most likely stable structure in the solid.

A microwave spectroscopic investigation<sup>11</sup> of natural abundance <sup>35</sup>Cl and <sup>37</sup>Cl isotopic species of methylchloroformate showed that the planar s-trans molecular structure was most consistent with the data. Small amounts of a second rotamer, however, could not be precluded.

Therefore, in an effort to clear up the ambiguities in the conformational preferences of methylchloroformate and to provide more complete structural data, the investigation of the microwave, Raman and far infrared spectra of the vapor were undertaken.

PRECEDING PAGE BLANK NOT FILMED

## EXPERIMENTAL

Practical grade methylchloroformate was purchased from Aldrich Chemical Company, Incorporated and was fractionally distilled before use. The isotopic derivatives were all prepared by the reaction of the appropriate isotopic form of methanol with phosgene.<sup>12</sup> The  $\text{CD}_3\text{OD}$  and  $\text{CH}_2\text{DOH}$  were obtained from Merck, Sharpe and Dohme, Canada Limited. The  $^{13}\text{CH}_3\text{OH}$  was purchased from Stohler Isotope Chemicals, Inc. The methanols were all used without further purification (Stated purities:  $\text{CD}_3\text{OD}$  and  $\text{CH}_2\text{DOH}$  > 95%;  $^{13}\text{CH}_3\text{OH}$  60-70%). The phosgene was obtained from Matheson and was used without further purification. The 1,2-dibromo-1,1,2,2-tetrafluoroethane was purchased from Columbia Organic Chemical Co. and was fractionally distilled before use.

The microwave spectra were recorded with a Hewlett-Packard 8460A MRR spectrometer in the 12.4-40.0  $\text{GHz}$  region. Samples were run both with the waveguide at room temperature and with the waveguide packed in dry ice. Sample pressures were 100 microns for survey spectra and typically 4-10 microns for measurement of the quadrupole hyperfine components. In general, the frequency accuracy should be no worse than  $\pm 0.05 \text{ MHz}$ .

The Raman spectra were recorded on a Cary Model 82 spectrophotometer with a Spectra Physics Model 171 argon ion laser. The  $5145\overset{\circ}{\text{A}}$  line was used for excitation of the samples. The gas was contained at its room temperature vapor pressure in a quartz cell and the laser beam was multipassed through the cell with the standard Cary accessory. Laser power for the gas sample was 5.5 W measured at the laser head. Laser power for the solid sample was 2.0 W measured at the laser head.

The cryostat used was a Spectrim Spectrometric Sample Conditioner controlled by a Lake Shore Electronics, Inc. Cryogenic Temperature Indicator/Controller. The temperature of the sample was maintained at  $18.7 \pm 0.1^\circ \text{K}$

and was measured with a calibrated Silicon Diode Sensor obtained from Cryogenic Technology, Inc.

The far infrared spectra of the vapor were recorded on a Digilab Model 15B Fourier Transform Interferometer with a 12.5 micron Mylar beamsplitter. The vapor was contained in a 10 cm. glass cell with polyethylene windows at its room temperature vapor pressure. Resolution was equal to  $0.5 \text{ cm}^{-1}$ .

All NMR measurements were made on a Varian XL-100-15 NMR spectrometer equipped with a 16K (1K = 1024 words) 620I computer and a 2 million word disc (VDM-36). Temperatures were measured with a thermocouple. All spectra were obtained in the pulse Fourier Transform mode.

The  $^{13}\text{C}$  spectra were obtained using broadband noise-modulated proton decoupling with the spectrometer locked to the  $^{19}\text{F}$  resonance of an external sample of  $\text{C}_6\text{F}_6$ . Samples for  $^{13}\text{C}$  measurements were 50%  $\text{CH}_3\text{-O-C}^{\text{O}}\text{-Cl}$ , 50%  $\text{CS}_2$ . Proton spectra were obtained with the spectrometer locked to the  $^2\text{H}$  resonance of internal  $\text{CD}_2\text{Cl}_2$ . The sample used for proton measurements was approximately 5 mM in  $\text{CH}_3\text{-O-C-Cl}$  in a 50/50 mixture of  $\text{CS}_2$  and  $\text{CD}_2\text{Cl}_2$ . Proton  $T_1$  measurements were obtained using the Freeman-Hill $^{13}$  inversion recovery pulse sequence. Peak heights were employed to determine the  $T_1$  values.

#### MICROWAVE RESULTS

##### Rotational Spectra

The rotational spectra of all the isotopic species of methylchloroformate were first assigned on the basis of the known spectral pattern of a prolate asymmetric ( $\kappa \approx -0.86$ ) rotor compared to high pressure ( $\sim 100$  Torr) survey spectra. Individual lines were then remeasured at a pressure (4-10 Torr) suitable for resolution of the hyperfine splittings that result from the nuclear quadrupole coupling. The quadrupole coupling constants were then fit to the observed splitting. The positions of all observed low pressure

transitions were then corrected for quadrupole shift and used to fit the rotational constants of a rigid rotor. Initial fits were made with a-type R branch transitions and were used to approximately locate the b-type R branches. The b-type R branches were then identified by their predicted quadrupole hyperfine splittings and were used to refine the final fit of the rotational constants. The frequencies from the low pressure measurements for all the isotopic species of methylchloroformate studied here are listed in Table I. The rotational constants, moments of inertia, inertial defect, and nuclear quadrupole coupling constants for these isotopic species are listed in Table II.

### STRUCTURE

The coordinates of the atoms for which isotopic substitutions were made have been calculated using the Kraitchman equations<sup>14</sup> and are listed in Table III. For a non-planar asymmetric top only N-3 substitutions are needed to determine the structure. This is true because there are nine moment relations:

$$I_{xx} = \sum_i m_i (y_i^2 + z_i^2) \quad (1)$$

$$I_{xy} = -I_{yx} = -\sum_i m_i x_i y_i \quad (2)$$

$$\sum_i m_i x_i = 0 \quad (3)$$

that may be solved simultaneously. The inertial defect (Table II) for all isotopic species is indicative of an essentially planar molecule with only two hydrogens symmetrically out of the plane. For that reason the z coordinate of the atoms in the plane has been taken as zero. The nine moment relations then yield six non-trivial equations of which only five are independent. To completely solve the structure we need to calculate the six in-plane coordinates of the three unsubstituted atoms. It is, therefore, possible to calculate

the complete structure assuming only one in-plane coordinate of one of the three unsubstituted atoms.

The electron diffraction study of methylchloroformate<sup>8</sup> placed the carbonyl bond length at 1.19Å. From microwave studies of some related structures  $r_{C=O}$  values have been calculated for COH<sub>2</sub> as 1.20<sup>15</sup>, for COCl<sub>2</sub> as 1.17<sup>16</sup>, for HCO<sub>2</sub>CH<sub>3</sub> as 1.20<sup>17</sup>, for HCO<sub>2</sub>H as 1.20<sup>18</sup>, for NH<sub>2</sub>COH as 1.19<sup>19</sup>, and for HFCO as 1.18<sup>20</sup>. Consideration of the carbonyl bond length listed strongly suggests that  $r_{C=O}$  for methylchloroformate should be between 1.18Å and 1.20Å.

In Table IV are listed the coordinates of the three unsubstituted atoms calculated from the moment relations with the x-coordinate of the carbonyl carbon chosen so that the carbonyl bond length equals 1.19Å. The coordinates listed in Table III and Table IV reproduce the observed rotational constants in Table II to within ± 0.02%. The structural parameters calculated from these coordinates are listed in Table V. The structure of methylchloroformate is depicted in Fig. 2.

#### NUCLEAR QUADRUPOLE COUPLING CONSTANT

The first order chlorine nuclear quadrupole coupling constants were evaluated by a least squares fit of the observed hyperfine splittings and are listed in Table II. The value of the coupling constant along the bond axis may be calculated if the angle,  $\theta$ , between the carbon-chlorine bond and the A principal axis is known using the equations:

$$x_{xx} \cos^2 \theta + x_{yy} \sin^2 \theta = x_{aa} \quad (4)$$

$$x_{xx} \sin^2 \theta + x_{yy} \cos^2 \theta = x_{bb} \quad (5)$$

where the x axis lies along the carbon chlorine bond and y is perpendicular to it in the plane of the molecule and  $x_{cc} = x_{zz}$ . This transformation yields

$\chi_{xx} = -73.7$  MHz,  $\chi_{yy} = 45.1$  MHz, and  $\chi_{zz} = 28.6$  for the parent molecule with  $\theta = 20^\circ$ .

These coupling constants along the bond may then be interpreted to give some information about the nature of the hybridization of the carbon-chlorine bond.<sup>21</sup> In the earlier microwave study of methylchloroformate the authors<sup>11</sup> presented such a discussion and that discussion is not significantly altered by the data presented here.

#### BARRIER TO INTERNAL ROTATION

The barrier to internal rotation of the methyl group was computed from the observed internal rotation splittings of the a-type and b-type R branches. The same method as adopted previously<sup>22</sup> was used to determine the barrier. The observed internal rotation splittings are listed in Table VI with the parameters calculated from the structure listed in Table III and IV. The value for  $V_3$  calculated here is within the error limits of the previously determined  $V_3$ <sup>11</sup> value and is consistent with that found for other substituted methylformates.<sup>23</sup>

#### DIPOLE MOMENT

The second order Stark effects were observed and measured for the following components:  $|M| = 0, 3, 4$  of the  $5_{05} + 4_{04}$  transition. Field calibration was accomplished by measuring the  $M=0$  component of OCS for the  $J = 2 \leftarrow 1$  transition.<sup>24</sup> The observed and calculated Stark coefficients for  $^{35}\text{Cl} \ ^{12}\text{C} \ ^{16}\text{O}_2 \ ^{12}\text{CH}_3$  are listed in Table VII. The microwave spectrum associated with this molecule is quite rich. The Stark components that were tracked could be followed as a function of voltage for only 2-3 MHz before being lost in other unidentified Stark component. Considerable effort was expended in an attempt to find other suitable  $|M|$  components

to track; none were found. The error limits on the dipole components are, therefore, quite high.

### VIBRATIONAL RESULTS

The Raman spectra of methylchloroformate as a vapor at room temperature and as a solid at 18.7°K are shown in Fig. 3. The far infrared spectra of gaseous methylchloroformate and methylchloroformate-d<sub>3</sub> are presented in Fig. 4. Transitions marked with an arrow correspond to a small amount of an impurity (HCl). The vibrational data are listed in Table VIII with their assignments.

The Raman spectrum of gaseous methylchloroformate (Fig. 4A) is largely interpretable in terms of a previously reported assignment.<sup>7</sup> The features at 2905 and 2845 cm<sup>-1</sup> correspond to transitions observed in the Raman spectrum of the liquid<sup>7</sup> at 2897 and 2844 cm<sup>-1</sup>. They are assigned as the overtones of the antisymmetric,  $\nu_4/\nu_{14}$ , and the symmetric,  $\nu_5$ , methyl deformations in Fermi resonance with the symmetric CH stretch,  $\nu_2$ .

The carbonyl stretch,  $\nu_2$ , is assigned at 1801 cm<sup>-1</sup> because it is the stronger of the two Q-branches in the region; the other Q branch (1792 cm<sup>-1</sup>) is ascribed to Fermi resonance of the combination of  $\nu_8$ , the COC symmetric stretch, and  $\nu_9$ , the carbon chlorine stretch, (970 + 823 = 1793) with  $\nu_2$  the carbonyl stretch.

The result of another Fermi resonance interaction is apparent in the pair of Q branches at 977/962 cm<sup>-1</sup>. One of the Q branches may be assigned to  $\nu_8$ , the COC symmetric stretch and the other Q branch is the overtone of  $\nu_{10}$ , an in-plane skeletal bend (2 X 487 = 974). The last feature of the Raman spectrum of the vapor of methylchloroformate is the Q branch series starting at 487 cm<sup>-1</sup>. The relative intensity of the Q branches at 487, 480, and 472 cm<sup>-1</sup> indicates they are hot bands of  $\nu_{12}$ , the COC in-plane bend.

The far infrared spectra of methylchloroformate- $d_0$  and  $d_3$  (Fig. 4) cover the region expected for  $\nu_{17}$  and  $\nu_{18}$ , the methoxy and the methyl torsions, respectively. The methoxy torsion is assigned to the Q branches at  $163.2 \text{ cm}^{-1}$  for the light compound and to the doublet at  $148.8/148.2 \text{ cm}^{-1}$  for the perdeutero compound. The apparent splitting of the Q branch for the perdeutero compound remains unexplained.

The methyl torsion,  $\nu_{18}$ , had been reported<sup>7</sup> as occurring at  $62 \text{ cm}^{-1}$  in the liquid phase. The three-fold barrier and the structure obtained from the microwave data indicate that it should be observed near  $134 \text{ cm}^{-1}$  in the spectra of the vapor. It is observed at  $129.2 \text{ cm}^{-1}$  in the spectrum of the "light" vapor (Fig. 4A) and at  $99.0 \text{ cm}^{-1}$  in the spectra of the perdeutero compound (Fig. 4B). The shift factor,  $\nu_H/\nu_D = 1.305$  is lower than should be expected if the torsion was not mixed with other normal modes.

The barrier to internal rotation of the methyl group,  $V_3$ , was calculated using a computer program similar to one described by Lewis *et al.*<sup>26</sup> The barrier for the  $\text{CH}_3$  rotor is  $1.15 \text{ kcal/mole}$  and for the  $\text{CD}_2$  rotor it is  $1.19 \text{ kcal/mole}$ . Both barriers are lower than that calculated by the microwave splitting method.

The Raman spectrum of methylchloroformate solid at  $18.7^\circ\text{K}$  is quite rich (Fig. 3B). It had been suggested previously<sup>7</sup> that the unit cell of methylchloroformate contained one molecule or that if it contained two molecules it should be considered centrosymmetric. Eight sharp lattice modes are apparent in the Raman spectrum below the lowest fundamental ( $\nu_{18}$ , methyl torsion,  $135 \text{ cm}^{-1}$ ). This number of lattice modes rules out the possibility of there being only one molecule per unit cell and also rules out the possibility of two molecules in a centrosymmetric unit cell. Crystal structure data remain unavailable for methylchloroformate but such



data would be of interest.

There are several apparent examples of crystal splitting evident in the spectrum, some of them quite striking. The pair of bands at  $3028/3016\text{ cm}^{-1}$  had also been reported in the infrared study of the solid.<sup>7</sup> The authors reported an isotopic dilution study of this band pair from which they concluded the pair did not result from crystal splitting. Further examples in the Raman spectrum that apparently result from crystal splitting are  $\nu_3$ , the carbonyl stretch ( $1774/1757/1752\text{ cm}^{-1}$ ),  $\nu_{12}$ , the COC in-plane bend ( $283/277\text{ cm}^{-1}$ ),  $\nu_{15}$ , the  $\text{CH}_3$  rock ( $1152/1147\text{ cm}^{-1}$ ) and  $\nu_{17}$ , the methoxy torsion ( $194/182\text{ cm}^{-1}$ ). These apparent examples of crystal splitting are quite reasonable in view of the crystal structure required by the large number of lattice modes evident in the Raman spectrum.

#### NUCLEAR MAGNETIC RESONANCE RESULTS

The proton decoupled  $^{13}\text{C}$  NMR spectrum of  $\text{ClCO}_2^{13}\text{CH}_3$  in 1,2-dibromo,tetra-fluoroethane at  $29^\circ\text{C}$  consists of a single resonance with a half height linewidth ( $\nu_{1/2}$ ) of 0.8 Hz. The spectrum was studied as a function of temperature to  $-105^\circ\text{C}$ . Throughout the temperature range covered, there was no significant line-broadening observed.

The  $^1\text{H}$  NMR spectrum of methylchloroformate in  $\text{CS}_2/\text{CD}_2\text{Cl}_2$  (50/50) at  $26^\circ\text{C}$  consists of a single resonance with  $\nu_{1/2} = 0.25\text{ Hz}$ .  $T_1$  was measured to be 15 seconds. The spectrum was again studied as a function of temperature to  $-100^\circ\text{C}$ . At  $-100^\circ\text{C}$ ,  $\nu_{1/2} = 0.55\text{ Hz}$  and  $T_1 = 2.36\text{ seconds}$ . The small amount of apparent line-broadening observed at the lower temperature can be completely attributed to the change in  $T_1$  and the fact that the magnetic field was more difficult to shim at  $-100^\circ\text{C}$  because of the  $^2\text{H}$  lock utilized on the Varian model XL-100. For these reasons the small apparent line-broadening in the  $^1\text{H}$  NMR spectrum at  $-100^\circ\text{C}$  should not be ascribed to any chemical exchange processes.<sup>9</sup>

## CONCLUSIONS

The microwave spectrum of methylchloroformate has been thoroughly reinvestigated. The data are interpreted to indicate that only one form of this molecule exists through, given the richness of the spectrum, small amounts of a second rotamer cannot positively be ruled out with these data alone. Further the microwave data are conclusively interpreted to demonstrate that the stable form of methylchloroformate is the s-trans form not the s-cis form as had been previously reported.<sup>9</sup> From this microwave study it has been possible to obtain the structural parameters of this stable form along with the dipole moment components.

The vibrational data include the first direct observation of the methyl torsion of gaseous methylchloroformate and that observation is consistent with the structure and barrier calculated from the rotational data. The Raman spectrum of the solid provides more specific data on which to base generalizations as to the crystal structure of methylchloroformate than the infrared spectrum of the solid alone<sup>7</sup> and again confirms the complimentary nature of the two sources of vibrational data.

The <sup>13</sup>C NMR temperature study is interpreted to be much more strongly suggestive of the likelihood that methylchloroformate has only one stable molecular conformation (s-trans) at room temperature and below. The <sup>1</sup>H NMR temperature study provided data that is felt to demonstrate the need for care in the performance and interpretation of such experiments.<sup>9</sup>

## ACKNOWLEDGEMENT

The authors gratefully acknowledge the financial support given this study by the National Aeronautics and Space Administration through grant NGL-41-002-003. The authors also gratefully acknowledge the services of Dr. Paul Ellis and Mr. David Bailly for providing the NMR data and the expertise necessary for their evaluation.

## REFERENCES

<sup>†</sup>Taken from the thesis of M. G. Griffin which will be submitted to the Department of Chemistry in partial fulfillment of the Ph.D. degree.

1. S. Mizushima and M. Kubo, *Bull. Chem. Soc. Jap.* 13, 174 (1938).
2. R. J. W. LeFevre and A. Sundaram, *J. Chem. Soc. (Lond.)* 1962, 3904.
3. E. Bock and D. Iwacha, *Can. J. Chem.* 45, 3177 (1967).
4. M. J. Aroney, R. J. W. LeFevre, R. K. Pierens, and H. L. K. The, *Aust. J. Chem.* 22, 1599 (1969).
5. B. Collingwood, H. Lee, and J. K. Wilmshurst, *Aust. J. Chem.* 19, 1637 (1966).
6. R. A. Nyquist, *Spectrochim. Acta* 28A, 285 (1972).
7. J. E. Katon and M. G. Griffin, *J. Chem. Phys.* 59, 5868 (1973).
8. J. M. O'Gormm, W. Shand, Jr., and V. Schomaker, *J. Amer. Chem. Soc.* 72, 4222 (1950).
9. M. Oki and H. Nakanishi, *Bull. Chem. Soc. Jap.* 45, 1552 (1972).
10. A. J. Cohen and M. A. Whitehead, *J. C. S. Faraday II*, 68, 649 (1972).
11. D. G. Lister and N. L. Owen, *J. C. S. Faraday II*, 69, 1036 (1973).
12. W. Hentschel, *Chem. Ber.* 18, 1177 (1885).
13. R. Freeman and H. D. W. Hill, Jr., *J. Chem. Phys.*, 53, 5103 (1970).
14. J. Kraitchman, *Am. J. Phys.* 21, 17 (1953).
15. T. Oka, *J. Phys. Soc. Jap.* 15, 2274 (1960).
16. G. W. Robinson, *J. Chem. Phys.* 21, 1741 (1953).
17. R. F. Curl, Jr., *J. Chem. Phys.* 30, 1529 (1959).
18. G. H. Kwei, and R. F. Curl, Jr., *J. Chem. Phys.* 32, 1592 (1960).
19. C. C. Costain, and J. M. Jowling, *J. Chem. Phys.* 32, 158 (1960).
20. O. H. LeBlanc, V. W. Laurie, and W. D. Gwinn, *J. Chem. Phys.* 33, 598 (1960).
21. W. Gordy, and R. L. Cook, *Microwave Molecular Spectra* (Wiley, New York, 1970) Chapt. 9.
22. J. R. Durig, K. L. Kizer, and Y. S. Li, *J. Amer. Chem. Soc.* 96, 7400 (1974).

23. G. Williams, N. L. Owen, and J. Sheridan, *Trans. Faraday Soc.* 67, 922 (1971).
24. J. S. Muentzer, *J. Chem. Phys.*, 48, 4544 (1968).
25. M. G. Griffin, "The Vibrational Assignment of Methylchloroformate", M. S. Thesis, Miami University, Oxford, Ohio, 1972.
26. J. D. Lewis, T. B. Malloy, Jr., T. H. Chao, and J. Laane, *J. Mol. Struct.* 12, 427 (1972).


Table I. The Microwave Spectra of Methylchloroformate

$^{35}\text{Cl}^{12}\text{C}^{16}\text{O}_2^{12}\text{CH}_3$		Observed	F' ← F		Calculated	Obs-Calc	Unperturbed Level	A or E
$J'_{k'_1 k'_2} ← J_{k_1 k_2}$								
$3_{03} ← 2_{02}$	14366.90	9/2 ← 7/2	14366.76	0.14	14366.21	A		
	14363.24	5/2 ← 3/2	14363.12	0.12		A		
$3_{13} ← 2_{12}$	13642.09	9/2 ← 7/2	13641.94	0.15	13640.52	A		
	13638.40	7/2 ← 5/2	13638.22	0.18		A		
$3_{12} ← 2_{11}$	15303.35	9/2 ← 7/2	15303.29	0.06	15301.74	A		
	15299.62	7/2 ← 5/2	15299.57	0.05		A		
$4_{04} ← 3_{03}$	19014.28	11/2+9/2 ← 9/2+7/2	19014.14	0.14	19013.84	A		
	19012.57	7/2+5/2 ← 5/2+3/2	19012.39	0.18		A		
$4_{14} ← 3_{13}$	18154.10	11/2 ← 9/2	18153.98	0.12	18153.33	A		
	18152.70	9/2 ← 7/2	18152.51	0.19		A		
	18152.15	7/2 ← 5/2	18151.77	0.38		A		
	18153.40	5/2 ← 3/2	18153.24	0.16		A		
$4_{13} ← 3_{12}$	20364.25	11/2 ← 9/2	20363.94	0.31	20363.35	A		
	20362.57	9/2 ← 7/2	20362.48	0.09		A		
	20362.11	7/2 ← 5/2	20361.65	0.46		A		
$4_{22} ← 3_{21}$	19605.73	11/2 ← 9/2	19605.76	-0.03	19594.42	A		
		9/2 ← 7/2	19599.69			A		
	19601.73	7/2 ← 5/2	19601.80	-0.07		A		
	19607.78	5/2 ← 3/2	19607.87	-0.09		A		
	19596.54	11/2 ← 9/2				E		
	19590.72	9/2 ← 7/2				E		
	19592.80	7/2 ← 5/2				E		
	5/2 ← 3/2			E				

Table I Continued

$^{35}\text{Cl}^{12}\text{C}^{16}\text{O}_2^{12}\text{CH}_3$		Observed	F'	←	F	Calculated	Obs-Calc	Unperturbed Level	A or E
$J'_{k'-1}k'+1$	$← J_{k-1}k+1$								
$5_{05} ← 4_{04}$		23553.51	13/2	←	11/2	23553.40	0.11	23553.25	A
		23552.58	9/2	←	7/2	23552.48	0.10		A
$5_{15} ← 4_{14}$		22640.36	13/2	←	11/2	22640.40	-0.04	22640.02	A
		22639.83	11/2	←	9/2	22639.68	0.15		A
		22639.41	9/2	←	7/2	22639.03	0.38		A
$5_{14} ← 4_{13}$		25387.83	13/2	←	11/2	25388.02	-0.19	25387.24	A
		25387.02	11/2	←	9/2	25387.32	-0.30		A
$5_{24} ← 4_{23}$		24082.85	13/2	←	11/2	24082.75	0.10	24081.70	A
		24080.12	11/2	←	9/2	24079.78	0.34		A
		24084.85	13/2	←	11/2			24083.74	E
		24082.19	11/2	←	9/2				E
$5_{23} ← 4_{22}$		24680.69	13/2	←	11/2	24680.41	0.28	24679.45	A
		24677.75	11/2	←	9/2	24677.32	0.43		A
		24678.69	13/2	←	11/2			24677.54	E
		24675.94	11/2	←	9/2				E
$6_{06} ← 5_{05}$		27981.37	15/2	←	13/2	27981.36	0.01	27981.18	A
		27980.77	11/2	←	9/2	27980.78	-0.01		A
$6_{16} ← 5_{15}$		27098.04	15/2	←	13/2	27098.08	-0.01	27097.80	A
		27097.79	13/2	←	11/2	27097.68	0.11		A
		27097.33	11/2	←	9/2	27097.18	0.15		A
$6_{15} ← 5_{14}$		30361.85	15/2	←	13/2	30361.83	0.02	30361.54	A
		30361.49	13/2	←	11/2	30361.46	0.03		A
		30361.08	11/2	←	9/2	30360.92	0.16		A

Table 1 Continued



$J'_{k-1}k'+1 \leftarrow J_{k-1}k+1$	Observed	F' $\leftarrow$	F	Calculated	Obs-Calc	Unperturbed Level	A or E
$6_{25} \leftarrow 5_{24}$	28841.62	15/2 $\leftarrow$	13/2	28841.50	0.12	28840.82	A
	28839.95	13/2 $\leftarrow$	11/2	28839.82	0.13		A
	28842.29	15/2 $\leftarrow$	13/2			28841.49	E
	28840.62	13/2 $\leftarrow$	11/2				E
$6_{24} \leftarrow 5_{23}$	29840.51	15/2 $\leftarrow$	13/2	29840.42	0.09	29839.65	A
	29838.72	13/2 $\leftarrow$	11/2	29838.09	0.03		A
	29839.98	15/2 $\leftarrow$	13/2			29839.14	E
	29838.23	13/2 $\leftarrow$	11/2				E
$7_{07} \leftarrow 6_{06}$	32313.35	17/2 $\leftarrow$	15/2	32313.31	0.04	32313.17	A
	32312.99	13/2 $\leftarrow$	11/2	32312.92	0.07		A
	32312.66	11/2 $\leftarrow$	9/2	32312.81	-0.15		A
$7_{17} \leftarrow 6_{16}$	31526.02	17/2 $\leftarrow$	15/2	31526.06	-0.04	31525.78	A
	31525.75	15/2 $\leftarrow$	13/2	31525.82	-0.07		A
	31525.48	13/2 $\leftarrow$	11/2	31525.43	0.05		A
$7_{16} \leftarrow 6_{15}$	35269.20	17/2 $\leftarrow$	15/2	35269.35	-0.15	35269.03	A
	35269.01	15/2 $\leftarrow$	13/2	35269.15	-0.14		A
	35268.82	13/2 $\leftarrow$	11/2	35268.93	-0.11		A
$7_{26} \leftarrow 6_{25}$	33529.92	17/2 $\leftarrow$	15/2	33569.88	0.04	33569.38	A
	33568.91	15/2 $\leftarrow$	13/2	33568.85	0.06		A
$7_{25} \leftarrow 6_{24}$	35064.18	17/2 $\leftarrow$	15/2	35064.29	-0.11	35063.55	A
	35062.98	15/2 $\leftarrow$	13/2	35063.15	-0.17		A
$7_{34} \leftarrow 6_{33}$	34143.18	17/2 $\leftarrow$	15/2	34143.41	-0.23	34142.13	A
	34140.89	15/2 $\leftarrow$	13/2	34140.96	-0.07		A
	34143.55	11/2 $\leftarrow$	9/2	34143.73	-0.18		A

Table I Continued

$^{35}\text{Cl}^{12}\text{C}^{16}\text{O}_2^{12}\text{CH}_3$							Unperturbed		
$J'_{k-1}k+1 \leftarrow J_{k-1}k+1$	Observed	$F'$	$\leftarrow$	$F$	Calculated	Obs-Calc	Level	A or E	
$8_{08} \leftarrow 7_{07}$	36578.20	19/2	$\leftarrow$	17/2	36578.36	-0.16	36578.11	A	
	36577.99	15/2	$\leftarrow$	13/2	56578.07	-0.08		A	
	36577.79	13/2	$\leftarrow$	11/2	36577.99	-0.20		A	
$8_{18} \leftarrow 7_{17}$	35924.89	19/2	$\leftarrow$	17/2	35925.15	-0.26	35924.70	A	
	35924.75	17/2	$\leftarrow$	15/2	35924.96	-0.21		A	
$8_{17} \leftarrow 7_{16}$	40092.32	19/2	$\leftarrow$	17/2	40092.53	-0.21	40092.09	A	
	40092.07	17/2	$\leftarrow$	15/2	40092.44	-0.37		A	
	40091.87	15/2	$\leftarrow$	13/2	40092.10	-0.23		A	
$8_{27} \leftarrow 7_{26}$	38263.35	19/2	$\leftarrow$	17/2	38263.48	-0.13	38263.00	A	
	38262.77	17/2	$\leftarrow$	15/2	38262.81	-0.04		A	
$6_{16} \leftarrow 5_{05}$	31148.83	15/2	$\leftarrow$	13/2	31148.73	0.10	31149.07	A	
	31150.00	13/2	$\leftarrow$	11/2	31149.86	0.14		A	
	31147.90	9/2	$\leftarrow$	7/2	31147.80	0.10		A	
	31146.81	15/2	$\leftarrow$	13/2				31147.00	E
	31147.90	13/2	$\leftarrow$	11/2					E
	31145.80	9/2	$\leftarrow$	7/2					E
$7_{17} \leftarrow 6_{06}$	34693.44	17/2	$\leftarrow$	15/2	34693.41	0.03	34693.55	A	
	34694.18	15/2	$\leftarrow$	13/2	34694.19	-0.01		A	
	34692.74	11/2	$\leftarrow$	9/2	34692.78	-0.04		A	
	34691.56	17/2	$\leftarrow$	15/2				34691.72	E
	34692.36	15/2	$\leftarrow$	13/2					E
	34690.93	11/2	$\leftarrow$	9/2					E



Table I Continued

$^{35}\text{Cl}^{12}\text{C}^{16}\text{O}_2^{12}\text{CH}_3$		Observed	F' +	F	Calculated	Obs-Calc	Unperturbed Level	A or E
$J'_{k'-1k'+1}$	$+ J_{k-1k+1}$							
$8_{18}$	$+ 7_{07}$	38305.20	19/2	$\leftarrow$ 17/2	38305.24	-0.04	38305.27	A
		38305.68	17/2	$\leftarrow$ 15/2	38305.77	-0.09		A
		38304.74	13/2	$\leftarrow$ 11/2	38304.79	-0.05		A
		38303.58	19/2	$\leftarrow$ 17/2			38303.61	E
		38304.01	17/2	$\leftarrow$ 15/2				E
		38303.06	13/2	$\leftarrow$ 11/2				E
<hr/>								
$^{37}\text{Cl}^{12}\text{C}^{16}\text{O}_2^{12}\text{CH}_3$								
$3_{03}$	$+ 2_{02}$	14034.80	9/2+7/2	$\leftarrow$ 7/2+5/2	14034.77	0.03	14034.24	A
		14031.90	5/2+3/2	$\leftarrow$ 3/2+1/2	14031.87	0.03		A
$6_{06}$	$+ 5_{05}$	27377.36	13/2	$\leftarrow$ 11/2	27377.32	0.04	27377.03	A
		27377.07	15/2	$\leftarrow$ 13/2	27377.18	-0.09		A
$6_{16}$	$+ 5_{15}$	26502.95	15/2	$\leftarrow$ 13/2	26502.90	0.05	26502.68	A
		26502.64	13/2	$\leftarrow$ 11/2	26502.60	0.04		A
$6_{15}$	$+ 5_{14}$	29630.17	15/2+13/2	$\leftarrow$ 13/2+11/2	29630.10	0.07	29630.03	A
$6_{24}$	$+ 5_{23}$	29085.75	15/2	$\leftarrow$ 13/2	29085.63	0.12	29085.07	A
		29084.33	13/2	$\leftarrow$ 11/2	29084.19	0.14		A
		29085.10	15/2	$\leftarrow$ 13/2			29084.50	E
		29083.84	13/2	$\leftarrow$ 11/2				E
$7_{07}$	$+ 6_{06}$	31629.18	15/2	$\leftarrow$ 13/2	31629.10	0.08	31628.94	A
		31629.00	17/2	$\leftarrow$ 15/2	31628.97	0.03		A
$7_{17}$	$+ 6_{16}$	30838.25	17/2+15/2	$\leftarrow$ 15/2+13/2	30838.27	-0.02	30838.15	A

Table I Continued

$^{37}\text{Cl}^{12}\text{C}^{16}\text{O}_2^{12}\text{CH}_3$		Observed	F'	←	F	Calculated	Obs-Calc	Unperturbed Level	A or E
$J'_{k'-1}k'+1$	← $J_{k-1}k+1$								
7 <sub>16</sub>	← 6 <sub>15</sub>	34430.71	17/2	←	15/2+15/2+13/2	34430.70	0.01	34430.59	A
7 <sub>25</sub>	← 6 <sub>24</sub>	34170.53	17/2	←	15/2	34170.54	-0.01	34170.04	A
		34169.60	19/2	←	17/2	34169.62	-0.02		A
8 <sub>08</sub>	← 7 <sub>07</sub>	35813.41	15/2	←	13/2	35813.40	0.01	35813.22	A
		35813.24	17/2	←	15/2	35813.28	-0.04		A
8 <sub>18</sub>	← 7 <sub>17</sub>	35146.57	19/2	←	17/2	35146.64	-0.07	35146.44	A
		35146.48	17/2	←	15/2	35146.54	-0.06		A
8 <sub>17</sub>	← 7 <sub>16</sub>	39155.34	19/2+17/2	←	17/2+15/2	39155.41	-0.07	39155.24	A
8 <sub>27</sub>	← 7 <sub>26</sub>	37382.34	19/2	←	17/2	37382.34	0.00	37382.06	A
		37381.88	17/2	←	15/2	37381.83	0.05		A
8 <sub>26</sub>	← 7 <sub>25</sub>	39287.67	19/2	←	17/2	39287.78	-0.11	39287.31	A
		39287.10	17/2	←	15/2	39287.16	-0.06		A
<hr/>									
$^{35}\text{Cl}^{12}\text{C}^{16}\text{O}_2^{12}\text{CD}_3$		Observed	F'	←	F	Calculated	Obs-Calc	Unperturbed Level	A or E
$J'_{k'-1}k'+1$	← $J_{k-1}k+1$								
6 <sub>06</sub>	← 5 <sub>05</sub>	25500.25	15/2	←	13/2	25500.18	0.07	25500.09	A
		25499.72	11/2	←	9/2	25499.63	0.09		A
6 <sub>15</sub>	← 5 <sub>14</sub>	27398.40	15/2	←	13/2	27398.34	0.06	27398.08	A
		27397.99	13/2+9/2	←	11/2+7/2	27397.90	0.09		A
		27397.59	11/2	←	9/2	27397.46	0.13		A
6 <sub>25</sub>	← 5 <sub>24</sub>	26128.59	15/2	←	13/2	26128.51	0.09	26127.84	A
		26127.04	13/2+11/2	←	11/2+9/2	26126.91	0.11		A

Table I Continued

$^{35}\text{Cl}^{12}\text{C}^{16}\text{O}_2^{12}\text{CD}_3$		Observed	$F' \leftarrow F$	Calculated	Obs-Calc	Unperturbed Level	A or E
$J'_{k'-1, k'+1} \leftarrow J_{k-1, k+1}$							
$6_{24} \leftarrow 5_{23}$		26855.95	$15/2 \leftarrow 13/2$	26855.88	0.07	26855.12	A
		26854.23	$13/2 \leftarrow 11/2$	26854.18	0.05		A
$7_{07} \leftarrow 6_{06}$		29494.20	$17/2 \leftarrow 15/2$	29494.17	0.03	29494.02	A
		29493.77	$15/2+13/2 \leftarrow 13/2+11/2$	29493.63	0.14		A
$7_{17} \leftarrow 6_{15}$		31858.88	$17/2 \leftarrow 15/2$	31858.92	-0.04	31858.67	A
		31858.70	$15/2 \leftarrow 13/2$	31858.72	-0.02		A
		31858.53	$11/2 \leftarrow 9/2$	31858.52	0.01		A
		31858.32	$13/2 \leftarrow 11/2$	31858.32	0.00		A
$7_{26} \leftarrow 6_{25}$		30426.94	$17/2 \leftarrow 15/2$	30427.02	-0.08	30426.42	A
		30425.91	$15/2+13/2 \leftarrow 13/2+11/2$	30425.98	-0.07		A
$7_{25} \leftarrow 6_{24}$		31530.17	$17/2 \leftarrow 15/2$	31530.23	-0.06	31529.60	A
		31529.03	$15/2+13/2 \leftarrow 13/2+11/2$	31529.08	-0.05		A
$7_{34} \leftarrow 6_{33}$		30829.38	$17/2 \leftarrow 15/2$	30829.49	-0.11	30828.38	A
		30827.08	$15/2 \leftarrow 13/2$	30827.18	-0.10		A
$6_{16} \leftarrow 5_{05}$		28881.94	$13/2 \leftarrow 11/2$	28881.95	-0.01	28881.03	A
		28880.78	$15/2 \leftarrow 13/2$	28880.79	-0.01		A
<hr/>							
$^{35}\text{Cl}^{12}\text{C}^{16}\text{O}_2^{13}\text{CH}_3$							
$4_{04} \leftarrow 3_{03}$		18566.43	$11/2+9/2 \leftarrow 9/2+7/2$	18566.42	0.01	18565.98	A
		18564.72	$7/2+5/2 \leftarrow 5/2+3/2$	18564.74	-0.02		A
$5_{05} \leftarrow 4_{04}$		23011.96	$13/2+11/2 \leftarrow 11/2+9/2$	23012.01	-0.05	23011.65	A
		23011.03	$9/2+7/2 \leftarrow 7/2+5/2$	23011.01	0.02		A

Table I Continued


$^{35}\text{Cl}^{12}\text{C}^{16}\text{O}_2^{13}\text{CH}_3$							Unperturbed	
$J'_{k'-1}k'+1$	$\leftrightarrow$	$J_{k-1}k+1$	Observed	$F' \leftrightarrow F$	Calculated	Obs-Calc	Level	A or E
$6_{25} \leftrightarrow 5_{24}$			28140.95	$15/2+9/2 \leftrightarrow 13/2+7/2$	28140.90	0.05	28140.16	A
			28139.31	$13/2+11/2 \leftrightarrow 11/2+9/2$	28139.30	0.01		A
			28141.67	$15/2+9/2 \leftrightarrow 13/2+7/2$			28140.88	E
			28140.02	$13/2+11/2 \leftrightarrow 11/2+9/2$				E
$6_{24} \leftrightarrow 5_{23}$			29054.61	$15/2+9/2 \leftrightarrow 13/2+7/2$	29054.53	0.08	29053.74	A
			29052.79	$13/2+11/2 \leftrightarrow 11/2+9/2$	29052.78	0.01		A
			29053.99	$15/2+9/2 \leftrightarrow 13/2+7/2$			29053.16	E
			29052.26	$13/2+11/2 \leftrightarrow 11/2+9/2$				E
$7_{26} \leftrightarrow 6_{25}$			32759.63	$17/2+11/2 \leftrightarrow 15/2+9/2$	32759.64	-0.01	32759.19	A
			32758.69	$15/2+13/2 \leftrightarrow 13/2+11/2$	32758.66	0.03		A
$7_{25} \leftrightarrow 6_{24}$			34133.36	$17/2+11/2 \leftrightarrow 15/2+9/2$	34133.46	-0.10	34132.88	A
			34132.30	$15/2+13/2 \leftrightarrow 13/2+11/3$	34132.33	-0.03		A
$8_{18} \leftrightarrow 7_{07}$			37646.46	$17/2 \leftrightarrow 15/2$	37646.63	-0.17	37646.14	A
			37646.12	$19/2+15/2 \leftrightarrow 17/2+13/2$	37646.10	0.02		A
			37645.69	$13/2 \leftrightarrow 11/2$	37645.55	0.14		A
<hr/>								
$^{35}\text{Cl}^{12}\text{C}^{16}\text{O}_2^{12}\text{CH}_2\text{D-OOP}$								
$3_{03} \leftrightarrow 2_{02}$			13923.95	$9/2+7/2 \leftrightarrow 7/2+5/2$	13923.88	0.07	13923.22	A
			13920.26	$5/2+3/2 \leftrightarrow 3/2+1/2$	13920.24	0.02		A
$3_{12} \leftrightarrow 2_{11}$			14783.69	$9/2+3/2 \leftrightarrow 7/2+1/2$	14783.50	0.19	14782.06	A
			14780.00	$7/2+5/2 \leftrightarrow 5/2+3/2$	14799.86	0.14		A

Table 1 Continued

$^{35}\text{C}1^{12}\text{C}^{16}\text{O}_2^{12}\text{CH}_2\text{D-OOP}$

$J'_{k-1}k'+1 \leftarrow J_{k-1}k+1$	Observed	F' $\leftarrow$ F	Calculated	Obs-Calc	Unperturbed Level	A or E
$4_{04} \leftarrow 3_{03}$	13923.95	9/2+7/2 $\leftarrow$ 7/2+5/2	13923.88	0.07	13923.22	A
	18440.64	7/2+5/2 $\leftarrow$ 5/2+3/2	18440.64	0.00		A
$4_{14} \leftarrow 3_{13}$	17632.27	11/2 $\leftarrow$ 9/2	17632.31	-0.04	17631.45	A
	17631.54	5/2 $\leftarrow$ 3/2	17631.55	-0.01		A
	17630.83	9/2 $\leftarrow$ 7/2	17630.86	-0.03		A
	17630.19	7/2 $\leftarrow$ 5/2	17630.10	0.09		A
$4_{13} \leftarrow 3_{12}$	19676.46	11/2 $\leftarrow$ 9/2	19676.52	-0.06	19675.65	A
	19675.62	5/2 $\leftarrow$ 3/2	19675.74	-0.12		A
	19675.13	9/2 $\leftarrow$ 7/2	19675.08	0.05		A
	19674.41	7/2 $\leftarrow$ 5/2	19674.30	0.11		A
$5_{05} \leftarrow 4_{04}$	22865.86	13/2+11/2 $\leftarrow$ 11/2+9/2	22865.86	0.00	22865.56	A
	22864.88	9/2+11/2 $\leftarrow$ 7/2+5/2	22864.84	0.04		A
$5_{14} \leftarrow 4_{13}$	24538.20	13/2 $\leftarrow$ 11/2	24538.28	-0.08	24537.72	A
	24537.62	11/2+7/2 $\leftarrow$ 9/2+5/2	24537.60	0.02		A
	24536.92	9/2 $\leftarrow$ 7/2	24536.91	0.01		A
$5_{24} \leftarrow 4_{23}$	23325.99	13/2+7/2 $\leftarrow$ 11/2+5/2	23326.04	-0.05	23324.54	A
	23323.10	11/2+9/2 $\leftarrow$ 9/2+7/2	23323.12	-0.02		A
$8_{18} \leftarrow 7_{07}$	37471.78	17/2 $\leftarrow$ 15/2	37471.73	0.05	37471.34	A
	37471.29	19/2 $\leftarrow$ 17/2	37471.28	0.01		A
	37470.81	13/2 $\leftarrow$ 11/2	37470.85	-0.04		A

Table I Continued



$J'_{k'_+} + J_{k_+}$	Observed	F' + F	Calculated	Obs-Calcd	Unperturbed Level	A or E
$3_{12} + 2_{11}$	14686.41	9/2 + 7/2	14686.29	0.12	14684.90	A
	14682.92	7/2+5/2+5/2+3/2	14682.61	0.31		A
$6_{15} + 5_{14}$	29144.92	15/2 + 13/2	29144.83	0.09	29144.66	A
	29144.56	13/2 + 11/2	29144.44	0.12		A
	29144.23	11/2 + 9/2	29143.93	0.30		A
$6_{24} + 5_{23}$	28628.59	5/2+9/2+3/2+7/2	28628.54	0.05	28627.74	A
	28626.87	13/2+11/2+11/2+9/2	28626.81	0.06		A
$6_{25} + 5_{24}$	27693.97	15/2+9/2+13/2+7/2	27694.00	-0.03	27693.20	A
	27692.38	13/2+11/2+11/2+9/2	27692.34	0.04		A
$7_{25} + 6_{24}$	33637.46	17/2+11/2+15/2+9/3	33637.45	0.01	33636.96	A
	33636.40	15/2+13/2+13/2+11/3	33636.37	0.03		A
$7_{26} + 6_{25}$	32236.10	17/2+11/2+15/2+9/2	32236.12	-0.02	32235.60	A
	32235.17	15/2+13/2+13/2+11/2	32235.11	0.06		A
$8_{26} + 7_{25}$	38675.55	19/2+13/2+17/2+11/2	38675.60	-0.05	38675.24	A
	38674.86	17/2+15/2+(15/2+13/2	38674.92	-0.06		A
$8_{27} + 7_{26}$	36745.89	19/2+13/2+13/2+11/2	36745.98	-0.09	36745.57	A
	36745.19	17/2+15/2+15/2+13/3	36745.30	-0.11		A
$7_{17} + 6_{06}$	33426.72	15/2 + 13/2	33426.76	-0.07	33426.23	A
	33426.19	17/2 + 15/2	33426.11	0.08		A
$8_{18} + 7_{07}$	36889.70	17/2 + 15/2	36889.72	-0.02	36889.31	A
	36889.25	19/2+15/2+17/2+13/2	36889.29	-0.04		A
	36888.81	13/2 + 11/2	36888.86	-0.05		A

TABLE II. Rotational constants, moments of inertia and nuclear quadrupole coupling constants for methylchloroformate

	$^{35}\text{Cl}^{12}\text{C}^{16}\text{O}_2^{12}\text{CH}_3$	$^{37}\text{Cl}^{12}\text{C}^{16}\text{O}_2^{12}\text{CH}_3$	$^{35}\text{Cl}^{12}\text{C}^{16}\text{O}_2^{12}\text{CD}_3$	$^{35}\text{Cl}^{12}\text{C}^{16}\text{O}_2^{13}\text{CH}_3$	$^{35}\text{Cl}^{12}\text{C}^{16}\text{O}_2^{12}\text{CH}_2\text{D-OOP}$	$^{35}\text{Cl}^{12}\text{C}^{16}\text{O}_2^{12}\text{CH}_2\text{D-IP}$
A	$9765.82 \pm 0.20$	$9738.39 \pm 0.29$	$9172.25 \pm 0.18$	$9738.40 \pm 0.30$	$9619.16 \pm 0.15$	$9465.36 \pm 0.30$
B	$2692.17 \pm 0.01$	$2623.00 \pm 0.01$	$2413.68 \pm 0.01$	$2620.09 \pm 0.01$	$2594.66 \pm 0.01$	$2582.49 \pm 0.01$
C	$2138.03 \pm 0.01$	$2092.92 \pm 0.01$	$1957.65 \pm 0.01$	$2091.18 \pm 0.01$	$2082.27 \pm 0.01$	$2054.86 \pm 0.01$
$I_a$	$51.7498 \pm 0.0011$	$51.8955 \pm 0.0016$	$55.0987 \pm 0.0011$	$51.8955 \pm 0.0016$	$52.5388 \pm 0.0009$	$53.3925 \pm 0.0010$
$I_b$	$187.722 \pm 0.001$	$192.671 \pm 0.001$	$209.381 \pm 0.001$	$192.886 \pm 0.001$	$194.777 \pm 0.001$	$195.694 \pm 0.001$
$I_c$	$236.376 \pm 0.001$	$241.471 \pm 0.001$	$258.156 \pm 0.001$	$241.672 \pm 0.001$	$242.706 \pm 0.001$	$245.943 \pm 0.001$
$\Delta$	3.096	3.097	6.324	3.110	4.610	
$\chi_{aa}$	$-59.8 \pm 0.5$	$046.4 \pm 0.5$	$-56.7 \pm 0.5$	$-57.2 \pm 0.5$	$-58.6 \pm 0.5$	$-56.2 \pm 0.5$
$\chi_{bb}$	$31.2 \pm 1.0$	$25.4 \pm 1.0$	$29.8 \pm 1.0$	$30.9 \pm 1.0$	$29.7 \pm 1.0$	$29.6 \pm 1.0$
$\chi_{cc}$	$28.6 \pm 1.0$	$21.0 \pm 1.0$	$26.9 \pm 1.0$	$26.3 \pm 1.0$	$28.9 \pm 1.0$	$26.6 \pm 1.0$

TABLE III

Coordinates of the atoms for which isotopic substitutions were made<sup>a</sup>

	Chlorine	Carbon (methyl)	Hydrogen (in plane)	Hydrogen (out of plane)
X	-1.590	2.278	2.808	2.510
Y	-0.278	-0.381	-1.312	0.182
Z	0.014 <sup>b</sup>	0.084 <sup>b</sup>	0.161 <sup>b</sup>	±0.887

<sup>a</sup>Error limits on the coordinates are never worse than  $\pm 0.005$ .

<sup>b</sup>The Z-coordinate of these atoms was taken as 0.0 for subsequent calculations.



TABLE IV

Coordinates of the unsubstituted atoms calculated from the moment relations

	Carbon (carbonyl)	Oxygen (carbonyl)	Oxygen (ether)
X	0.034 <sup>a</sup>	0.353	0.895
Y	0.312	1.458	-0.740
Z	0.0 <sup>b</sup>	0.0 <sup>b</sup>	0.0 <sup>b</sup>

<sup>a</sup>This value was chosen so that the carbonyl bond length would equal 1.190 Å.

<sup>b</sup>These values were assumed based on a consideration of the inertial defect.

TABLE V

Structural parameters for methylchloroformate<sup>a</sup>

	Bond Lengths ( $\text{\AA}$ )	Bond Angles ( $^{\circ}$ )
Cl-C	1.73	Cl-C=O 126
C=O	1.19	O=C-O 125
(O=)C-O	1.36	C-O-C 115
O-C(H <sub>3</sub> )	1.43	O-C-H(IP) 105
C-H(IP)	1.07	O-C-H(OOP) 110
C-H(OOP)	1.08	H-C-H(OOP-OOP) 111
		H-C-H(IP-OOP) 110

<sup>a</sup>Calculated from the coordinates in Table III and IV. The errors associated with these parameters because of the assumptions involved in their calculation are probably no worse than  $\pm 0.02 \text{ \AA}$  for the bond lengths and  $\pm 2^{\circ}$  for the bond angles.

TABLE VI

Internal rotation splitting (MHz) and parameters of methylchloroformate

Transition	<sup>35</sup> Cl	<sup>12</sup> C	<sup>16</sup> O <sub>2</sub>	<sup>12</sup> CH <sub>3</sub>	<sup>37</sup> Cl	<sup>12</sup> C	<sup>16</sup> O <sub>2</sub>	<sup>12</sup> CH <sub>3</sub>	<sup>35</sup> Cl	<sup>12</sup> C	<sup>16</sup> O <sub>2</sub>	<sup>13</sup> CH <sub>3</sub>
	$\nu_E - \nu_A$	$V_3$ (cal/mol)		$\nu_E - \nu_A$	$V_3$ (cal/mol)		$\nu_E - \nu_A$	$V_3$ (cal/mol)				
4 <sub>22</sub> ← 3 <sub>21</sub>	-9.06	1190										
5 <sub>24</sub> ← 4 <sub>23</sub>	2.04	1230										
5 <sub>23</sub> ← 4 <sub>22</sub>	-1.91	1240										
6 <sub>25</sub> ← 5 <sub>24</sub>	0.67	1230					0.72	1230				
6 <sub>24</sub> ← 5 <sub>23</sub>	-0.51	1270		-0.57	1270		-0.58	1270				
6 <sub>16</sub> ← 5 <sub>05</sub>	-2.09	1220										
7 <sub>17</sub> ← 6 <sub>06</sub>	-1.83	1220										
8 <sub>18</sub> ← 7 <sub>07</sub>	-1.66	1220										
$I_\alpha$	2.954 amu · Å <sup>2</sup>		2.954 amu · Å <sup>2</sup>		2.955 amu · Å <sup>2</sup>							
F	171.1 GHz <sub>3</sub>		171.1 GHz <sub>3</sub>		171.0 GHz <sub>3</sub>							
$\alpha$	0.05598		0.05945		0.05926							
$\beta$	0.00432		0.00439		0.00456							
$\gamma$	0.0		0.0		0.0							
$V_3$	1230 ± 30 cal/mol											

TABLE VII

Stark coefficients  $[(\text{MHz cm}^2)/\text{V}^2]$  and dipole moments of methylchloroformate<sup>a</sup>

Transition	M	$\Delta\nu/E^2$ ( $\times 10^5$ ) <sup>c</sup>	
		Obsd.	Calcd.
$5_{05} \leftarrow 4_{04}$	0	0.09	0.07
	3	0.28	0.41
	4	0.59	0.67

$$|\mu_a| = 1.7 \pm 0.2$$

$$|\mu_b| = 1.0 \pm 0.2$$

$$|\mu_c| = 0.0^a$$

$$|\mu_t| = 2.0 \pm 0.2$$

a.) By symmetry



Table VIII. Con't.

Vapor				Solid				Assignment		
Infrared <sup>a</sup>		Raman		Infrared <sup>a,b</sup>		Raman <sup>c</sup>				
cm <sup>-1</sup>	type	Δcm <sup>-1</sup>	Rel Int.	A'	cm <sup>-1</sup>	A''	Rel Int	Δcm <sup>-1</sup>	Rel Int	
								2640	vvw	
2610	?				2601	w		~2603	vvw	A' ν <sub>3</sub> + ν <sub>9</sub> (2599)
					2582	w		~2592	vvw	
					2570	w	2570 w	2577	vw	A' 2ν <sub>6</sub> (2578)
					2418	w				
2402					2406	w				A' 2ν <sub>7</sub> (2408)
					2371	vw				A'
					2316	m	2316 m			A' 2ν <sub>15</sub> (2304)
								2297	vw	A' 2ν <sub>15</sub> (2296)
								2273	vw	A'/A'' ν <sub>4</sub> /ν <sub>14</sub> + ν <sub>9</sub> (2271)
								2178	w	A' ν <sub>3</sub> + ν <sub>11</sub> (2183)
					2142	w				
								2120	vvw	A' ν <sub>7</sub> + ν <sub>8</sub> (2116)
					2095	w		~2100	vvw	A'' ν <sub>8</sub> + ν <sub>15</sub> (2099)
					2089	w		2079	vvw	
2027					2025	vw				A' ν <sub>7</sub> + ν <sub>9</sub> (2025)
2022										
1973	?				1960	vw				A'' ν <sub>3</sub> + ν <sub>17</sub> (1956)
1924					1905	w				
					1890	w				A' 2ν <sub>8</sub> (1894)
					1882	ms		~1880	vvw	

Table VIII. Con't.

Vapor				Solid				Assignment				
Infrared <sup>a</sup>		Raman		Infrared <sup>a,b</sup>		Raman <sup>c</sup>						
cm <sup>-1</sup>	type	Δcm <sup>-1</sup>	Rel Int	cm <sup>-1</sup>	A' Rel Int	A'' cm <sup>-1</sup>	Rel Int	Δcm <sup>-1</sup>	Rel Int			
1806	B	1601	p	1789	s			1774	w	ν <sub>3</sub> C=O stretch		
1794		1792	p	to				1757	mw			
1788				1755				1752	m			
				1725		m						
				1716		s			1717		vw	A' ν <sub>12</sub> + ν <sub>5</sub> (1717)
				1646		mw			1652		vw	A' 2ν <sub>9</sub> (1642)
			1622	mw			~1625	vw	A' ν <sub>14</sub> + ν <sub>17</sub> (1622)			
1464	?	1466	p					1458	m	A'/A'' ν <sub>4</sub> /ν <sub>14</sub> CH <sub>3</sub> asymmetric deformation		
1459								1450	mw			
1448			1459	dp	1448.6	s	1449.2	s				
1441	?											
1433												
				1431	s	1431	sh	1432	mw	A' ν <sub>5</sub> CH <sub>3</sub> symmetric deformation		
1396	?							~1405	vw			
1382			1389	p	1387	s	1387	vw		A' 2ν <sub>16</sub> (1378)		
					1326	mw				A' ν <sub>15</sub> + ν <sub>17</sub> (1322)		
1313	AB											
1308					1306	s	1306	w		A' ν <sub>9</sub> + ν <sub>10</sub> (1302)		
1300					1297	s				A' ν <sub>9</sub> + ν <sub>10</sub> (1296)		

Table VIII. Con't.

Vapor				Solid				Assignment	
Infrared <sup>a</sup>		Raman		Infrared <sup>a,h</sup>		Raman <sup>c</sup>			
cm <sup>-1</sup>	type	Δcm <sup>-1</sup>	Rel Int	cm <sup>-1</sup>	Rel Int	cm <sup>-1</sup>	Rel Int	Δcm <sup>-1</sup>	Rel Int
				A'	A''				
1233				1250		1250	w		
1210				to		1226	vw		
1202	?	1157	p	1210	s	1211	vw	1204	mw
1196									A' ν <sub>6</sub> COC asymmetric stretch
				1186	s	1180	w		
1162	AB			1174	s			1169	mw
1154		1124	dp	1152	sh	1152	ms	1152	w
				1142	m	1142	w	1148	w
								1144	sh
1060								1087	vvw
								1056	vvw
1036								1046	vvw
1007									
								988	vw
				961	w			972	vw
				958	w			960	vw
971		977	p					951	sh
966	?	962	p	947	s	947	w	947	m
957								932	vvw
								924	vvw
				920	w			918	vvw
892				889	mw			~893	vvw
									A' ν <sub>10</sub> + ν <sub>11</sub> (895)



Table VIII. Con't.

Vapor				Solid				Assignment		
Infrared <sup>a</sup>		Raman		Infrared <sup>a,b</sup>		Raman <sup>c</sup>				
cm <sup>-1</sup>	type	Δcm <sup>-1</sup>	Rel Int	cm <sup>-1</sup>	Rel Int	cm <sup>-1</sup>	Rel Int	Δcm <sup>-1</sup>	Rel Int	
882				881	w					
830	A	823	p	825	s	825	w			
824				821	s	821	w	821	ms	A' ν <sub>9</sub> C-Cl stretch
816				817	ms					
				758	w			766	vw	
706	C			689	w	689	s	702	vvw	A'' ν <sub>16</sub> C-Cl bend
694				672	w	672	m			
677										
668										
542										
527								518	vvw	
494	A	487	p	481	s	481	vw	486	vs	A' ν <sub>10</sub> OCO bend
488				477	m	477	vw	479	ms	
477				472	p					
413	AB	406	p	407	s	407	w	409	ms	A' ν <sub>11</sub> C-Cl bend
405										
397										
								392	vvw	A' 2ν <sub>17</sub> (388)
354								367	vvw	A' 2ν <sub>17</sub> (364)

Table VIII. Con't.

Vapor				Solid				Assignment		
Infrared <sup>a</sup>		Raman		Infrared <sup>a,b</sup>		Raman <sup>c</sup>				
cm <sup>-1</sup>	type	Δcm <sup>-1</sup>	Rel Int	A'	cm <sup>-1</sup>	Rel Int	A''	cm <sup>-1</sup>	Rel Int	
325								341	vvw	
314	?									
304										
279	?	274	p	270	s			283	w	A' ν <sub>12</sub> COC bend
268								277	w	
Methylchloroformate-d <sub>3</sub>										
Infrared Vapor										
163.2	s	148.8	s					202	vs	
		148.2	s					194	w	A'' ν <sub>17</sub> Methoxy torsion
								182	w	
129.2	w	99.0	w					135	vvw	A'' ν <sub>18</sub> Methyl torsion
								122	ms	Lattice mode
								100	sh	Lattice mode
								94	s	Lattice mode
								81	w	Lattice mode
								67	mw	Lattice mode
								60	w	Lattice mode
								52	mw	Lattice mode
								45	m	Lattice mode

a.) Taken from Ref. 2<sup>7</sup>


b.) Recorded at ~77°K

c.) Recorded at 187°K


d.) vw = very weak, w = weak, m = medium, s = strong,  
vs = very strong, b = broad, p = polarized, dp = depolarized

## FIGURE CAPTIONS


- Fig. 1 Possible conformations of methylchloroformate.
- Fig. 2 The structure of methylchloroformate in the principal axis system.
- Fig. 3 The Raman spectra of methylchloroformate. [A] The room temperature vapor. [B] The solid at 18.7°K.
- Fig. 4 The far infrared spectra of [A] methylchloroformate and [B] methylchloroformate-d<sub>3</sub>. Transitions marked with arrows correspond to a small amount of HCl impurity.






S - CIS



S - TRANS







## APPENDIX II

### SPECTRA AND STRUCTURE OF ORGANOPHOSPHORUS COMPOUNDS. XVII.

#### INFRARED AND RAMAN SPECTRA, VIBRATIONAL ASSIGNMENT AND BARRIERS TO INTERNAL ROTATION FOR t-BUTYLPHOSPHINE

##### Abstract

The infrared spectra of gaseous and solid tertiary-butylphosphine,  $[(\text{CH}_3)_3\text{CPH}_2]$ , have been recorded from  $50\text{ cm}^{-1}$  to  $3500\text{ cm}^{-1}$ . The Raman spectra of gaseous, liquid and solid  $(\text{CH}_3)_3\text{CPH}_2$  have been recorded from 10 to  $3500\text{ cm}^{-1}$ . A vibrational assignment of the 42 normal modes has been made. A harmonic approximation of the methyl torsional barrier from observed transitions in the solid state gave a result of 4.22 kcal/mole and 3.81 kcal/mole in the gaseous state. Hot band transitions for the phosphino torsional mode have been observed. The potential function for internal rotation about the C-P bond has been calculated. The two potential constants were determined to be:

$$V_3 = 2.79 \pm 0.01 \text{ kcal/mole and } V_6 = 0.07 \pm 0.01 \text{ kcal/mole.}$$



## INTRODUCTION

Vibrational spectroscopy has been shown to be a valuable tool in the determination of intramolecular potential functions governing internal rotation about single bonds.<sup>1, 2</sup> This method has recently been extended to the study of asymmetric tops rotating against a molecular frame, as opposed to the more familiar symmetric tops such as  $-\text{CH}_3$  and  $-\text{SiH}_3$ .<sup>3</sup> If the molecular frame is asymmetric as well, it is possible for there to be spectroscopically distinct rotational isomers or conformers. Both ethylphosphine ( $\text{CH}_3\text{CH}_2\text{PH}_2$ ) and isopropylphosphine [ $(\text{CH}_3)_2\text{CHPH}_2$ ] are examples of this type of system since the  $\text{PH}_2$  group constitutes an asymmetric internal rotor. The internal rotation barrier for the  $\text{PH}_2$  top has been determined for both of these molecules.<sup>4, 5</sup> If, however, the molecular frame is symmetric, as is the case in tertiary-butylphosphine, [ $(\text{CH}_3)_3\text{CPH}_2$ ], only one conformation is possible and the potential function governing internal rotation is considerably simplified, becoming three-fold symmetric as in the case of methylphosphine ( $\text{CH}_3\text{PH}_2$ ). Thus, the determination of this potential function for t-butylphosphine is a natural extension of the prior studies of ethyl-<sup>4</sup> and isopropylphosphine.<sup>5</sup> Also, it would be of interest to determine the barriers to internal rotation of the methyl groups in this molecule, as part of continuing studies<sup>6, 7</sup> of the top-top interactions in such "three-top" molecules.

Finally, the assignment of the vibrational spectrum of t-butylphosphine would finish the series of singly alkyl-substituted phosphines: methyl, ethyl and isopropyl. Group frequencies determined

for the  $\text{PH}_2$  moiety from these studies could be expected to apply to almost any hydrocarbon derivative.

### EXPERIMENTAL

In general, all sample preparations and manipulations were performed using standard high vacuum techniques where possible. This was done in view of the relative instability of trivalent phosphorous compounds with respect to oxidation as well as to minimize exposure to the compounds, which are probably quite toxic and undeniably unpleasant smelling. Sample purifications were performed using fractional condensation (trap-to-trap) methods or more generally a low temperature, low pressure fractionating column.

The sample of  $(\text{CH}_3)_3\text{CPH}_2$  was prepared by the reduction of  $(\text{CH}_3)_2\text{CHPCl}_2$  with  $\text{LiAlH}_4$  in di-n-butyl ether. After low temperature distillation, the purity of the  $(\text{CH}_3)_3\text{CPH}_2$  was checked by mass spectrometry and  $^1\text{H}$ ,  $^{13}\text{C}$  and  $^{31}\text{P}$  NMR.

Mid-infrared spectra of  $(\text{CH}_3)_3\text{CPH}_2$  were obtained using a Digilab FTS-15B Fourier transform interferometric spectrometer. A germanium beamsplitter on a KBr substrate gave spectral coverage in the range from  $3800\text{ cm}^{-1}$  to  $400\text{ cm}^{-1}$ . Spectra of the gaseous phase were obtained by using a 10 cm cell with KBr windows and typically less than 10 torr of sample pressure.

Atmospheric water vapor was removed from the spectrometer housing by purging with dry nitrogen. Since the FTS system uses a He-Ne laser as a frequency reference of relatively high stability, calibration is achieved by adjusting a software parameter, the laser wave length, to reproduce an observed vibrational transition of a suitable standard at its correct frequency. As a result, the accuracy of a measured frequency can be expected to be at least  $0.1\text{ cm}^{-1}$  throughout the spectrum.

Both apodization function and resolution were varied for each sample to attain the best quality spectrum, but a typical experiment would be run at an effective resolution of better than  $0.5 \text{ cm}^{-1}$  using a modified "boxcar" apodization function for photometric accuracy.

Far infrared spectra were obtained by using the FTS-15B and substituting a mylar beamsplitter for the germanium one to obtain a spectral range of 400 to  $80 \text{ cm}^{-1}$ . A high pressure Hg arc lamp source was substituted for the glower used in the mid-infrared, but otherwise the operation of the instrument was the same as described above. Spectra of compounds in the gas phase were obtained by using 10 cm cells equipped with polyethylene windows. Typical sample pressures ranged from 100 to 300 torr, and spectra were taken at various resolutions between  $0.5$  and  $2.0 \text{ cm}^{-1}$ . The spectrum of the solid phase sample was obtained by condensing the sample onto a wedged silicon plate cooled to  $\sim 15^\circ\text{K}$  by a closed cycle He refrigerator.

Raman spectra were recorded to  $4000 \text{ cm}^{-1}$  using a Cary 82 spectrometer equipped with either a Spectra Physics model 171 or a Coherent Radiation Labs model 53A argon ion laser operating on the  $5145 \text{ \AA}$  line. Spectra of the gaseous phase were obtained using the Cary multipass accessory at sample pressures of  $\sim 300$  torr. The spectrum of the liquid was recorded from samples in sealed glass capillaries. The spectrum of the solid was obtained by condensing the sample on a copper block maintained at  $\sim -190^\circ\text{C}$  by boiling nitrogen. Polarization measurements were made using the standard Cary accessories. Frequencies measured for sharp, resolvable bands are expected to be accurate to at least  $\pm 2 \text{ cm}^{-1}$ .

## VIBRATIONAL ASSIGNMENT

Tertiary-butylphosphine was assumed to adopt a fully staggered molecular conformation, and as such would possess a single plane of symmetry and belong to the point group  $C_s$ . However, if one ignores the hydrogen atoms attached to the phosphorus atom, the molecular symmetry would be  $C_{3v}$ . If the major portion of the molecule follows this "local" symmetry, the assignment of the vibrational spectra would be somewhat simplified. Pritchard and Nelson<sup>8</sup> were able to analyze the spectrum of t-butanol using the higher symmetry model for all but a few vibrational modes. Hence for  $(CH_3)_3CP$  there would be 24 normal modes, distributed  $8A_1$ ,  $4A_2$  and  $12E$ . Under  $C_s$  symmetry,  $(CH_3)_3CPH_2$  would exhibit  $23A'$  and  $19A''$  vibrational modes. Spectra taken of  $(CH_3)_3CPH_2$  in the fluid phases might be expected then to show groups of unresolved bands. However, spectra taken of the sample in the solid phase should allow observation of all 42 fundamentals since in the crystal closely spaced fundamentals should be resolvable. Where applicable, the origin of the vibrational mode under the hypothetical  $C_{3v}$  symmetry will be indicated in parenthesis as a guide to the assignment. To aid in the analysis of gas phase band contours, the principal moments of inertia for  $(CH_3)_3CPH_2$  were calculated using a reasonable assumed structure. These calculations indicate that the molecule is very nearly a symmetric top, with the A and C inertial axes in the symmetry plane. Thus the  $A'$  vibrations are expected to show A/C

hybrid band contours with a sharp Q branch, and A" vibrations B type bands with no central maximum. The vibrational frequencies given in Table I for the A" modes in the infrared spectra of gaseous  $(\text{CH}_3)_3\text{CPH}_2$  therefore reflect estimates of the band center. The vibrational assignments were guided by those of Pritchard and Nelson<sup>8</sup> for  $(\text{CH}_3)_3\text{COH}$  and those of Evans and Lo<sup>9</sup> for  $(\text{CH}_3)_3\text{CCl}$ .

The Raman spectra of  $(\text{CH}_3)_3\text{CPH}_2$  in the gaseous, liquid and solid phases are shown in Fig. 1, and the mid-infrared spectra of  $(\text{CH}_3)_3\text{CPH}_2$  in the gaseous and solid phases are shown in Fig. 2. Fig. 3 depicts the far-infrared spectra of gaseous and solid  $(\text{CH}_3)_3\text{CPH}_2$ . The observed frequencies and their assignments are given in Table I.

The C-H stretching region, 2870 to 3000  $\text{cm}^{-1}$ , showed a dense cluster of bands in all sample phases in both the Raman and infrared spectra. In addition to the 9 fundamentals expected in this region, Durig, *et al.*,<sup>10</sup> have pointed out that additional strong bands may be expected to occur in this region, attributable to overtones of the antisymmetric methyl deformations in Fermi resonance with the stretching fundamentals. The bands in this region were not resolved even in the spectra of the solid phase, and no additional isotopic species were available for study. In view of these difficulties, attempts to assign this portion of the vibrational spectra were abandoned and the observed band maxima tabulated.

The two very strong bands at 2293  $\text{cm}^{-1}$  and 2278  $\text{cm}^{-1}$  in the spectra of the solid sample are undoubtedly the antisymmetric and symmetric P-H stretches,  $\nu_{28}$  and  $\nu_6$ , respectively. As previous experience has indicated,<sup>4, 5</sup> only the A' mode is seen in the Raman spectra of the gas phase.

The advantage of recording spectra of the solid phase is easily seen in the analysis of the  $\text{CH}_3$  antisymmetric deformations  $\nu_{29}$ ,  $\nu_7$  (E),  $\nu_{30}$ ,  $\nu_8$  (E),  $\nu_9$  ( $A_1$ ) and  $\nu_{31}$  ( $A_2$ ). The Raman spectrum of gaseous  $(\text{CH}_3)_3\text{CPH}_2$  shows only a broad, featureless band centered near 1450, but the Raman spectrum of the solid clearly shows all six fundamentals. The assignment of the  $\text{CH}_3$  symmetric deformations,  $\nu_{10}$  ( $A_1$ ),  $\nu_{11}$  and  $\nu_{32}$  (E) is also straightforward, although these modes are not seen in the Raman spectra of the fluid phases.

Assignment of the vibrational spectrum in the region between 1200  $\text{cm}^{-1}$  and 800  $\text{cm}^{-1}$  is clearly the most difficult. Twelve fundamental vibrations; 6 methyl rocks, 3 C-C stretches and the 3  $\text{PH}_2$  deformations, are all expected to occur in this region. In addition, severe mixing of several vibrational modes, especially the C-C stretches and methyl rocking modes, could make relative intensities unreliable as assignment aids. Three fundamentals of this group should fall considerably lower than the rest;  $\nu_{17}$  ( $A_1$ ), the C-C symmetric stretch should be below 900  $\text{cm}^{-1}$ , as should  $\nu_{37}$ , the  $\text{PH}_2$  twist, and  $\nu_{18}$ , the  $\text{PH}_2$  wag. Only  $\nu_{17}$  and  $\nu_{18}$  should exhibit sharp Q branches in the gaseous phase Raman spectrum, and only three bands are apparent in this region in the Raman spectrum of the solid phase. Thus,  $\nu_{17}$ ,  $\nu_{37}$  and  $\nu_{18}$  are assigned to the bands at 836  $\text{cm}^{-1}$ , 818  $\text{cm}^{-1}$  and 795  $\text{cm}^{-1}$  in the Raman spectrum of the solid sample. The frequency of  $\nu_{15}$ , the  $\text{PH}_2$  scissoring mode, is well established<sup>4, 5, 11</sup> and is assigned to the band at 1074  $\text{cm}^{-1}$  in the mid-infrared spectrum of solid  $(\text{CH}_3)_3\text{CPH}_2$ . The C-C antisymmetric stretches,  $\nu_{33}$  and  $\nu_{12}$  (E), occurred near 1240  $\text{cm}^{-1}$  in both  $(\text{CH}_3)\text{CCl}^9$  and  $(\text{CH}_3)\text{COH}$ ;<sup>8</sup> however, there are no bands in this region for  $(\text{CH}_3)_3\text{CPH}_2$ .

There are two strong bands at  $1211\text{ cm}^{-1}$  and  $1192\text{ cm}^{-1}$  in the Raman spectrum of the solid, and have been assigned to  $\nu_{33}$  and  $\nu_{12}$ , respectively. The slightly lower, weaker band at  $1170$  is assigned to  $\nu_{13}$  ( $A_1$ ), a  $\text{CH}_3$  rocking mode. The complex series of bands seen on the low frequency side of  $\nu_{13}$  in the gas phase spectra are most probably hot bands due to the methyl torsions or phosphino torsion. Although this assignment of  $\nu_{13}$  may seem to place this rocking mode quite high, the corresponding modes in  $(\text{CH}_3)_3\text{COH}$ <sup>8</sup> and  $(\text{CH}_3)_3\text{CCl}$ <sup>9</sup> were assigned to bands at  $1189\text{ cm}^{-1}$  and  $1115\text{ cm}^{-1}$ , respectively. The sharp, polarized peak seen at  $1029\text{ cm}^{-1}$  in the Raman spectra of the gaseous and liquid phases must be due to the  $A'$  methyl rocking mode,  $\nu_{16}$ . The sharp, moderately intense Q branch observed at  $1090\text{ cm}^{-1}$  in the mid-infrared spectrum of gaseous  $(\text{CH}_3)_3\text{CPH}_2$  must then belong to the remaining  $A'$  methyl rocking mode,  $\nu_{14}$ . In terms of the  $C_{3v}$  model symmetry,  $\nu_{14}$  and  $\nu_{34}$  are derived from a degenerate E vibration, as are  $\nu_{16}$  and  $\nu_{35}$ . Therefore  $\nu_{34}$  and  $\nu_{35}$  which would be expected to be clearly discernable only in the solid phase spectra, should fall near their respective  $A'$  modes,  $\nu_{14}$  and  $\nu_{16}$ . On this basis,  $\nu_{34}$  is assigned to a band seen at  $1065\text{ cm}^{-1}$  in the spectra of the solid phase, and  $\nu_{35}$  to the band at  $1036\text{ cm}^{-1}$ . The remaining rocking mode  $\nu_{36}$  ( $A_2$ ) can then be assigned to the broad, depolarized band seen near  $940\text{ cm}^{-1}$ .

The assignment of  $\nu_{19}$ , the C-P stretch, is unambiguous, as it is usually<sup>4, 5</sup> the most intense band in the Raman spectrum below  $2000\text{ cm}^{-1}$ . In contrast, this band, located near  $590\text{ cm}^{-1}$ , is quite weak in the mid-infrared spectrum.

In the region below  $400\text{ cm}^{-1}$  there should be 5 skeletal deformations

as well as the 4 torsional vibrations. In the far infrared spectrum of gaseous  $(\text{CH}_3)_3\text{CPH}_2$  there are apparently two sharp Q branches at  $352\text{ cm}^{-1}$  and  $346\text{ cm}^{-1}$ , also seen in the Raman spectrum of the gas. However, this structure disappears upon condensation, and only a single, strong, sharp band is seen in the solid phase spectra at  $353\text{ cm}^{-1}$ . This band has been assigned to  $\nu_{21}$  ( $A_1$ ), the CCC symmetric deformation, with the  $346\text{ cm}^{-1}$  band assigned as a hot band. The other two CCC deformations,  $\nu_{38}$  and  $\nu_{20}$  (E), are assigned to the bands at  $394\text{ cm}^{-1}$  and  $389\text{ cm}^{-1}$ , respectively, in the Raman spectrum of the solid. The CCP deformational modes,  $\nu_{39}$  and  $\nu_{22}$  (E), have been assigned to the bands at  $297\text{ cm}^{-1}$  and  $295\text{ cm}^{-1}$ , respectively, in the Raman spectrum of the solid. The anomalously low intensity of  $\nu_{38}$ ,  $\nu_{20}$ ,  $\nu_{39}$  and  $\nu_{22}$  in the far infrared spectrum of solid  $(\text{CH}_3)_3\text{CPH}_2$  is possibly attributable to crystal effects. Following the assignments of Durig, et al.,<sup>7</sup> for a series of t-butyl compounds, the bands at  $278\text{ cm}^{-1}$ ,  $273\text{ cm}^{-1}$ , and  $248\text{ cm}^{-1}$  in the far infrared spectrum of the solid have been assigned to the methyl torsional modes,  $\nu_{23}$ ,  $\nu_{40}$  (E) and  $\nu_{41}$  ( $A_2$ ). The strong band system centered near  $176\text{ cm}^{-1}$  in the far infrared spectrum of the gas is undoubtedly due to the fundamental  $\text{PH}_2$  torsion,  $\nu_{42}$ , and its hot bands. This band becomes a single sharp band centered at  $200\text{ cm}^{-1}$  upon solidification.

#### DETERMINATION OF THE TORSIONAL BARRIERS

In order to treat the problem of the barriers to internal rotation of the methyl groups in t-butylphosphine, the assumption was made that, so far as the methyl torsions are concerned, the molecule has  $C_{3v}$



symmetry. Lide and Mann<sup>12</sup> have expressed the internal potential energy of such a system in a three dimensional Fourier expansion in terms of  $3n\phi_1$ ,  $3n\phi_2$ , and  $3n\phi_3$ , where  $\phi_j$  is the internal rotation angle of the  $j$ th top. If terms with  $n>1$  are ignored, the potential energy becomes:

$$2V = V_0 - V_1 \sum_{i=1}^3 \cos 3\phi_i - V_2 \sum_{i>j} \cos 3\phi_i \cos 3\phi_j - V_3 \cos 3\phi_1 \cos 3\phi_2 \cos 3\phi_3 + V_4 \sum_{i>j} \sin 3\phi_i \sin 3\phi_j + V_5 \sum_{i \neq j \neq k} \cos 3\phi_i \sin 3\phi_j \sin 3\phi_k$$

with the inclusion of the proper kinetic energy terms, the torsional frequencies are given in the harmonic approximation by:<sup>12</sup>

$$\nu_A = [2(F-F')(K + 2L)]^{1/2}$$

$$\nu_E = [2(F + 2F')(K - L)]^{1/2}$$

where

$$K = (9/2) (V_1 + 2V_2 + V_3)$$

$$L = (9/2)(V_4 + V_5)$$

The kinetic terms for the present case were obtained by averaging the terms for each top, which were only slightly different. The results of a calculation using this approximation are shown in Table II for both the solid phase and gas phase data. If the interaction constant  $L$  is interpreted as a measure of the forces between hydrogen atoms on different  $\text{CH}_3$  groups, then the negative sign found for  $L$  indicates that the net force is repulsive. Thus, the  $A_2$  - type motion, in which the three tops move in phase, is more effective in keeping the hydrogen atoms apart than the  $E$  torsion, where one top is out of phase. In view of the small value found for  $L$ , the cosine interaction terms are probably negligible also and  $(2/9)K$  is a good approximation of the total barrier.

To determine the barrier to internal rotation of the  $\text{PH}_2$  top, a computer program was used which is similar to one described by Lewis, et al.<sup>13</sup> It is assumed that the torsional potential can be represented as a Fourier cosine series in the internal rotation angle:

$$V(\phi) = \left(\frac{1}{2n}\right) V_n (1 - \cos n\phi).$$

Since the potential function must be three-fold symmetric, only terms where  $n=3$  and  $6$  were used. The band centers of the  $1 \leftarrow 0$  and  $2 \leftarrow 0$  phosphino torsions were estimated from the observed far infrared spectrum of the gaseous phase. At this point, the program was allowed to iterate on the two potential constants, and a prediction made for the location of the  $3 \leftarrow 2$  transition using the resulting potential function. A slight change in the band contour at  $162 \text{ cm}^{-1}$  had been noted previously and coincided reasonably well with the prediction. This band was then included in the program input and the potential function re-evaluated. The calculated  $4 \leftarrow 3$  transition splits by  $\approx 3 \text{ cm}^{-1}$  as the energy levels near the top of the well, due to quantum mechanical tunnelling.<sup>14</sup> This predicted transition roughly coincided with another contour change seen at about  $151 \text{ cm}^{-1}$ ; however, this transition was not included in the fit. The results of this calculation are shown in Table III.

#### DISCUSSION

With regard to the vibrational assignment, several small points are probably ambiguous, but overall it is felt that the given assignment is quite reasonable. A vibrational study of the  $(\text{CH}_3)_2\text{CPD}_2$  isotope would confirm the assignment of  $\text{PH}_2$  deformations, and might clarify the

assignment of the skeletal deformations if vibrational coupling of these modes is sufficient to cause an appreciable frequency shift.

The methyl barrier potential constants determined in this work may be compared with those found by Durig, *et al.*,<sup>7</sup> for some similar molecules observed in the solid state. The (2/9)K term for  $(\text{CH}_3)_3\text{COH}$  of 4.13 kcal/mole goes to 4.41 kcal/mole in  $(\text{CH}_3)_3\text{CSH}$ , while the (2/9)L terms are 0.44 and 0.49 kcal/mole, respectively. In comparison, the (2/9)K term for  $(\text{CH}_3)_3\text{CNH}_2$ , is 4.28 kcal/mole and 4.22 kcal/mole for  $(\text{CH}_3)_3\text{CPH}_2$ , with the (2/9)L terms 0.70 and 0.15 kcal/mole, respectively. The general agreement in the magnitude of (2/9)K is reassuring; however, the trends in the magnitude of (2/9)L for the amine and phosphine are not clearly understood. The magnitude of the (2/9)L term is derived from the frequency splitting of the pseudo-E torsional mode and the pseudo-A<sub>2</sub> torsional mode. Differing degrees of intermolecular effects in the solid, such as hydrogen bonding, might be expected to produce some variation in this interaction term, but the trend should then also be apparent in the  $(\text{CH}_3)_3\text{COH} \rightarrow (\text{CH}_3)_3\text{CSH}$  series. Similar arguments concerning lone pair effects or lengths of the tertiary carbon to unique substituent bond should be evident in the  $(\text{CH}_3)_3\text{COH} \rightarrow (\text{CH}_3)_3\text{CSH}$  series, also. The most probable explanation is that the methyl torsional modes in  $(\text{CH}_3)_3\text{CNH}_2$  are severely coupled with the amine torsional vibration which falls between the two methyl torsional regions. Thus the large interaction term seen for  $(\text{CH}_3)_3\text{CNH}_2$  is due in part to interactions of the methyl torsions with the NH<sub>2</sub> torsion.

The barrier to internal rotation of the PH<sub>2</sub> group in  $(\text{CH}_3)_3\text{CPH}_2$ , 2.79 kcal/mole, is somewhat higher than that determined by Tsuboi,

et al.,<sup>15</sup> for  $(\text{CH}_3)_3\text{CNH}_2$ , 2.36 kcal/mole. The barrier for t-butylphosphine is nearly a kilocalorie/mole higher than the barrier for  $\text{CH}_3\text{PH}_2$ ,<sup>11</sup> determined to be 1.96 kcal/mole. Similarly the barrier in  $\text{CH}_3\text{NH}_2$  is 1.95 kcal/mole.<sup>16</sup> From these results, one might infer that electrostatic interactions of the PH bonds and phosphorus lone pair are greater with CC bonds than with CH bonds, if these interactions are predominantly repulsive.

#### ACKNOWLEDGEMENT

The authors gratefully acknowledge the financial support given this study by the National Aeronautics and Space Administration through grant NGL-41-002-003. Acknowledgement is also given to the National Science Foundation for funds to purchase the Digilab FTS-15B interferometer by grant MPS-75-06926.

## BIBLIOGRAPHY

1. J. R. Durig, S. M. Craven, and W. C. Harris, Vibrational Spectra and Structure, Vol. 1, J. R. Durig, ed., Marcel Dekker, New York, N.Y., (1972) 73-179.
2. A. V. Cunliffe, Internal Rotation in Molecules, W. J. Orville-Thomas, ed., John Wiley and Sons, New York, N.Y., (1974) 217-252.
3. W. G. Fateley, Pure App. Chem., 36, (1973) 119.
4. J. R. Durig and A. W. Cox, Jr., J. Chem. Phys., 63, (1975) 2303.
5. J. R. Durig and A. W. Cox, Jr., J. Phys. Chem., 80, (1976) 000.
6. J. R. Durig, S. M. Craven and J. Bragin, J. Chem. Phys., 51, (1969) 5663.
7. J. R. Durig, S. M. Craven, J. H. Mulligan and C. W. Hawley, J. Chem. Phys., 58, (1973) 1281.
8. J. G. Pritchard and H. M. Nelson, J. Phys. Chem., 64, (1960) 795.
9. J. C. Evans and G. Y.-S. Lo, J. Amer. Chem. Soc., 88, (1966) 2118.
10. J. R. Durig, J. W. Thompson, V. W. Thyagesan and J. D. Witt, J. Mol. Struct., 24, (1975) 41.
11. T. Kojima, E. L. Breig and C. C. Lin, J. Chem. Phys., 35, (1961) 2139.
12. D. R. Lide, Jr., and D. E. Mann, J. Chem. Phys., 28, (1958) 572.
13. J. D. Lewis, T. B. Malloy, Jr., T. H. Chao and J. Leane, J. Mol. Struct., 12, (1972) 427.
14. M. D. Harmony, Chem. Soc. Rev., Vol. 1, The Chemical Society, London, (1972) 211.
15. M. Tsuboi, A. Y. Hirakawa, K. Tamagake, Nippon Kagaku Zasshi, 89 (a), (1968) 821.
16. D. Kivelson and D. R. Lide, Jr., J. Chem. Phys., 27, (1957) 353.

ORIGINAL PAGE IS  
OF POOR QUALITY

Table I Vibrational Frequencies ( $\text{cm}^{-1}$ ) and their Assignments for t-Butylphosphine.<sup>a</sup>

Infrared gas	Rel. Int.	Infrared solid	Rel. Int.	Raman gas	Rel. Int.	Raman Liquid	Rel. Int.	Raman Solid	Rel. Int.	Assignment
2967	vs							2959	s,sh	} $\nu_1, \nu_2, \nu_3, \nu_4, \nu_5$ $\nu\text{C-H}$ sym. } $\nu_{24}, \nu_{25}, \nu_{26}, \nu_{27}$ $\nu\text{C-H}$ asym.
2960	vs									
2954	vs									
		2949	vs	2961	s p	2953	s p	2949	vvs	
2946	vs,sh			2947	s,sh p	2936	s,sh p	2943	s,sh	
		2937	vs							
				2934	vs p	2925	vs p	2930	s,sh	
		2924	s					2910	m,sn	
		2892	s					2894	vvs	
2906	s			2907	vvs p					
		2877	s					2878	s,sh	
		2857	vs					2861	s	
2873	s			2875	vs p	2865	vs p			
								2850	s,sh	
				2786	m p	2767	m p	2769	m	
				2724	s p	2715	m p	2707	m	$2\nu_{10}?$
		2279	vvs					2293	vvs	$\nu_{28}$
2292	vvs	2293	vvs	2292	vvs p	2284		2278	vvs	$\nu_6$

ORIGINAL PAGE IS  
OF POOR QUALITY

Table I (continued)

Infrared gas	Rel. Int.	Infrared solid	Rel. Int.	Raman gas	Rel. Int.	Raman Liquid	Rel. Int.	Raman Solid	Rel. Int.	Assignment	
								1473	w	$\nu_{29}$	
1476	s,sh	1467	m,sh					1463	m	$\nu_7$	
		1462	vs					1462	m,sh	$\nu_{30}$	
				1450	w,brd dp?	1458	w,brd dp?			} $\delta\text{CH}_3$ asym.	
1466	s	1457	s,sh					1457	s		$\nu_8$
1456	s,sh	1442	m,sh			1445	w,brd	1442	s		$\nu_9$
		1436	w,sh					1438	m,sh		$\nu_{31}$
1370	s	1368	m,sh							$\nu_{10}$	
		1363	s,sh					1363	vw	$\nu_{11}$	
1361	s,sh									} $\delta\text{CH}_3$ sym.	
		1361	s					1361	vw,sh		$\nu_{32}$
1204	m	1211	w,sh	1205	w,sh			1211	s	$\nu_{33}$	
1198	m	1192	m	1199	m p	1192	m,brd p	1192	s	$\nu_{12}$	
1192	m	1169	w	1192	m p	1173	w,sh p	1170	w,sh	$\nu_{13}$ $\rho\text{CH}_3$	
1183	m			1184	w,sh					} "hot" bands of $\nu_{12}$	
1172	w			1173	w,sh						
1090	m	1096	s			1089	vw,sh p	1092	vw	$\nu_{14}$ $\rho\text{CH}_3$	
1074	m	1074	m	1080	vw,brd p	1072	vw,brd p	1075	m	$\nu_{15}$ $\rho\text{PH}_2$ sic	
		1065	w,sh					1065	w,sh	$\nu_{34}$	
		1036	w,sh					1036	w,sh	$\nu_{35}$	
				1029	vw p	1029	vw p			} $\rho\text{CH}_3$	
1029	m	1029	m					1032	w		$\nu_{16}$
940	w	938	w	940	vw dp	938	vw,brd dp	937	m	$\nu_{3E}$	

Table I (continued)

Infrared gas	Rel. Int.	Infrared gas	Rel. Int.	Raman gas	Rel. Int.	Raman Liquid	Rel. Int.	Raman Solid	Rel. Int.	Assignment
834	m	834	w	835	m p	833	w p	836	vs	$\nu_{17}$ $\nu_{C-C}$ sym.
826	s	816	m	825	w dp	820	vw,sh dp?	818	m	$\nu_{37}$ $\delta PH_2$ twist
799	s	793	vs	800	s p	798	m p	795	m	$\nu_{18}$ $\delta PH_2$ wag
589 <sup>b</sup>	m	592	w	590	vs p	589	vs p	593	vs	$\nu_{19}$ $\nu_{C-P}$
394	w							394	vw,sh	$\nu_{36}$ $\delta CCC$ asym.
		383 <sup>b</sup>	vw	382	vw,sh	386	vw,sh			
389	w							389	m	$\nu_{20}$ $\delta CCC$ asym.
352	s	354	s	354	s p	354	m p	353	s	$\nu_{21}$ $\delta CCC$ sym.
346	m			347	w,sh p					"hot" band of $\nu_{21}$
								297	w,sh	$\nu_{39}$ $\delta CCP$ sym.
285	w	295	vw			287	vw,brd dp?			
								295	w	$\nu_{22}$ $\delta CCP$ asym.
		278	s,sh	280	w,brd					$\nu_{23}$ $\tau CH_3$
266	w					272	vw,sh p?	276	w	
		273	s							$\nu_{40}$ $\tau CH_3$
227	w	248	w,brd			239	vvw,brd dp	248	vw	$\nu_{41}$ $\tau CH_3$
181	s	200	vs	172	vvw,brd dp	182	vvw,brd dp	199	vw	$\nu_{42}$ $\tau PH_2$
171	s									"hot"band of $\nu_{42}$
								77		} Lattice modes
								63		
								50		
								42		
								32		

<sup>a</sup>Abbreviations used: s, strong; m, medium; w, weak; v, very; brd, broad; sh, shoulder; p, polarized, dp, depolarized.

<sup>b</sup>This and lower frequencies obtained from the far infrared spectra.

ORIGINAL PAGE IS  
OF POOR QUALITY



TABLE II METHYL TORSIONAL FREQUENCIES AND POTENTIAL  
CONSTANTS FOR T-BUTYLPHOSPHINE

	Solid	Gas
$\nu_E^a$ (cm <sup>-1</sup> )	275.5	266
$\nu_A$ (cm <sup>-1</sup> )	248	227
F (cm <sup>-1</sup> )	5.3559	5.3559
F' (cm <sup>-1</sup> )	0.0278	0.0278
2/9K (kcal/mole)	4.22	3.81
2/9L (kcal/mole)	-0.15	-0.20

a)  $\frac{1}{2}(A' + A'')$

TABLE III PHOSPHINO TORSIONAL FREQUENCIES  
( $\text{cm}^{-1}$ ) AND POTENTIAL CONSTANTS

Transition	frequency		
	obs.	calc.	obs.-calc.
1 $\leftarrow$ 0	181	180.9	0.1
2 $\leftarrow$ 1	172	172.2	-0.2
3 $\leftarrow$ 2	162	161.8	0.2
4 $\leftarrow$ 3 <sup>-</sup>		146.1	4.9
4 $\leftarrow$ 3 <sup>+</sup>	(151) <sup>2</sup>	149.1	0.9
V <sub>3</sub>	976 $\pm$ 4 <sup>b</sup> $\text{cm}^{-1}$ or 2.79 $\pm$ 0.01 kcal/mole		
V <sub>6</sub>	-25 $\pm$ 2 $\text{cm}^{-1}$ or 0.07 $\pm$ 0.01 kcal/mole		

a) estimated

b) quoted error limits represent one standard deviation

## FIGURE CAPTIONS

- Figure 1 Raman spectra of t-butylphosphine
- A) gaseous
  - B) liquid
  - C) solid
- Figure 2 Mid-infrared spectra of t-butylphosphine
- A) gaseous, approximately 2 torr sample pressure
  - B) gaseous, approximately 40 torr sample pressure
  - C) solid
  - D) solid, more sample deposited
- Figure 3 Far infrared spectra of t-butylphosphine
- A) gaseous, approximately 400 torr
  - B) solid<sup>a</sup>
- a) pronounced baseline "roll" is due to interference fringes from the sample film

