

General Disclaimer

One or more of the Following Statements may affect this Document

- This document has been reproduced from the best copy furnished by the organizational source. It is being released in the interest of making available as much information as possible.
- This document may contain data, which exceeds the sheet parameters. It was furnished in this condition by the organizational source and is the best copy available.
- This document may contain tone-on-tone or color graphs, charts and/or pictures, which have been reproduced in black and white.
- This document is paginated as submitted by the original source.
- Portions of this document are not fully legible due to the historical nature of some of the material. However, it is the best reproduction available from the original submission.

04

E78-10373 II
CR-147966
6/16/76

Type II Progress Report

Evaluation of Landsat-2 Data
for
Selected Hydrologic Applications

"Made available under NASA sponsorship
in the interest of early and wide dis-
semination of Earth Resources Survey
Program information and without liability
for any use made thereof."

Donald R. Wiesnet, Principal Investigator
David F. McGinnis, Jr., Coinvestigator
Michael C. McMillan, Associate
Michael Matson, Associate

NOAA/NESS/ESG
WWB, Room 810
Washington, D.C. 20233

(E78-10373) EVALUATION OF LANDSAT-2 DATA
FOR SELECTED HYDROLOGIC APPLICATIONS
PROGRESS REPORT (NATIONAL OCEANIC AND
ATMOSPHERIC ADMINISTRATION) 3 P HC \$3.50

N76-25618

UNCLAS
CSCL 08H G3/43 00373

Landsat Investigation No. 23170


Contract No. NAS⁵-35991A

23170

RECEIVED

JUN 21 1976

SIS/902.6



A. Problems

None

B. Accomplishments

1. Complete basic data set on March 1975 Phoenix test site soil survey received from Texas A&M.

2. Landsat MSS 4 and MSS 6 digital data were compared with averaged soil moisture values for bare and vegetated fields from the Phoenix test site. Regression analyses produced a correlation coefficient of $r^2 = 0.71$ ($r=.84$) for a power fit of satellite MSS 4 radiance values vs. averaged soil moisture for bare fields, and an $r^2 = .69$ ($r=.83$) for a similar fit of MSS 6 values. For vegetated fields, MSS 4 values remained almost constant, not affected by vegetation type or soil moisture. Using MSS 6 data, alfalfa fields increased in brightness as soil moisture increased, providing an $r^2 = .88$ ($r=.94$) for a parabolic fit. Wheat field MSS 6 brightness varied little with moisture changes.

3. Aircraft M²S tapes received from NASA for October 1975 flight over Luverne.

4. Snow mapping of American River basin for February 20, 1976 and March 9, 1976 using Landsat imagery completed. Results and comparisons are listed below:

	<u>% Snow Cover</u>		
<u>Date</u>	<u>MSS 5</u>	<u>MSS 7</u>	<u>SMS-2</u>
Feb. 20, 1976	54%	51%	55%
Mar. 9, 1976	46%	41%	46%

C. Significant Results

None

D. Publications

Received from Texas A&M

Blanchard, Bruce J., 1975, Ground Truth Report 1975 Phoenix Microwave Experiment, Remote Sensing Center, College Station, Texas

Received from EG&G

Feimster, E.L. and Fritzsche, A.E., 1975, Soil Moisture Survey
Experiment at Luverne, Minnesota, EG&G, Las Vegas, Nevada

E. Recommendations

None

F. Funds Expended

\$7.6K remains of funds available for investigation.

G. Data Use

CCT tapes used in obtaining MSS 4 and MSS 7 radiance values for Phoenix test site in March, 1975. 70 mm positive transparencies used in mapping American River Basin snow cover for February 20, 1976 and March 9, 1976.

REPRODUCIBILITY OF THE
ORIGINAL PAGE IS POOR