

General Disclaimer

One or more of the Following Statements may affect this Document

- This document has been reproduced from the best copy furnished by the organizational source. It is being released in the interest of making available as much information as possible.
- This document may contain data, which exceeds the sheet parameters. It was furnished in this condition by the organizational source and is the best copy available.
- This document may contain tone-on-tone or color graphs, charts and/or pictures, which have been reproduced in black and white.
- This document is paginated as submitted by the original source.
- Portions of this document are not fully legible due to the historical nature of some of the material. However, it is the best reproduction available from the original submission.

"Made available under NASA sponsorship
in the interest of early and wide dis-
semination of Earth Resources Survey
Program information and without liability
for any use made thereof."

N76-21648

Unclas
00281

G3/43

(E76-10281) THE FRENCH ATLANTIC LITTORAL
AND THE MASSIF ARMORICAIN Progress Report,
Jan. - Mar. 1976 (Ecole Pratique des Hautes
Etudes) 28 p HC \$4.00
CSCL 08E

I.D. No 29690

E76-10281

CR-146651

THE FRENCH ATLANTIC LITTORAL

AND THE MASSIF ARMORICAIN

Prof. Fernand VERGER

Dr. Jean-Marie MONGET

Dr. Jean-Yves SCANVIC

ECOLE PRATIQUE DES HAUTES ETUDES 3

61, rue Buffon

75005 PARIS-FRANCE

RECEIVED
NASA STI FACILITY
ACQ. BR.

Excl
APR 19 1976

March 1976

SCAF# 1002583
1 2 3 4 5

Type I Report for period

January - March 1976

CENTRE NATIONAL D'ETUDES SPATIALES

129, rue de l'Université

75007 PARIS-FRANCE

Original photography may be purchased from:
EROS Data Center
10th and Dakota Avenue
Sioux Falls, SD 57198

ORIGINAL CONTAINS
COLOR ILLUSTRATIONS

1. D. Number 29.690	2. Type of report 1	3. Recipient's catalog No.
4. Title THE FRENCH ATLANTIC LITTORAL AND THE MASSIF ARMORICAIN		5. Report Date : March 1976
		6. Period Covered : January - March 1976
7. Principal investigator Prof. Fernand <u>VERGER</u> <i>ent</i>		8. No of pages : 25
9. Name and Address of Principal investigator's Organization ECOLE PRATIQUE DES HAUTES ETUDES 61, rue Buffon 75005 - PARIS. France		10. Principal Investigat. Rept. No 2
		11. GSFC Technical Monitor CRUMP Edward W.
12. Sponsoring Agency Name and Address CENTRE NATIONAL D'ETUDES SPATIALES 129, rue de l'Université 75007 - PARIS. France		13. Key Words (Selected by Principal Investigator) Tidal marsh; littoral transport, estuary ; beaches ; wetlands ; computer analysis
14. Supplementary Notes		Part 2 : faulting ; ore deposits.

Prepared in cooperation with J. M. MONGET and J.-Y. SCANVIC

15. Abstract :

This second progress report gives initial results on Landsat - 2 data received in the corresponding period. During this period 22 scenes were received and evaluated. Most of them covered the summer of 1975 and cloud cover was on the whole quite low.

Two more scenes have been ordered in CCT Format :

- Loire Estuary
- Islands of the bay of Biscay.

These have been chosen to complete or extend time-series covering significant test areas.

Part 2 of the report covers research done by the B.R.G.M.

1 - INTRODUCTION

This report covers the first quarter of 1976 during which a number of significant new techniques has been made operational using, LANDSAT-1 data.

Over the same period, LANDSAT-2 data has been received and reformatted to our databank standard (assembly of 4 strips to single scene recording) on direct-access units.

Interpretation, classification and mapping of LANDSAT-2 data has been initiated, singly or in connection with LANDSAT-A data, with emphasis on coastal areas, extending seaward to the extent of coastal sediment transport and landward into saltmarshes, dunes and polders.

2 - TECHNIQUES

2-1 - Photographic : 70 mm B/W negatives were examined as received, and after enlargement to 1/1000 000 th scale. Particular attention was directed to scenes suitable for ordering in CCT format.

2-2-1 : 4 strip records were copied on disk using 3240 pixel/line format to facilitate data processing, especially in areas bisected by strip borders.

2-2-2 : Various improvements were carried out on our supervised automatic mapping procedures (using IBM 360 and Benson Offline Plotting System) to save program execution time.

2-2-3 : Time sharing (Tektronix 4013 output linked to IBM 360) : further development of the FRACAM package have now made this unsupervised interactive classification system fully operational.

2-2-4 - The CLAMS unsupervised automatic classification program provides graphic output on Calcomp microfilm.

2-2-5 : A test run for evaluation of the stand-alone GE-IMAGE-100 supervised system was conducted by Dr. J.M. MONGET at the NASA-JSC remote sensing facility. Sample color displays one shown in Fig. 2.2.5.1 and 2.2.5.2. Special attention was given to the software/hardware trade-offs.

2-2-6 : A smoothing technique is being tested based on using each pixel in the mapped output from programs (2.2.2. and 2.2.3) as the centre of a rectangular or square matrix and replacing this pixel with the modal signature found in the surrounding matrix.

3 - ACCOMPLISHMENTS

3-1 - Ground truth : for LANDSAT-2 data, we have continued to draft maps of coastal areas showing directions of currents and tide heights (following pages). Times given are T.U. (Fig. 3.1.1. to 3.1.6).

3-2 - Mapping of offshore currents (Fig. 3-2)

The map of the Island of Jersey and surrounding marine approaches was obtained from LANDSAT-1 imagery by automatic mapping techniques described previously. Green areas are land, black and brown show clouds and cloud

shadow (a major interpretation hindrance). Blue and yellow show waters of increasing turbidity (symbols 1 through 4) and in particular the strange shapes found in the mixing area of the turbidity front. No explanation has been found for this phenomenon, actually under investigation, but it is a typical example of scientific observation which could not be obtained by any other means, and for which the scale if not the spectra of MSS imagery is ideally suited.

3-3 - Application of techniques developed with LANDSAT-1 data to the fromentine test-site, chosen for its extensive homogenous landscape.

Fig. 3.3.1 is an example of graphic output after supervised classification water was separated using MSS band 7 and then differentiated by turbidity detected in band 5 (symbols 1-6).

Shoreline sediment has been classified according to type : sand (symbol 8) and mudflats (symbols 9 through 12).

For the mudflats, varying degrees of wetness (decreasing from 9 to 12) have been identified. In the north-eastern corner, freshly ploughed polders show up as sandy areas. Finally symbols 13 to 18 show various types of vegetation, particularly 13 and 14 (psammophytes more or less covering dunes), maritime pine forests (15), and cultivated marshes (16).

The following illustrations are an example of data smoothing described in 2.2.6. From the original 1/80 000 scale classification, which used data from two LANDSAT-1 scenes, the FRALISSE program devised Fig. 3.3.3. (smoothing by retraining modal values in a 3 x 3 matrix round each pixel) and, further Fig. 3.3.4 where a 5 x 5 matrix was used.

Here again, darkening shades of blue show water of decreasing turbidity (1 through 5) and brown and yellow are mud and sand. For these last two types, data from 2 scenes have been used as an 8 band classifying instrument. The lighter symbols (5,7) show areas above tide levels at both dates and the darker ones (6,8) are areas uncovered on Sept 27, 1972 but covered on March 8, 1973.

Finally Fig. 3.3.5 is an example of 3 scene diachronic classification, using 12 MSS channels and whereas water was drawn from a single scene, it is now possible to identify 3 tide levels in sandy and mudflat areas and to draw more accurate distinctions between 3 types of vegetation.

All these maps have been drawn up over a fairly long period due to data delivery, research and publication delays. But we are presently applying similar techniques to LANDSAT-2 data, particularly a scene covering the Fromentine test-site and preliminary results show that a 3 year span is sufficient to allow efficient monitoring of coastline dynamics.




Fig. 2.2.5.1





Fig. 2.2.5.2

GE IMAGE 100 colour output for the Fromentine test site.





Tidal currents (1 cm = 1 m/s)

Tidal range and water level (2 mm = 1 m)

June 22, 1975



Fig. 3.1.1. : Hydrological ground - truth data.



Tidal currents (1 cm = 1 m/s)

Tidal range and water level (2 mm = 1 m)

July 28, 1975

→ S H O M

Maximum high water springs





Fig. 3.1.2. : Hydrological ground - truth data.

REPRODUCIBILITY OF THE ORIGINAL PAGE IS POOR



Tidal currents (1 cm = 1 m/s)


Tidal range and water level (2 mm = 1 m)

July 28, 1975

Maximum high water springs

→ S H O M

falling tide
 rising tide




High water

Tide level 10^h13 TU

Low water

Chart datum 0

Fig. 3.1.3. : Hydrological ground - truth data.



Tidal currents (1 cm = 1 m/s)

Tidal range and water level (2 mm = 1 m)

July 29, 1975

→ S H O M

Maximum high water springs

▼ falling tide
▲ rising tide




High water
Tide level 10h18T
Low water

Chart datum 0

Fig. 3.1.4. : Hydrological ground - truth data.

REPRODUCIBILITY OF THE ORIGINAL PAGE IS POOR



Tidal currents (1 cm = 1 m/s)

Tidal range and water level (2 mm = 1 m)

July 29, 1975

→ S H O M

Maximum high water springs



↘ falling tide
↙ rising tide



High water
Tide level 10^h18 TU
Low water

Chart datum 0

Fig. 3.1.5. : Hydrological ground - truth data.



Tidal currents (1 cm = 1 m/s)

Tidal range and water level (1 mm = 1 m)

July 30, 1975

Maximum high water springs

→ S H O M

↘ falling tide
↗ rising tide



High water
Tide level 10^h23 TU
Low water

Chart datum 0

Fig. 3.1.6. : Hydrological ground - truth data.

FORMES DE TURBIDITE DANS LE GOLFE NORMAND BRETON

DONNEES ERTS-1 DU 8 MARS 1973

REPRODUCIBILITY OF THE ORIGINAL PAGE IS POOR



Fig. 3.2. : Sediment transport round the Island of Jersey.

Revue "PHOTO-INTERPRETATION" © 6-1974 (4^e tr.) Editions TECHNIP



REPRODUCTION OF THE
COPYRIGHT PAGE IS POOR

donnees NASA - carte automatique FRALIT


NOMBRE DE CARACTERES	SURFACE EN KM ²	NUMERO DE REFERENCE	NOMBRE DE CARACTERES	SURFACE EN KM ²	NUMERO DE REFERENCE	NOMBRE DE CARACTERES	SURFACE EN KM ²	NUMERO DE REFERENCE			
	1958	8.83	1		225	1.01	7		104	0.47	13
	3991	17.99	2		480	2.16	8		1117	5.04	14
	10245	46.18	3		976	4.40	9		1838	8.29	15
	2945	13.28	4		1697	7.65	10		1683	7.59	16
	2940	13.25	5		503	2.27	11		7597	34.25	17
	1366	6.16	6		77	0.35	12		1364	6.15	18

P
5 3 2 0 6 1 0 0 0

1 : 50 000 IGNF	1 : 75 000	A 0,6-0,7 μm	A 0,8-1,1 μm	27-9-72	10 h 29	NASA-LANDSAT 1 MSS 5 et 7	Image n° 1066-10294-5 et 7	F FRANCE
feuilles NORMOUTIER 24 X	feuilles CHALLANS 25 X et XI							

Fig. 3.3.1. : The Fromentine test site : scale = 1/75 000.

CARTE DIACHRONIQUE DU GOULET DE FROMENTINE D'APRES LES DONNEES DE LANDSAT-1
ATLAS INFOGRAPHIQUE DU LITTORAL FRANCAIS (PROGRAMME FRALIT) CONVENTION 75/CNES/222




ÉCHELLE : 1 : 80000

Fig. 3.3.2. : The Fromentine test site : scale = 1/80 000.

REPRODUCIBILITY OF THE ORIGINAL PAGE IS POOR


CARTE DIACHRONIQUE LISSEE DU GOULET DE FROMENTINE D'APRES LES DONNEES DE LANDSAT-1
ATLAS INFOGRAPHIQUE D' LITTORAL FRANCAIS (PROGRAMME FRALIT) CONVENTION 75/CNES/222



ÉCHELLE : 1 : 80 000

Fig. 3.3.3. : The Fromentine test site : scale = 1/80 000, 3 x 3 smoothing.

CARTE DIACHRONIQUE LISSEE DU GOULET DE FROMENTINE D'APRES LES DONNEES DE LANDSAT-1
ATLAS INFOGRAPHIQUE DU LITTORAL FRANCAIS (PROGRAMME FRALIT) CONVENTION 75/CNES/222



ÉCHELLE : 1 : 80 000

Fig. 3.3.4. : The Fromentine test site : scale = 1/80 000, 5 x 5 smoothing.

4 - SIGNIFICANT RESULTS

Diachronic use of LANDSAT data time series will in time allow us to study statistically submersion frequencies in tidal areas. This is an essential element of coastal geomorphology and of coastal zone management bening particularly useful in siting shellfish farms. The maps we are obtaining, at useable scales and simple, user oriented legends should become an essential document for coastal planning agencies.

5 : LIST OF PUBLICATIONS :

- Fernand VERGER : Une cartographie automatique des données de LANDSAT - 1.
Photo-Interprétation. XIII^e année, 1974, fasc. 6, p. 1-6.
- Fernand VERGER , Gérard JOLY : Cartographie diachronique à partir des données numériques de LANDSAT - 1.
Photo-Interprétation. XIII^e année, 1974, fasc. 6, p. 7-13.
- Odile GUERIN , Jean-Marie MONGET : Intérêt de la télédétection pour la connaissance du milieu estuarien ; l'exemple de l'estuaire de la Loire.
Bull. Union Océanogr. de France, 1975, vol. 7, n° 12, p. 47-54,
2 fig.

6 : DATA QUALITY AND DELIVERY.

Since the last report, 22 frames of B/W 70 mm imagery have been received. Despite certain delays we find that data has been reading us more regularly (2 to 3 months after coverage). Despite extensive inland cloud coverage negative quality has been good in coastal areas but too contrasted for photographic interpretation in marine zones.

The five CCTS ordered during the last quarter of 1975 have been received and treatment has begun using techniques described above and previously in connection with LANDSAT - 1 imagery. Delay was reasonable (2 months from order to reception) but one tape (scene no E 2133 - 10125, quality given as G, F, F, G) appears to be unusable since repeated attempts at reading it cause numerous parity errors.

DATE	PRODUCT ID	BLACK AND WHITE PRODUCTS				CLOUD COVER (%)	TAPE PRO-DUCT 0=order received 1=received	ORBIT NUM-BER	CODE OF FRAMES (L = Littoral)
		MSS (Quality)							
		4	5	6	7				
10 April 75	E2 078-10055	F	P	F	F	30		1083	FLANDRES (L)
"	E2 078-10062	F	F	F	F	40		"	PICARDIE (L)
"	E2 078-10064	F	G	G	G	70		"	BEAUCE
"	E2 078-10071	G	G	G	G	60		"	ANJOU
"	E2 078-10073	G	G	G	G	60		"	CHARENTE-GIRONDE
"	E2 078-10080	G	G	G	F	50		"	LANDES (L)
"	E2 078-10082	G	G	F	G	60		"	GOLFE GASCOGNE (L)
3 June 75	E2 132-10070	P	P	P	P	80		1836	ANJOU-VENDEE
"	E2 132-10073	G	G	G	G	60		"	GIRONDE
"	E2 132-10075	F	G	G	F	70		"	LANDES (L)
"	E2 132-10082	F	G	G	G	60		"	PYRENEES
4 June 75	E2 133-10113	G	P	F	G	40		1850	ARTOIS (L)
"	E2 133-10120	F	F	F	G	30		"	BAIE DE SEINE (L)
"	E2 133-10122	G	P	G	G	60		"	BAIE DE SEINE (L)
"	E2 133-10125	G	F	G	G	50		"	VENDEE (L)
"	E2 133-10131	G	F	F	G	50	⊕	"	GIRONDE (L)
"	E2 133-10134	F	P	G	G	40	⊕	"	LANDES (L)
6 June 75	E2 135-10233	F	P	P	F	20		1878	JERSEY-COTENTIN (L)
"	E2 135-10235	F	F	G	F	30	⊕	"	BAIE ST MICHEL (L)
"	E2 135-10242	F	F	G	F	60		"	MORBIHAN (L)
8 June 75	E2 137-10352	F	G	G	G	50		1906	OUESSANT (L)
21 June 75	E2 150-10060	F	F	F	F	40		2087	FLANDRES (L)
"	E2 150-10071	F	F	G	G	50		"	TOURAIN
"	E2 150-10074	G	G	G	G	(a) 30		"	GIRONDE (L)
"	E2 150-10080	F	F	G	G	10		"	ARCACHON (L)
"	E2 150-10083	G	G	G	G	10		"	GOLFE DE GASCOGNE (L)
22 June 75	E2 151-10135	F	F	F	G	20		2101	ARCACHON (L)
26 June 75	E2 155-10353	F	F	F	F	40		2157	MER CELTE (L)
9 July 75	E2 168-10073	G	G	G	G	50		2338	SAINTONGE
"	E2 168-10080	G	G	G	G	50		"	ARCACHON (L)
"	E2 168-10082	F	G	G	G	40		"	GOLFE DE GASCOGNE(L)
14 July 75	E2 173-10352	G	F	G	G	30		2408	OUESSANT (L)
27 July 75	E2 186-10071	G	G	G	G	0		2589	CHARENTE
"	E2 186-10073	G	F	G	G	10		"	ARCACHON (L)
"	E2 186-10080	G	G	G	G	40		"	COTE D'ARGENT (L)
28 July 75	E2 187-10111	F	P	G	X	30		2603	PAS-DE-CALAIS (L)
"	E2 187-10114	F	F	G	G	30		"	PICARDIE (L)
"	E2 187-10120	G	F	G	G	50		"	BAIE DE SEINE (L)
"	E2 187-10123	G	F	G	G	20		"	ANJOU
"	E2 187-10125	G	G	G	G	20		"	CHARENTE (L)
"	E2 187-10132	G	G	F	G	10		"	ARCACHON (L)
"	E2 187-10134	F	G	G	F	10		"	GOLFE DE GASCOGNE (L)
29 July 75	E2 188-10172	F	F	F	F	10		2617	MANCHE/BAIE DE SEINE (L)
"	E2 188-10181	F	F	F	G	10	o	"	ESTUAIRE LOIRE (L)
"	E2 188-10183	F	G	F	F	0	o	"	VENDEE-CHARENTE (L)
"	E2 188-10190	F	F	F	G	0		"	LARGE DES LANDES
30 July 75	E2 189-10230	F	F	F	F	10		2631	N COTENTIN (L)
"	E2 189-10233	F	F	F	G	10		"	GOLFE ST MALO (L)
"	E2 189-10235	F	F	F	G	10		"	MORBIHAN (L)
"	E2 189-10242	F	F	F	F	10		"	GOLFE DE GASCOGNE
4 Sept. 75	E2 225-10222	G	G	F	G	50		3133	ANGLETERRE/COTENTIN (L)
10 Sept. 75	E2 241-10120	F	G	G	G	50		3356	CHARENTE/GIRONDE (L)
"	E2 241-10123	F	F	G	G	40		"	ARCACHON/LANDES (L)
"	E2 241-10125	F	F	F	F	30		"	ST JEAN DE LUZ (L)

REPRODUCIBILITY OF THE
 ORIGINAL PAGE IS
 POOR

PART 2 : FIRST PRELIMINARY REPORT ON LANDSAT 2

SPATIAL IMAGES INTERPRETATION : L'ARMORIQUE

I : TECHNIQUES

A - Photographic processing

We have generated black and white imagery products from negatives sent by Professor Verger, principal investigator. The chosen scale is 1 million.

B - Work procedure

B. 1. - Interpretation of the different available spectral bands (on overlay), generally bands five and/or seven was carried out under a mirror stereoscope (Wild ST4), not to have a stereoscopic effect, but because it allows a binocular study ; we feel it is a good way to do the interpretation.

First of all we do the interpretation in using two similar spectral bands, for instance band five. Then we compare, always under a stereoscope, the results obtained on different spectral bands.

As far as possible, we also compare and complete the interpretation done on adjoining images, using the stereoscopic effect.

B. 2. - Identification of the Landsat image is obtained by reference to the 1 million scale topographical and geological maps of France, Unfortunately on the later edition of the geological map coordinates are in grades out but this is a minor problem since images can generally be located at a glance.


Then, to obtain an accurate location of observed data we trace on the overlays all visible hydrographic detail.

B. 3. - "Ground truth" techniques utilized mainly consist, in this first stage, in a search of available scientific papers concerning proposed subjects and new data found in the area of interest. This point concerns, geology, geophysics, drilling, etc... as points of interest are raised interpretation and correlation, field geologists mapping the area at a 1/50 000 scale are advised.

II : ACCOMPLISHMENTS

The actual french atlantic coast is often a cloudy area and good quality images are not numerous. Nevertheless we have been able, using both Landsat 1 and 2 images, to investigate some of the problems we proposed to N.A.S.A. for investigation with space data.

SOUTH BRITANNY



ID N° 1066-10294

Scale 1/1,000,000

Sept. 27.72 Mss Band 5

REPRODUCIBILITY OF THE ORIGINAL PAGE IS POOR

1 - Structural analysis

a - Boundaries between folded and not folded Cadomian area in central Armorique : this point is not yet investigated with the received documents because of the lack of complete coverage of the area.

b - Mapping of fractures is in progress in the process of being. The main lineament known in southern Armorique are visible well observed on ERTS 1 imagery. New directions of fracturing are also visible. (see fig.)

c - Some of the granitized axes known through ground mapping in the northern Armorique show very typical tone and pattern on Landsat 2 images and locally allow a differentiation between younger and ancient granites.

d - Several small sedimentary basins (Tertiary in age) exist in the " Massif Armoricaïn ". A first attempt to map them, between the towns of Rennes and Nantes, on Landsat images, was unsuccessful.

e - Extent of fracturing - The gravimetric map of France (1 million scale), recently published by B.R.G.M., geophysical department, reveals several extensions of the southern armoricaïn lineament toward the south east, to the Massif Central. South of Nantes, gravimetric data cannot indicate which of the three observed geophysical discontinuities corresponds to the above mentioned lineament. A careful interpretation of four ERTS 1 and 2 images gives a preliminary answer to the question.

2 - Mineral exploration

a - Pb - Zn - Ba deposits.

Pb, Zn, Ba deposits are found, outside Massif Armoricaïn and Massif Central, in the sedimentary levels of the Poitou area. Three of them are important : Melle Alloue-Ambazac and Nontron-Ehebrinas and this last one is located along a fault trending NW-SE, the main Armoricaïn direction. Our purpose was to look for similar faults in Melle and Alloue-Ambazac, and to find similar directions which could direct new prospection. First results outline such faulting in Melle and Alloue-Ambazac, according to ERTS - 2 image interpretation. This faulting is parallel to the main armoricaïn trends.

b - Sn deposits are located in or near Cadomian and Hercynian granite, along first order lineaments. Landsat 1 image show, near Nozay, one of the main deposits, why ore occurrences are disappear eastward : an unknown north-south fault cuts the Sn bearing formation, disturbing the continuity of the deposit.

The problem of mineralized Cadomian granite - Hercynian sterile granite differentiation is on the way (see above).

c - Kaolin deposits.

Kaolin deposits are known everywhere in the Armorique. We plan to test the ability of colour composite imagery to enhance . Cromalin process is to be used. The work is in progress in the northern part of Armorique.

3 - Ground water

We have proposed to survey water supply possibilities in two types of structure.

- The sandstone layers : their water bearing ability partly depends on their. The first results obtained in this domain confirm Landsat image mapping possibilities of fractures in this formation.

- Tertiary basins, where thick, could have important water bearing possibilities. We have seen they are not differenciated by tone on the ERTS images but, if they are linked with north-south fractures, as mentioned by Jaeger, the similar directions outlined by space data are promising to direct future water supply research.

4 - Other results

Circular structures detected in the vicinity of Limoges on Landsat 1 image : (Structural investigation in the Massif Central using ERTS - 1 multispectral imagery - N.A.S.A. report 1974) are also visible on ERTS - 2 images (2186 - 10071 - July 1975). We would like to point out specially the Rochechouart meteorite, this new image confirming its possible southern extend and its general shape, as visible on ERTS - 1 image and mapped on the field.

III : SIGNIFICANT RESULTS

The main results concern :

- The extent of the armorican southern lineament toward the south east until Montalembert where north 20 faulting disrupts it: On a global tectonic point of view this is a significant result.

- The existence of armorican fractures directions near the Melle and Alloue-Ambazac Pb, Zn, Ba deposits, new data confirming genesis hypothesis.

- The discovery of similar other possible faults which can be a guide for further Pb-Zn prospection in the Poitou area.

IV : PUBLICATIONS

No publications are in progress on these first results but since the B.R.G.M. prepared the final N.A.S.A. report (ERTS - 1) on the Massif Central some papers were published, using ERTS - 1 image interpretation results. They concern :

M. LAMBERT - La structure impactitique de Rochechouart (Limousin) et du contexte regional par interpretation des photos-satellites ERTS Bull. B.R.G.M., section 1, n° 4, 1974.

J.-M. BROSSE - Apport de la télédetection en géologie structurale (Massif Central - France) Thèse Orléans et Bull. B.R.G.M., 2è série, Section II, n° 6, 1975.

N. DEBEGLIA, A. GERARD - Apport de l'aéromagnétisme à l'étude du Massif Central français.
Bull. B.R.G.M., section II, n° 4, 1975.

J.-Y. SCANVIC - Apport de l'imagerie spatiale multi-spectrale à la compréhension tectonique du Massif Central français et de son environnement sédimentaire.
Bull. B.R.G.M., section II, n° 4, 1975.

V : DATA QUALITY

- Bande 7 is often unusable along the sea side (very pale), by contrast.

- On some scenes, for instance 2186 - 10071 - 27 July 1975, covering both sedimentary and basement formations, while the sedimentary area is a very good quality image the basement area is generally too dark to be well interpreted.

VI : CONCLUSIONS

Having used Landsat 1 imagery to investigate tectonic features we can, assume it is a useful complementary tool : comparison between Landsat tectonic interpretation and geophysical, air photographs and ground data seems to be the right method to improve our global knowledge. First Landsat 2 results in the tectonic domain confirm this idea. Practical applications of the method is in progress :

- Establish a sismo-tectonic map of France, to select power station sites.

- Characterize the location of ore deposits.

- Complete photogeological studies realized as a preparation for geological mapping of France at a 1/50 000.

The technical impact of this new data source on the B.R.G.M. activities is evident : engineering geologists, mining geologists and mapping geologists ask for ERTS interpretation and have already found in the field some evidence of its practical interest.