

General Disclaimer

One or more of the Following Statements may affect this Document

- This document has been reproduced from the best copy furnished by the organizational source. It is being released in the interest of making available as much information as possible.
- This document may contain data, which exceeds the sheet parameters. It was furnished in this condition by the organizational source and is the best copy available.
- This document may contain tone-on-tone or color graphs, charts and/or pictures, which have been reproduced in black and white.
- This document is paginated as submitted by the original source.
- Portions of this document are not fully legible due to the historical nature of some of the material. However, it is the best reproduction available from the original submission.

E7.6-10.0.80
CR-145885

APPLICATION OF REMOTE SENSING
FOR FISHERY RESOURCE
ASSESSMENT AND MONITORING

"Made available under NASA sponsorship
in the interest of early and wide dis-
semination of Earth Resources Survey
Program information and without liability
for any use made thereof."

SKYLAB EXPERIMENT NO. 240

CONTRACT No. T-8217B

PROGRESS REPORT NO. 20

REPORTING PERIOD: 1 September to 15 November 1975

Approved: 
K. J. Swadlow

Date Submitted: 11/26/75

(E76-10080) APPLICATION OF REMOTE SENSING
FOR FISHERY RESOURCE ASSESSMENT AND
MONITORING Progress Report, 1 Sep. - 15
Nov. 1975 (Mississippi Test Facility) 3 p
HC \$3.50

N76-14563

CSSL 08A G3/43

Unclas
00080

Technical Monitor: KH Faller
NASA/JSC Earth Resources Laboratory
National Space Technology Laboratories
Bay Saint Louis, Mississippi 39520

**APPLICATION OF REMOTE SENSING
FOR FISHERY RESOURCE
ASSESSMENT AND MONITORING**

INTRODUCTION

This is report #21 of a series of progress reports required by the Statement of Work for Skylab Experiment #240 entitled "Application of Remote Sensing for Oceanic Gamefish Assessment and Monitoring" under Contract No. T-8217B.

OVERALL STATUS

The S192 processing and analysis has been completed. The data from spectral bands 2, 3, 6 and 7 were used to develop a white marlin distribution model. The correlation coefficient for this model was .892 which was an improvement over .489 for a model constructed entirely with sea truth data (D_5). The significance level for this model was 90 percent as compared to 60% for the model D_5 . Therefore, the increase in precision of the model can be attributed to the data from the S192 sensor.