

General Disclaimer


One or more of the Following Statements may affect this Document

- This document has been reproduced from the best copy furnished by the organizational source. It is being released in the interest of making available as much information as possible.
- This document may contain data, which exceeds the sheet parameters. It was furnished in this condition by the organizational source and is the best copy available.
- This document may contain tone-on-tone or color graphs, charts and/or pictures, which have been reproduced in black and white.
- This document is paginated as submitted by the original source.
- Portions of this document are not fully legible due to the historical nature of some of the material. However, it is the best reproduction available from the original submission.

**NASA TECHNICAL
MEMORANDUM**

NASA TM X-71837

NASA TM X-71837



**A PRELIMINARY REPORT OF MULTISPECTRAL SCANNER
DATA FROM THE CLEVELAND HARBOR STUDY**

by Don Shook, Charles Raquet, Roger Svehla, Douglas Wachter,
Jack Salzman, Tom Coney, and Dick Gedney

Lewis Research Center
Cleveland, Ohio 44135

November, 1975

(NASA-TM-X-71837) A PRELIMINARY REPORT OF
MULTISPECTRAL SCANNER DATA FROM THE
CLEVELAND HARBOR STUDY (NASA) 40 P HC \$4.00
CSCI 13E

N76-13638

Unclas
05639
63/45

1. Report No.	2. Government Accession No.	3. Recipient's Catalog No.	
4. Title and Subtitle A PRELIMINARY REPORT OF MULTISPECTRAL SCANNER DATA FROM THE CLEVELAND HARBOR STUDY		5. Report Date	
		6. Performing Organization Code	
7. Author(s) Don Shook, Charles Raquet, Roger Svěhla, Douglas Wachter, Jack Salzman, Tom Coney, and Dick Gedney		8. Performing Organization Report No. E-8550	
		10. Work Unit No.	
9. Performing Organization Name and Address Lewis Research Center National Aeronautics and Space Administration Cleveland, Ohio 44135		11. Contract or Grant No.	
		13. Type of Report and Period Covered Technical Memorandum	
12. Sponsoring Agency Name and Address National Aeronautics and Space Administration Washington, D. C. 20546		14. Sponsoring Agency Code	
		15. Supplementary Notes	
16. Abstract A report on the remote sensing part of an E. P. A. study of the water quality in the Cleveland harbor area is presented. The study was performed for E. P. A. by the Ohio State University Center for Lake Erie Research, Case Western Reserve University, and the NASA Lewis Research Center. Imagery obtained from an airborne multispectral scanner is presented. A synoptic view of the entire study area is shown for a number of time periods and for a number of spectral bands. Using several bands, sediment distributions, thermal plumes, and Rhodamine B dye distributions are shown.			
17. Key Words (Suggested by Author(s))		18. Distribution Statement Unclassified - unlimited STAR Category	
19. Security Classif. (of this report) Unclassified	20. Security Classif. (of this page) Unclassified	21. No. of Pages	22. Price*

* For sale by the National Technical Information Service, Springfield, Virginia 22161

A PRELIMINARY REPORT OF MULTISPECTRAL SCANNER

DATA FROM THE CLEVELAND HARBOR STUDY

by Don Shook, Charles Raquet, Roger Svehla, Douglas Wachter,
Jack Salzman, Tom Coney, and Dick Gedney

Lewis Research Center

SUMMARY

A report on the remote sensing part of an E.P.A. study of the water quality in the Cleveland harbor area is presented. The study was performed for E.P.A. by the Ohio State University Center for Lake Erie Research, Case Western Reserve University and the NASA Lewis Research Center. Imagery obtained from an airborne multispectral scanner is presented. A synoptic view of the entire study area is shown for a number of time periods and for a number of spectral bands. Using several bands, sediment distributions, thermal plumes, and Rhodamine B dye distributions are shown.

INTRODUCTION

The Cleveland Harbor Study took place in the time period July 17 to July 29, 1975. The participants in this E.P.A. sponsored study were Case Western Reserve University, Ohio State University, and NASA Lewis Research Center. The objective of the study was the verification of E.P.A. models of flow and mixing of river pollutants in lake waters. To achieve this objective, in situ measurements were made of water quality, temperature, and flow patterns by both O.S.U. and C.W.R.U. personnel.

At the request of the E.P.A., multispectral imagery of the study area was taken by Lewis personnel while the in-the-water measurements were being made. This data was taken using the NASA C-47 aircraft. Imagery from the multispectral data was made available to O.S.U. and C.W.R.U. personnel within approximately 2 to 6 hours after completion of the flights. This imagery was used for planning the next set of measurements.

The study area was laid out on a grid, and sampling stations were chosen prior to the study. Measurements were made at these stations daily by O.S.U. personnel using the 60-foot research vessel Hydra. To aid in the river-flow pattern measurements, Rhodamine B dye was dumped into the water from the C.W.R.U. boat on three separate days during the study period.

SCANNER FLIGHTS AND SHIP OPERATIONS

A summary of activities during the study is shown in table I. As seen in the table, as many as twenty-seven flight lines were flown in one day. The days of maximum activity were July 25, 28, and 29. Three scanner flights were made prior to any O.S.U. ship operation. The first of these was a test flight, and the second and third flights were made to provide imagery to aid in choosing sampling sites. Sixteen sampling sites were used. These sites were in the vicinity of the river mouth and the breakwall. The actual locations are shown in figure 1. The time lines in the figure indicate the direction of the flight lines for the time of day flown. Table II shows in detail the flight lines that were flown during the study. The table contains all information on the flights such as time, direction, altitude, some notes on the weather, and camera information.

SCANNER IMAGERY

The scanner flown was a Bendix Modular Multispectral Scanner (M²S). Eleven channels of digital data were recorded simultaneously on tape. The central wavelength and bandwidth for each scanner band are the following:

Band number	Central wavelength nanometers	Bandwidth nanometers
1	410	60
2	465	50
3	515	40
4	560	40
5	600	40
6	640	40
7	680	40
8	720	40
9	810	100
10	1 015	90
11	11 500	1000

Images to be presented were obtained from the scanner high-density digital tape using a digital to analog converter and an EDO Western Model 572A fiber optics recorder using 3M type 7869 Dry Silver film. The EDO Western operator adjusted the gains and bias to give brightest areas (white) where the water reflectance was indicated to be the highest and the dark areas (black) where the water reflectance was lowest. This provided the greatest possible number of gray scales for areas of intermediate water reflectance. Since this was independently done for each band, it is not possible to accurately compare the brightness of different bands with each other.

The scanner imagery has been divided into three categories to show three properties of major interest to the E.P.A. of the water in the Cleveland harbor area during the study. These three properties are obtained directly from the scanner imagery without recourse to a detailed computer-aided analysis of the data. The first property to be discussed is total suspended solids. A synoptic view of the sediment distribution will be shown. Secondly the initial Rhodamine B dye distribution and degree of mixing with the sediment plume will be shown. Thirdly the surface temperature in the harbor area will be presented.

Sediment Plume

Figure 2 shows the harbor area on July 25 between 10:00 and 10:30 using band 4. The image is a mosaic of three flight lines made at 5000 feet. The flight line over the river mouth was made at 10:00 just before the Rhodamine B dye was dumped there. Referring to the sampling sites in figure 1, the figure 2 data indicates that relative to sites 10, 11, and 12, the total suspended solids at sites 1, 3, and 16 should be high. Sites 2, 4, 15, and 13 lie close to the plume edge so that their exact location and the sampling time is important. Sites 6, 7, and 8 should have an intermediate suspended solid concentration.

Figure 3 shows the harbor area also on July 25 but at 3:41 and 4:32 p.m. using band 6. The data show that the sediment plume has changed substantially from what it was in figure 2. The direction of the plume is affected by wind direction and lake currents. This imagery indicates that sites 13, 15, 16, and possibly 3 will show relatively less sediment in this distribution than the one at 10:00. The bright area in the image to the right of the harbor entrance and outside the breakwall could be the remnant of the morning plume. Figures 2 and 3 were made with different scanner bands. However, as shown in reference 1, bands in the green-to-red wavelength range show sediment plumes more or less equally well.

Figures 4 to 7 show the sediment distributions on the remaining days of the study. The images show the day to day variations in the sediment distribution. In figure 5(a) band 4 (green) and band 7 (red) image of the 10:02 flight data is shown. The bright area along the edge of the band 4 image is due to air scattering. The overall variation of the reflectance is similar for the two bands but with band 4 perhaps showing a greater difference in the water east of the river mouth compared to that west of the river mouth inside the breakwall. Figures 8 to 18 also show the sediment plume during the study. However, these will be discussed in terms of the dye distribution.

Dye Distribution

Rhodamine B dye was dumped in the river near the mouth on the morn-

ing of July 25. The dye was easily visible from the air during the morning flight and traces along the breakwall could be seen in the afternoon. Figure 8 shows three images made from data taken at 10:15, 10:27, and 10:38. In the 10:15 band 4 image, the dye is just visible as a small dark spot at the river mouth. In the 10:27 band 5 image, the dye appears as a bright spot at the river mouth and in the 10:38 band 4 image a dark spot. In figure 9 the 10:50 flight line using bands 4 and 5 shows the dye to have dispersed some and moved to the center of the breakwall area.

The reason the area of water containing the concentrated dye appears as a dark spot in the band 4 image and a bright spot in the band 5 image is the following: The peak of the dye fluorescence emission is in band 4 (ref. 2) but the absorption is also the highest. There is always a positive contribution by the dye fluorescence to the radiance even with extreme self-absorption. However, the net radiance in the dye area is much lower because the dye is absorbing the light that would have been reflected by the sediment and water. In band 5 the dye absorptance is less for the same concentration and the net radiance is higher than the surrounding water. Band 6 lies further in the wings of the dye fluorescence wavelength region and the concentrated dye area appears bright on the image. The dye was not detectable on the band 7 images. The variation of reflectance with dye concentration is, of course, very nonlinear.

Figure 10 shows images made from bands 4 and 8 from the 3:35 p.m. flight on July 25. A comparison of these images indicates that the dye has completely dispersed into the sediment distribution which is seen to be essentially the same for band 8 with no dye fluorescence contribution and band 4. Some difference in these images is apparent along the shore line.

On July 28 dye was dumped at the harbor entrance and in the harbor in the vicinity of sampling site 5. In figures 11 to 18 bands 5 and 7 imagery is shown which indicates areas of high dye concentration. The dye in the harbor area is difficult to see while that at the breakwall entrance is easily seen in the band 5 imagery. In reference 2 it is shown that the dye could be detected at concentrations as small as 1 PPB. Computer analysis of our multispectral data must be made to determine the sensitivity of the scanner. The final data will be only relative and will require the C.W.R.U. data in order to be put on an absolute basis.

Temperature Distribution

Band 11 thermal infrared imagery for July 25, 28, and 29 are shown in figures 19 to 21. Included with each image is a gray scale with the temperature corresponding to each shade of gray in the scale and in the image. This can be done because the band 11 detector views two known temperature sources during each rotation of the scan mirror. In order to apply this calibration to the scene below, air absorption must be accounted for (ref. 3). This effect causes the indicated surface tempera-

ture from 5000 feet to be approximately 3° C low depending on atmospheric conditions. The surface temperature differences should be given accurately by the gray scale.

The warm water plumes shown in figures 19 and 20 are similar in structure to the sediment plumes shown previously. As a general comment on the images, there appears to be little warm water flowing from the river mouth compared to that from the Cleveland Municipal Power Plant. During the time of the flights the land was much warmer than the water and therefore most land temperature variations are beyond the dynamic range of the film and are lost in the images. These land-temperature variations are contained in the data and could be displayed at the expense of losing the small temperature variations in the water.

SUMMARY OF RESULTS

Imagery made from multispectral scanner data taken during the Cleveland Harbor Study has been presented. The images provide a synoptic view of the study area and show the areas of high water reflectance due to high sediment loading.

Since the area was overflown repeatedly, a number of images of the study area were obtained. These showed that the sediment distribution can change in a few hours due to wind direction or other changes.

During the time of the study when a high concentration of Rhodamine B dye was present in the harbor area, the dye could be seen in images using bands 4, 5, and 6. In very high concentrations the dye area was darker than the surrounding water which dramatically demonstrated the nonlinearity of reflectance with dye concentration. Low concentrations of dye could not be distinguished from the sediment plume using film images.

It will be necessary to use computer analysis such as classification or band subtraction to investigate the dye plume and sediment plume in a quantitative manner.

Thermal infrared images for three of the days during the study showed the thermal plume from the Cuyahoga River. The river plume was significantly smaller than the plume from the Cleveland Municipal Power Plant, which could also be seen in the images.

REFERENCES

1. Svehla, Roger, et al.: Remote Sensing Study of Maumee River Effects on Lake Erie. NASA TM X-71780, 1975.

2. Eliason, J. R.; Foote, H. P.; and Doyle, M. J.: Surface Water Movement Studies Utilizing a Tracer Dye Imaging System. Proceedings of the Seventh International Symposium on Remote Sensing of Environment, Vol. 1, Univ. of Michigan, 1971, pp. 731-748.
3. Boudreau, Robert D.: Correcting Airborne Scanning Infrared Radiometer Measurements for Atmospheric Effects. NASA TM X-69940, 1972.

TABLE I. - SUMMARY OF ACTIVITIES

Date	Number of data flight lines	Ship operation		
		Hydra (OSU)	CWRU boat	NASA boat
7/17	4		(Dye dumped)	
7/18	13			
7/24	4			
7/25	22	X	(Dye dumped)	
7/26	20	X	X	X
7/27	12	X	X	
7/28	27	X	(Dye dumped)	X
7/29	21	X	X	X

TABLE II. - INSTRUMENT AND FLIGHT LOG

Flight designation	Date	Time	Run	Flight line from fig. 1	Offset right or left from flight line, n mi	Altitude, ft	Ground speed, knot	Camera frame number and time interval, sec	Camera notes	Notes	Weather
WQ 34	7/17	1442	1	1500 E→W	---	1500	135	26		Wheels up Late start.	
		1504						4.7			
		1505									
		1509						40			
		1513	2	1500 W→E	---	1500	142	4.7		[Plume moving to west; no dye apparent]	Hot sultry day; fairly uniform, hazy sky. Much haze.
		1519	3	1500 W→E	---	1500	139	36			
1523	4.7										
1530	57										
1534	4.3										
		1545	4	1500 W→E	---	1500	142				
WQ 35	7/18	942	1	1015	0.7 L	6000	140		Some shots taken	Wheels down	
		1002									
		1005									
		1014									
		1017	2	1015	1.0 R	6000	133				Bright hazy sky; no clouds, much haze

TABLE II. - Continued.

Flight designation	Date	Time	Run	Flight line from fig. 1	Offset right or left from flight line, n mi	Altitude, ft	Ground speed, knot	Camera frame number and time interval, sec	Camera notes	Notes	Weather		
WQ 35	7/18	1155	13	1115	3.5 R	1 500	138						
		1157											
WQ 36	7/24	1205											
		907											
		943	1	H1000	1.0 L	1 500	118	22			Wheels down.		
		945						5.7				Wheels up.	
		951	2	H1000	0.5 L	1 500	123	21					
		953						5.3					
WQ 37A	7/25	959	3	H1000	0	1 500	124	30					
		1002						5.3				Very hazy, overcast	
		1007	4	H1000	0.5 R	1 500	121	37					
		1010						5.3					
		1025											
		842											
		909	1	H0915	0.0	10 000	117	20					
		915						30.0					
919	2	H0915	0.5 R	10 000	180	13							
921		(Rev)				20.0				[Plume moving toward east]			
926	3	H0915	0.5 L	10 000	118	22							
931						22.0					Only scattered high clouds.		

TABLE II. - Continued.

Flight designation	Date	Time	Run	Flight line from fig. 1	Offset right or left from flight line, n mi	Altitude, ft	Ground speed, knot	Camera frame number and time interval, sec	Camera notes	Notes	Weather
WQ 38B	7/26	1509	1	H1530	0.2 L	10 000	165	11 25.7	III-4	Wheels up.	Very clear skies, just like s.m.
		1529	2	H1530	3.1 L	10 000	168	10 23.7			
		1533	3	H1600	1.3- 1.6 L	5 000	154	18 12.7			
		1543	4	H1600	3.1 L	5 000	159	18 12.7			
		1547									
		1601	5	H1630	0.0- 0.1 L	10 000	169	10 21.7	II5-6		
		1604	6	H1630	2.4 L	10 000	166	10 21.7			
		1614	7	C1700	-						
		1617	8	C1700	1.1 L	5 000	159	12 10.0			
		1620	9	C1700	1.3 L 14° L TKE	500	123	55 1.7			
		1628									
		1632									
		1640									
		1643									
1653											
1654											
1702											
1703											
1709											
1710											
1720											
-940	7/27									Wheels down. Wheels up.	
1601		1	H1015	0.6 L	5 000	127	16 14.7				
1004											

TABLE II. - Continued.

Flight designation	Date	Time	Run	Flight line from fig. 1	Offset right or left from flight line, n mi	Altitude, ft	Ground speed, knot	Camera frame number and time interval, sec	Camera notes	Notes	Weather	
WQ39A	7/27	1011	2	H1015	0.9 K	5 000	127	14			Generally clear day, some haze.	
		1015					14.7					
		1022	3	H1015	2.4 R	5 000	129	11				
		1027						13.7				
		1036	4	C1045	0.6 L	5 800	147	10				
		1037						14.7				
		1047	5	C1045	0.6 L	2 000	129	20				
		1048						4.2				
WQ 39B	7/27	1054	6	C1045	-0.5 L	500	126	34			Slightly improved visibility, but many more clouds.	
		1055						1.6				
		1104	7	H1115	-0.5 R 15° L TKE	2 000	118	35				
		1106						4.4				
		1116	8	H1115	-0.6 R 10° L TKE	5 000	128	17				
		1119						11.6				
		1129										
		1508										
		1521	1	H1530	0.9 R	5 000	164	26				
		1524						10.0				
1535	2	H1530	0.8 R	5 000	164	24						
1538						11.0						
1549	3	H1530	1.6 R	2 000	155	53						
1553						5.0						

TABLE II. - Continued.

Flight designation	Date	Time	Run	Flight line from fig. 1	Offset right or left from flight line, n mi	Altitude, ft	Ground speed, knot	Camera frame number and time interval, sec	Camera notes	Notes	Weather	
WQ 39B	7/27	1603	4	H1550	1.3 L	2 000	154	12				
		1604		(Overrib)				5.2				
WQ 40A	7/28	1614										
		851										
		918	1	C0930	0.8 L	10 000	130	14		III12-13	Wheels down. Wheels up.	
		921							25.5			
		926	2	H0930		10 000	176	14				
		929							20.0			
		938	3	C0930		5 000	147	31		III14-15		
		940							10.0			
		943	4	H0945		5 000	155	17				
		946							10.0			
		955	5	C0945		500	134	131				
956							1.7					
1000	6	H1000		1 500	142	43						
1002							4.3					
1012	7	H1015		5 000	144	20						
1015							12.7					
1023	8	H1015		5 000	139	20			III16			
1026							12.7					
1034	9	H1015		5 000	136	19			III17			
1038							12.7		III18			
1046	10	H1015		5 000	139	11						
1049							12.7					
1053	11	H1100		3 000	167	31			III19			
1055							4.8					

Clear day with some haze. Visibility 15-20 mi. However not as clear as 7/26.

TABLE II. - Continued.

Flight designation	Date	Time	Run	Flight line from fig. 1	Offset right or left from flight line, n mi	Altitude, ft	Ground speed, knot	Camera frame number and time interval, sec	Camera notes	Notes	Weather
WQ 40C	7/28	1552	3	H1545	0.1 R	10 000	165	12			Quite clear; visibility slightly better than a.m.
		1556						24.2			
		1606	4	C1600	0.0	10 000	170	4			
		1607						23.1			
		1616	5	C1630	0.1 L	5 000	152	8			
		1617						13.2			
		1622	6	C1630	0.3 L	1 500	155	25			
		1623			10° L TKE			2.5			
		1628	7	C1630	0.3 L	500	149	55			
		1629						1.4			
WQ 41A	7/29	1637	8	C1630	---	100	148	52			Wind out of west Wheels down. Wheels up. Little or no wind.
		1637						0.6			
		1647	9	H1645	0.1 R	5 000	160	21			
		1650						12.8			
		1656	10	H1645	0.8 L	5 000	154	17			
		1659						12.8			
		1710									
		840									
		859	1	H0915	0.2 L	5 000	143	12			
		902						14.7			
911	2	C0915	1.0 R	5 000	147	10					
912						13.7					

TABLE II. - Continued.

Flight designation	Date	Time	Run	Flight line from fig. 1	Offset right or left from flight line, n mi	Altitude, ft	Ground speed, knot	Camera frame number and time interval, sec	Camera notes	Notes	Weather	
WQ 41A	7/29	923	3	C0915	0.6 R	2 000	132	18			Clear skies, considerable haze; thick cloud from Myny drifting over harbor.	
		924						6.3				
		931	4	C0915	0.4 R	500	109	10				
		932							1.8			
		943	5	H1000	1.1 L	2 000	134	30				
		945							6.5			
		954	6	H1000	0.7 L	2 000	133	25				
		957							6.5			
		1006	7	H1000	0.1 L	2 000	132	27				
		1009							6.5			
		1018	8	H1000	0.4 R	2 000	132	25				
		1021							6.5			
1030	9	H1000	1.1 R	2 000	134	29						
1033							5.8					
1040	10	H1000	1.7 R	2 000	135	32						
1043							5.8					
1053	11	H1100	0.3 R	5 000	145	15						
1056			5° L TKE				13.7					
1105	12	H1100	1.0 L	5 000	145	12						
1107							13.7					
1118										Wheels down.		
1518	7/29											
1530		1	H1530	0.1 R	5 000	141	17				Wheels up.	
1534				4° TKE				14.6				

ORIGINAL PAGE IS
OF POOR QUALITY




Figure 1. - Cleveland area of Vermilion, Ohio chart showing numbered sampling stations and flight lines for the indicated time of day.

ORIGINAL PAGE IS
OF POOR QUALITY



Figure 2. - 7/25/75, Runs 6, 7, and 8, 10:15 AM Band 4

E - 6330





Figure 3. - 7/25/75, Band 6



4:03 PM, 10,000 ft. Altitude




4:24 PM, 5,000 ft. Altitude

Figure 4. - 7/26/75, Band 4

ORIGINAL PAGE IS
OF POOR QUALITY



Band 4



Band 7

Figure 5. - 7/27/75, 10:02 AM

ORIGINAL PAGE IS
OF POOR QUALITY




Figure 8. - 7/28/75, 3:24 PM, Band 5

ORIGINAL PAGE IS
OF POOR QUALITY





Figure 7. - 7/29/75, 3:30 PM, Band 4



Band 4, 10:15 AM



Band 5, 10:27 AM



Band 4, 10:38 AM

ORIGINAL PAGE IS
OF POOR QUALITY

Figure 8. - 7/25/75

F-8550



Band 5



Band 4

Figure 9. - 7/25/75, 10:50 AM

ORIGINAL PAGE IS
OF POOR QUALITY






Figure 10. - 7/25/75, 3:35 PM



Band 5

ORIGINAL PAGE IS
OF POOR QUALITY



Band 7

Figure 11. - 7/28/75, 9:28 AM

E-8550




Band 5

ORIGINAL PAGE IS
OF POOR QUALITY




Band 7

Figure 12. - 7/28/75, 9:45 AM



Band 5

E-6550




Band 7

Figure 13. - 7/28/75, 10:01 AM



Band 5




Band 7

Figure 14. - 7/28/75, 10:13 AM




Band 5




Band 7

Figure 15. - 7/28/75, 10:25 AM

E-8550



Band 5



Band 7


Figure 16. - 7/28/75, 10:54 AM

E-8550



Band 5

ORIGINAL PAGE IS
OF POOR QUALITY




Band 7

Figure 17. - 7/28/75, 11:00 AM



Band 5

ORIGINAL PAGE IS
OF NOE QUANTITY



Band 7

Figure 18. - 7/28/75, 4:49 PM

E-8550



ORIGINAL PAGE IS
OF POOR QUALITY




Figure 19. - 7/25/75, 3:35 PM, Thermal Infrared, Band 11





Figure 20. - 7/28/75, 3:22 PM, Thermal Infrared, Band 11

E-8550



26.4° C or higher

25.6

24.8

24.0

23.2

22.4

21.6

20.8

20.0 or lower




Figure 21. - 7/29/75, 3:30 PM, Thermal Infrared, Band 11