

Grant NGR 39-011-113

National Aeronautics and Space Administration

(NASA-CR-140960) STUDIES OF THE AIRGLOW, N75-12473
 THE AURORA, THE ION AND NEUTRAL
 COMPOSITION, AND THE CHEMISTRY OF THE
 TERRESTRIAL ATMOSPHERE (Pittsburgh Univ.) Unclas
 30 p HC \$3.75 CACL 04A G3/46 17002

Studies of the Airglow, the Aurora, the Ion and Neutral Composition
 and the Chemistry of the Terrestrial Atmosphere

[Previously: "Helium Geocorona Rocket Experiment"]

October 1, 1974

The University of Pittsburgh
 Department of Physics



Approved by:

Submitted by:

J. Rosenberg, Dean
 Faculty of Arts and Sciences

E. C. Zipf, Jr.
 E. C. Zipf, Jr.
 Principal Investigator

Paul Solyan
 Comptroller and Assistant Treasurer

P. Stehle
 P. Stehle
 Chairman, Department of Physics

I. A Report On

Studies of the Airglow, the Aurora, the Ion and Neutral Composition,
and the Chemistry of the Terrestrial Atmosphere

A familiar proverb reminds us that "A picture is worth a thousand words!" Figure 1 succinctly demonstrates this point for it shows the first altitude profile of atomic oxygen and N_2 in the mesosphere obtained by an optical mass spectrometer [OMS]. This instrument was launched on 14 August 1973 at 11:45 pm MST into the atmosphere above White Sands, N. M. on board Nike-Apache rocket 14.512 UA. This experiment was one of five conducted successfully this year by the University of Pittsburgh aeronomy group:

- (1) Javelin rocket (8.61 UA), 17 July 1973, high altitude (200 - 740 km) OMS and 1304\AA airglow experiment, Wallops Island, Va.
- (2) Nike-Apache rocket (14.512 UA), 14 August 1973, low altitude (90 - 115 km) OMS experiment, White Sands, N. M.
- (3) Black Brant V vehicle (AMD-VB-35), January 1974, double-mode mass spectrometer - 1304\AA resonance fluorescence/absorption experiment, and nitric oxide auroral studies, Fort Churchill, Manitoba.
- (4) Nike-Apache rocket (14.526 UA), 21 February 1974, double-mode mass spectrometer and multiple photometer for auroral composition and excitation studies, Fort Churchill, Manitoba.




Figure 1





Figure 2

- (5) Nike-Apache rocket (14.528 UA), 2 March 1974, double-mode mass spectrometer and multiple photometer for auroral composition and excitation, Fort Churchill, Manitoba.

1. Composition Measurements

The optical mass spectrometer is a new and unique instrument. The OMS combines a laboratory-quality, focused electron gun and a 1/8 - meter UV monochromator (Figure 3) which analyses the gas excited by the electron beam. As the monochromator scans the $\lambda 1200$, 1215 , and $1304\overset{\circ}{\text{A}}$ resonance lines by N, H, and O respectively, the energy of the electron beam is switched from 14.5 eV to 75 eV every 20 msec to exploit the difference between direct and dissociative excitation. In the low energy mode the OMS measures the density of the atomic species without interference from the related, parent molecular gases. The O_2 , N_2 , and H_2 concentrations are then measured separately in the high energy mode. The instrument does not require active pumping of the sampled gas and uses an open end, electron-gun-design with a very large conductance.

This instrument has been further improved to increase its sensitivity so that atomic N, O, or H concentrations as little as 10^5 atoms/cm³ can now be detected at mesospheric altitudes. The modified OMS will be flown as part of Project Alladin from Wallops Island, Va. during the week of 17 June 1974. This complex series of rocket-borne experiments will provide us with the opportunity to cross-compare a variety of techniques for measuring atomic oxygen at



$\frac{1}{8}$ - METER UV MONOCHROMATOR

Figure 3

mesospheric altitudes and above, and hopefully resolve several vexing problems concerning this highly reactive species. However, only the OMS has the necessary sensitivity and discrimination to measure the N and H concentrations in the D and E regions.

Project Alladin is a natural extension of our joint efforts with scientists from York University (G. Shephard and R. A. Young) to compare atomic oxygen concentrations inferred from conventional quadrupole mass spectrometer data with the results of a resonance absorption/fluorescence experiment on board the same rocket. This study was carried out successfully this winter at Fort Churchill on board a Black Brant V rocket (AMD-VB-35) launched during January (1974). These experiments will be continued in the fall with the launch of an additional Black Brant V payload weighing nearly 600 lbs. (AMD-VB-34).

The composition data of Figure 1 agrees well with the results obtained by Von Zahn and his collaborators using a liquid helium cooled mass spectrometer. The OMS data is in relatively poor agreement with the atomic oxygen profiles inferred from OGO VI $\lambda 5577$ airglow studies. The OI green problem is clearly still very much with us both in the airglow and in the aurora! Figure 2 compares our OMS data with a recent Jacchia model (1971). Both the magnitude and altitude profile for N_2 are in good agreement with the model values; however, the O data are not. In order to facilitate a comparison between the OMS results and the Jacchia model, we have reduced the magnitude of the O concentration in the model and in Figure 2 by about a factor of 2. The agreement is acceptable near 120 km but the observed O density near 100 km is still considerably smaller and the O concentration gradient noticeably less than that

predicted by the model atmosphere. Future OMS experiments will help bring these apparent discrepancies into sharper focus, and perhaps explain their origin.

The launch date for the initial test of the OMS instrument at mesospheric altitudes was chosen to coincide with the seasonal minimum predicted on the basis of the OGO VI airglow experiment. Project Alladin will give us an additional opportunity to see whether other OGO VI predictions are borne out under quite geophysical circumstances.

2. Auroral Studies

Four rocket experiments involving complex payloads were carried out at the Churchill Research Range during January and February 1974. The first vehicle, a Black Brant V, was prepared in collaboration with York University and was launched into a IBC II aurora during January. The University of Pittsburgh contributed a double-mode mass spectrometer in order (1) to continue our studies of nitric oxide enhanced aurora and (2) to compare atomic oxygen concentrations measured by this instrument with similar results obtained using $\lambda 1304\text{\AA}$ resonance absorption/fluorescence techniques. Both instruments performed flawlessly throughout the flight and a detailed analysis of the results is currently underway. A followup experiment has been planned and will be flown from Fort Churchill on board Black Brant V vehicle AMD-VB-34.

This year was notable in several respects. It began, for example, with the first successful OMS experiment. It moved on to the successful execution of a two-payload, three-rocket mission to the auroral zone.

And it concludes with active participation in Project Alladin. The auroral mission deserves special comment. In previous reports and proposals we had lamented the fact that progress in auroral physics was being impeded by the one experiment per year philosophy, and that a more effective approach would be (1) to construct two different, but complementary, payloads, (2) take them both to Churchill and launch them into suitable aurora, (3) recover both payloads, (4) repair them in the field and then (5) re-fly the payload that promised the greatest potential dividends based on an "in situ" review of what was learned on the first two flights. This new approach is actually more economical over the long run and makes good scientific sense. A mission of this kind was approved for the February 1974 auroral campaign.

In spite of the severe conditions that prevail at Churchill during the winter months we were able to refurbish and recalibrate completely a recovered payload (14.526 UA) in two days amply demonstrating the feasibility of this *modus operandi* when the operation is in the hands of an experienced group of scientific, vehicle and telemetry personnel. The breakup of Nike-Apache rocket 14.527 UA, although very unfortunate, does not in any way diminish this tactical accomplishment. Each vehicle was in fact launched into a major auroral substorm characterized by auroral forms 50 kR or more in intensity, and the mission was executed precisely as planned. The ultimate fate of 14.527 UA, however, was not what we had hoped for, and its disintegration was particularly painful because it carried our new extreme ultraviolet [EUV] experiment. No data was




Figure 4

obtained.

The impact of this development was felt acutely because at the same time we had been making outstanding progress at measuring the absolute cross sections for exciting auroral EUV feature in our complementary laboratory program. Figures 4 and 5 of this report show some typical EUV spectra in the 900Å and 1010Å region produced by electron impact excitation of N_2 . The laboratory spectrum is complex and the EUV features are quite bright; the same is likely to be true in a well-developed auroral storm. Although we were disappointed by the untimely demise of 14.527 UA, the data of Figures 4 and 5 show that the experiment was and still is a very good one! We are planning to return to Churchill this winter in a continuing quest for this elusive prize: the measurement of the intensity and altitude distribution of auroral EUV radiation in the spectral range 500Å to 1300Å.

The 1974 auroral expedition was an outstanding success overall because two rockets were launched successfully into auroras that were ideal for the continuance of our studies of nitric oxide enhancements at northern latitudes. Nike-Apache payload 14.526 UA was launched into an intense magnetospheric storm in which the OI green line ($\lambda 5577\text{Å}$) reached an intensity of nearly 100 kR and the H component of the earth's magnetic field exhibited a negative bay in excess of 600 gammas. Enhanced NO concentrations ($\sim 2 \times 10^9 \text{ NO/cm}^3$) were encountered. The buildup was similar in magnitude to earlier nitric oxide results obtained in a comparable IBC II⁺ aurora (see Figure 6). Characteristically perturbed ion profiles (large NO^+/O_2^+

II




Figure 5

ratios) were also observed. Flight 14.528 UA was launched into an aurora approximately as bright, but not as active magnetically, and with a shorter integration time. In this case comparatively little NO was observed. These flights further delineate a trend that we had noted earlier; namely, the magnitude of the NO enhancement is in proportion to the total energy deposited in the region through which the rocket passes. And, if we distinguish between the energy deposited by precipitating electrons versus the energy dissipated in the ionosphere by the auroral electrojet via Joule heating and other related electric field phenomena, these flights clearly show that the energy associated with the electrojet and with the perturbation of the earth's magnetic field is by far the most important factor.

Figure 7 presents some preliminary data showing this general trend. It is clear that the NO buildup is directly proportional to $\int \Delta H^2 \cdot \Delta t$; this quantity is a measure of the total energy deposited by Joule heating. The 1974 winter campaign to Churchill has provided data that will allow us to separate quantitatively the two components leading to NO enhancements. Equally important, our results stress the importance of having adequate magnetographic coverage. This point has been made privately to the Air Force Cambridge research group responsible for their auroral studies [Project Icecap]. Even they are convinced now that large NO enhancements do in fact occur (an advance over their attitude one year ago). But like ourselves, they are not sure how this remarkable production feat is accomplished so efficiently.

Our laboratory program has, however, suggested a partial explanation of this phenomena. Our findings are summarized in the

following abstract of a paper that we presented at the Washington meeting of the AGU (Spring 1974).

Comment on the Production of N(²D) Atoms by Photo-absorption Processes and by Electron-Impact Excitation of N₂. Laboratory studies on the production of extreme ultraviolet radiation by electron-impact excitation of N₂ show that N(²D) atoms are produced very efficiently by the predissociation of N₂ Rydberg and valence states with ¹Π_u symmetry. This dissociation process involves a nonradiative transition to the C' ³Π_u continuum: $e + N_2 \rightarrow N_2(^1\Pi_u) + e$ followed by $N_2(^1\Pi_u) \rightarrow N_2(C' \ ^3\Pi_u) \rightarrow N(^2D) + N(^4S)$. Under auroral conditions these predissociating ¹Π_u states are populated by electron-impact at a rate comparable to the N₂⁺ ionization rate. Thus, the total N(²D) population rate from dissociative excitation, from dissociative recombination of NO⁺ ions, and from ion molecule reactions could be as much as 50 - 100 times the λ3914 volume emission rate. This excitation rate would suffice to account for the large N(²D) concentrations observed recently by Rusch and Sharp in an IBC I⁺ aurora, and for the production of unusually large nitric oxide densities in some auroral forms as a result of the [N(²D) + O₂] quenching reaction. Under daytime conditions these same N₂(¹Π_u) states are populated by absorption of solar photons and the N(²D) production rate from this process is shown to be comparable to that from photoelectron impact and from the dissociation

LOGARITHMIC SCALE




Figure 6






Figure 7

recombination of NO^+ ions.

Once again in this work we have a very good example of the remarkable effectiveness of complementary laboratory and rocket-borne programs.

These developments have prompted us to formulate an ambitious set of plans for our return to the auroral zone this winter. The measurement of auroral EUV fluxes and the assessment of the role of these photons in a variety of photoionization and photodissociation processes (e.g., the excitation of $\text{O}(^1\text{S})$ atoms and the green line) is a major goal. Further work on the excitation and ultimate fate of $\text{N}(^2\text{D})$ atoms in the aurora, on the role of joule heating, and on the possibility that plasma instabilities (anomalous resistivity) may be involved are also high on our list of major projects. Our nitric oxide studies will be expanded further to include an investigation of the so-called auroral mystery feature which is observed occasionally at $\lambda 2150\text{\AA}$. This band has been observed by rocket-borne and satellite (OGO-IV) spectrometer, but not in every aurora. It is commonly believed that this emission feature is actually the (1,0) gamma band of NO excited by the deactivation of metastable $\text{N}_2(\text{A}^3\Sigma_u^+)$ molecules,




The critical experiment, in which the three principal actors in the drama [$\text{N}_2(\text{A}^3\Sigma_u^+)$, NO, and the intensity of the 2150\AA feature] are

measured simultaneously and Reaction (1) verified, has yet to be performed. If this reaction sequence could be verified, then the OGO IV spectrometer data could be used to study the morphology of nitric oxide aurora. This development would lead to a much clearer picture of the scope of the whole NO phenomena as well as give a new meaning and significance to unused OGO data. Thus the 1975 auroral campaign has impressive potential.

3. New and Bizarre Developments of Aeronomic Importance

One of the charms and ultimate frustrations of aeronomy is the regularity with which "established" theories become casualties of new experimental advances. In the jargon of the young, this is an endeavor that requires "staying loose". Figure 8 illustrates this point.

Figure 8 shows the vacuum ultraviolet spectrum of a recombining plasma in which vibrationally excited CO_2^+ ions predominate; this is the first measurement of its kind. Our results indicate strong excitation of the CO fourth positive system ($A^1\Pi \rightarrow X^1\Sigma$) due to the dissociative recombination process:



It is important to note that only the $v' = 0$ and 1 levels of the $A^1\Pi$ state can be populated if the recombining CO_2^+ ions are in thermal equilibrium at 300°K. It is quite clear from the data shown in Figure 8 and from other spectra that we have obtained, that $A^1\Pi$




Figure 8

vibrational levels up to at least $v' = 5$ are in fact being populated. Complementary electron heating experiments have shown conclusively that process (2) is exciting all of these levels (Gutcheck and Zipf, 1973).

The implications of the results are quite clear: CO_2^+ laboratory plasmas, which had previously been thought to be equilibrated at 300°K, are in fact vibrationally very hot. This finding raises serious questions about previous total recombination-coefficient measurements involving the important atmospheric ions O_2^+ , NO^+ , and N_2^+ . Were these plasmas also vibrationally hot? Can these laboratory results be used now in the analysis of ionospheric data without reservation? Does, in fact, the nominal factor of 2 - 3 agreement achieved by aeronomists when they model daytime ion composition data, really tell us that our own ionosphere, as well as that of Venus has a substantial population of vibrationally excited ions?

Our preliminary data contain another striking result. Figure 8 shows a very intense spectral feature near 1400Å which does not appear conspicuously in electron impact excitation of CO or CO_2 (Mamma and Zipf, 1972; Mamma et al., 1973). This is the first laboratory observation of the so-called "mystery" spectral feature at 1400Å that appears so strikingly in the Venus airglow spectra obtained by Rottman and Moos (JGR 78, 8033, 1973). We have been able to show (Unger and Zipf, 1974) that the 1400Å feature is produced by the dissociative recombination of vibrationally excited CO_2^+ ions. However, at this time we do not know what specific quantum transition is involved.

Our experiment provides virtually conclusive evidence for the

presence of substantial concentrations of vibrationally excited CO_2^+ ions in the Venus ionosphere. The Mars data in the same wavelength region shows a complete absence of this VUV feature implying a relatively cool ionosphere (and CO_2^+ plasma) [Moos, 1974]. It is not clear at this point why the Venus atmosphere is capable of sustaining a vibrationally hot ionosphere while Mars does not or can not.

What about our own ionosphere? Could it be that some of the so-called anomalies in the ion chemistry of the aurora and of other disturbed atmosphere phenomena are really consequences of vibrationally excited ions? Laboratory results of the caliber of those shown in Figure 8 stir the geophysical pot in a profound manner. They remind us that in spite of our best efforts, some popular ionospheric theories, like the smile of the Cheshire Cat, will ultimately fade away. Our oversimplified models of a planetary ionosphere are in need of an overhaul, a major goal of our research program.

The data for Figure 8 was obtained in a plasma spectroscopy experiment carried out recently by our group. These results have already influenced our plans for Churchill next winter. In effect early access to this kind of knowledge gives our rocket program a significant competitive edge - we know where the future action is going to be thanks to the symbiotic relationship that exists between both efforts and we are aware of the techniques needed to study these problems using rocket-borne instrumentation. It is to the credit of the National Aeronautics and Space Administration that the effectiveness of the approach was recognized and the effort nurtured.

II. REFERENCES

Gutcheck, R. A., and E. C. Zipf, "Excitation of the CO Fourth Positive System by the Dissociative Recombination of CO_2^+ Ions", J. Geophys. Res., 78, 5429, 1973.

Mumma, M. J., E. J. Stone and E. C. Zipf, "Excitation of the CO Fourth Positive Band System by Electron Impact on Carbon Monoxide and Carbon Dioxide", J. Chem. Phys., 54, No. 6, 2627, 1971.

Mumma, M. J. and E. C. Zipf, "Calibration of Vacuum Ultraviolet Monochromators by the Molecular Branching-Ratio Technique", J. Opt. Soc., 61, 83, 1971.

Mumma, M. J. and E. C. Zipf, "Dissociative Excitation of Vacuum Ultraviolet Emission Features by Electron Impact on Molecular Gases*: I. H_2 and O_2 ", J. Chem. Phys., 55, 1661, 1971.

Mumma, M. J. and E. C. Zipf, "Dissociative Excitation of Vacuum-Ultraviolet Emission Features by Electron Impact on Molecular Gases II. N_2 ", J. Chem. Phys., 55, 5582, 1971.

Mumma, M. J., E. J. Stone, W. L. Borst and E. C. Zipf, "Dissociative Excitation of Vacuum Ultraviolet Emission Features by Electron Impact on Molecular Gases III. CO_2 ", J. Chem. Phys., 57, 68, 1972.

Rottman, G. J. and W. H. Moos, "UV Spectrum of Venus", J. Geophys. Res., 78, 8033, 1973.

VI. PUBLICATIONS AND TECHNICAL PRESENTATIONS

1. E. C. Zipf, Jr., "Automatic Charge Information Devices", Technical Memorandum #56-5144-18, Bell Telephone Laboratories, New Jersey, 1956.
2. E. C. Zipf, Jr., "Transient Studies of De-Ionization Processes in the Nitrogen Afterglow", Johns Hopkins University, Spectroscopic Report No. 20, Baltimore, Maryland, 1961.
3. H. M. Crosswhite, E. C. Zipf, Jr., and W. G. Fastie, "Far Ultra-violet Auroral Spectra", J. Opt. Soc. Am., 52, 643, 1962.
4. E. C. Zipf, Jr., and D. F. Heath, "Photometer Measurements of the Day and Night Airglow from 80 to 210 km", Trans. Am. Geophys. Union, 43, 436, 1962.
5. E. C. Zipf, Jr., "Ambipolar Diffusion in Low-Energy Nitrogen Plasmas", Bull. Am. Phys. Soc., 8, 311, 1963.
6. E. C. Zipf, Jr., and W. G. Fastie, "An Observation of Day Airglow Emission at 6300Å", J. Geophys. Res., 63, 6208, 1963.
7. E. C. Zipf, Jr., "Measurements of the Diffusion Coefficient and Radiative Lifetime of Nitrogen Molecules in the $A^3\Sigma_u^+$ State", J. Chem. Phys., 38, 2034, 1963.
8. E. C. Zipf, Jr., "Afterglow Studies on the Destruction of the Metastable $A^3\Sigma_u^+$ State of Nitrogen", Bull. Am. Phys. Soc., 9, 185, 1964.
9. W. G. Fastie and E. C. Zipf, Jr., "Airglow Emission During a Total Solar Eclipse", Trans. Am. Geophys. Union 45, 40, 1964.
10. E. C. Zipf, Jr., and W. G. Fastie, "An Observation of the (0,0) First Negative Band of N_2^+ in the Dayglow", J. Geophys. Res., 69, 2357, 1964.
11. E. C. Zipf and W. G. Fastie, "The Concentration of N_2^+ Ions in the Daytime E and F_1 Regions", Trans. Am. Geophys. Union, 45, 49, 1964.
12. E. C. Zipf, "Excitation of the Dayglow by Solar Radiation", Bull. Am. Phys. Soc. 9, 539, 1964.
13. E. C. Zipf, "Metastable-Metastable Collisions in Nitrogen", Bull. Am. Phys. Soc., 10, 179, 1965.
14. E. C. Zipf, "On the Measurement of Recombination Radiation in the Afterglow", Bull. Am. Phys. Soc., 11, 494, 1966.

15. E. C. Zipf, "Rocket Measurements of the Visible Dayglow", J. Geomagnetism and Geoelectricity, 18, 301, 1966.
16. E. C. Zipf, "Time Dependent Measurements of Metastable Oxygen Atoms in the Afterglow", Bull. Am. Phys. Soc., 12, 200, 1967.
17. T. M. Donahue, T. D. Parkinson, E. C. Zipf, J. P. Doering, W. G. Fastie, and R. E. Miller, "Excitation of the Auroral Green Line by the Dissociative Recombination of Oxygen Molecular Ions: Analysis of Two Rocket Experiments", Planet. Space Sci., 16, 737, 1968.
18. L. F. Keyser, F. Kaufman and E. C. Zipf, "Radiative Lifetime and Collisional Deactivation of NO₂ Fluorescence and Nature of the Air Afterglow", Chem. Phys. Letters 2, 523 (1969).
19. E. C. Zipf, "The Collisional Deactivation of Metastable Atoms and Molecules in the Upper Atmosphere", Canad. J. Chem. 47, 1863 (1969).
20. T. D. Parkinson, E. C. Zipf and T. M. Donahue, "Rocket Observation of a Temporal Variation in an Aurora", Trans. Am. Geophys. Union 50, 258 (1969).
21. T. D. Parkinson, E. C. Zipf and T. M. Donahue, "Phase and Amplitude Studies of the $\lambda 3914$ and $\lambda 5577$ Emission in Pulsating Auroras", Trans. Am. Geophys. Union 50, 272 (1969).
22. T. D. Parkinson, E. C. Zipf and T. M. Donahue, "Rocket Investigation of the Auroral Oxygen Red Line", Trans. Am. Geophys. Union, 50, 257 (1969).
23. E. C. Zipf, T. D. Parkinson and T. M. Donahue, "Positive Ion Composition in an IBC I Aurora", Trans. Am. Geophys. Union 50, 258 (1969).
24. E. C. Zipf, "Transport Properties of Metastable O(¹S) Atoms", Bull. Am. Phys. Soc., 14, 264 (1969).
25. E. C. Zipf, W. L. Borst and T. M. Donahue, "A Mass Spectrometer Observation of NO in an Auroral Arc", J. Geophys. Res., 75, 6371 (1970).
26. T. D. Parkinson, E. C. Zipf and T. M. Donahue, "The Dissociative Excitation of the Auroral Oxygen Green Line by Electron Impact", Planet. Space Sci., 18, 187, 1970.
27. T. D. Parkinson, E. C. Zipf, and K. A. Dick, "An Observation In Situ of an Auroral Pulsation", J. Geophys. Res., 75, 1334 (1970).
28. W. L. Borst and E. C. Zipf, "Excitation of O₂⁺ First Negative Bands by Electron Impact on O₂", Phys. Rev. A, 1, No. 5, 1410, 1970.

29. W. L. Borst and E. C. Zipf, "Cross Section for Electron Impact Excitation of the (0,0) First Negative Band of N_2^+ from Threshold to 3 keV", Phys. Rev., 1A, 834 (1970).
30. W. L. Borst and E. C. Zipf, "Electron Excitation Cross Sections for the First Negative Bands of Oxygen", Phys. Rev., 1A, 1410 (1970).
31. R. Fralick and E. C. Zipf, "A Versatile Electrical Shutter for Optical Pumping Experiments", Rev. Sci. Instruments, 41, 47 (1970).
32. T. M. Donahue, E. C. Zipf and T. D. Parkinson, "Ion Composition and Ion Chemistry in an Aurora", Planet. Space Sci., 18, 171, 1970.
33. T. D. Parkinson, E. C. Zipf and T. M. Donahue, "Rocket Investigation of the Auroral Green Line", Planet. Space Sci., 18, 187, 1970.
34. T. D. Parkinson and E. C. Zipf, "Energy Transfer from $N_2(A^3\Sigma_u^+)$ as a Source of $O(^1S)$ in an Aurora", Planet. Space Sci., 18, 895 (1970).
35. E. C. Zipf, "The Dissociative Recombination of O_2^+ Ions into Specifically Identified Final Atomic States", Bull. Am. Phys. Soc., 15, 418 (1970).
36. M. J. Mumma and E. C. Zipf, "The Dissociative Excitation of Atomic Resonance Lines by Electron Impact", Bull. Am. Phys. Soc., 15, 422 (1970).
37. W. L. Borst and E. C. Zipf, "Electron Excitation Cross Sections of the First Negative Bands of Oxygen and Nitrogen", Bull. Am. Phys. Soc., 15, 421 (1970).
38. E. C. Zipf, "A Voltage Programmed Oscillator for Quadrupole Mass Spectrometers", Rev. Sci. Instruments, 41, 1236, 1970.
39. P. D. Feldman, J. P. Doering, W. G. Fastie and E. C. Zipf, "Excitation of $O(^1S)$ Atoms in the Day Airglow", Trans. Am. Geophys. Union 50, 750 (1970).
40. E. C. Zipf, W. L. Borst and T. M. Donahue, "Some Implications of a Large NO Abundance in an Auroral Arc", Trans. AGU., 51, 790, 1970.
41. W. C. Wells, W. L. Borst and E. C. Zipf, "Translational Spectroscopy of Fast Metastables Produced by Electron Impact Dissociation of Atmospheric Gases", Bull. Am. Phys. Soc., 17, 397, 1972.
42. E. C. Zipf and E. J. Stone, "Electron Impact Excitation of Atomic Nitrogen", Bull. Am. Phys. Soc., 17, 397, 1972.
43. R. A. Gutcheck and E. C. Zipf, "Collisional Deactivation of Metastable Ar($^3P_{0,2}$) Atoms and $N_2(^3\Sigma_u^+)$ Molecules", Bull. Am. Phys. Soc., 17, 395, 1972.

44. E. J. Stone and E. C. Zipf, "Electron Impact Excitation of the Werner Bands of H_2 ", Bull. Am. Phys. Soc., 17, 397, 1972.
45. P. D. Feldman, J. P. Doering and E. C. Zipf, "Excitation of $O(^1S)$ Atoms in the Day Airglow", J. Geophys. Res., 76, 3087, 1971.
46. T. M. Donahue, E. C. Zipf, and D. E. Shemansky, "Deactivation of $N_2 A^3\Sigma_u^+$ Molecules in the Aurora", Trans. AGU, 52, 309 (1971).
47. D. E. Shemansky, T. M. Donahue and E. C. Zipf, "Differential Electron Flux as Determined by Auroral Observations of the N_2 Positive and N_2^+ Systems", Trans. AGU, 52, 305 (1971).
48. E. C. Zipf, E. J. Stone, M. J. Mumma, W. C. Wells, and W. L. Borst, "Excitation of the $OI(^3S)$ and $OI(^5S)$ States by Electron Impact on O and O_2 in the Aurora, Dayglow and EEUV", Trans. AGU, 52, 308 (1971).
49. M. J. Mumma and E. C. Zipf, "Dissociative Excitation of Vacuum Ultraviolet Emission Features by Electron Impact on Molecular Gases: I. H_2 and O_2 ", J. Chem. Phys., 55, 1661, (1971).
50. W. C. Wells, W. L. Borst and E. C. Zipf, "Production of Energetic Long-Lived High-Rydberg States of Atomic Nitrogen by Electron Impact Dissociation of N_2 ", Proc. ICPEAC VII, North-Holland Publishing Co. (Amsterdam, 1971) p. 576.
51. E. C. Zipf, E. J. Stone, "Photoelectron Excitation of Atomic Resonance Radiation in the Terrestrial Airglow", J. Geophys. Res., 76, 6865, 1971.
52. W. C. Wells, W. L. Borst and E. C. Zipf, "Absolute Cross Section for the Production of $O(^5S^0)$ by Electron Impact Dissociation of O_2 ", Chem. Phys. Letters, 12, 288, 1971.
53. E. J. Stone and E. C. Zipf, "Excitation of the $OI(^3S)$ and $NI(^4P)$ Resonance States by Electron Impact on O and N ", Phys. Rev., A 4, 610 (1971).
54. M. J. Mumma, E. J. Stone and E. C. Zipf, "Non-Thermal Rotational Population of the $CO(A^1\Pi)$ State Produced by Dissociative Excitation of CO_2 by Electron Impact", Proc. ICPEAC VII, North-Holland Publishing Co. (Amsterdam, 1971) p. 858.
55. D. E. Shemansky, E. C. Zipf and T. M. Donahue, "Deactivation of $N_2 A^3\Sigma_u^+$ Molecules in the Aurora", Planet. Space Sci., 19, 1669, 1971.
56. T. D. Parkinson and E. C. Zipf, "The Auroral Intensity Ratio of $OI(6300\text{\AA})$ and $OI(5577\text{\AA})$ ", Planet. Space Sci., 19, 267 (1971).

57. D. E. Shemansky, T. M. Donahue and E. C. Zipf, "Differential Electron Flux as Determined by Auroral Observations of the N_2 Positive and N_2^+ Systems", submitted to J. Geophys. Res., (1971).
58. M. J. Mumma, E. J. Stone, and E. C. Zipf, "Excitation of the CO Fourth Positive Band System by Electron Impact on CO and CO_2 ", Bull. Am. Phys. Soc., 16, 218 (1971).
59. W. L. Borst and E. C. Zipf, "Energies of Metastables Produced in Electron Impact Dissociation of Atmospheric Gases", Bull. Am. Phys. Soc., 16, 205 (1971).
60. E. C. Zipf, "The Collisional Deactivation of Metastable $O(^1S)$ Atoms by O_2 and CO_2 ", J. Chem. Phys., submitted (1971).
61. E. J. Stone and E. C. Zipf, "Excitation of Resonance States in Electron Impact on Atomic Oxygen and Nitrogen", Bull. Am. Phys. Soc., 16, 217 (1971).
62. W. L. Borst and E. C. Zipf, "Lifetimes of Metastable CO and N_2 Molecules", Bull. Am. Phys. Soc., 16, 205, 1971.
63. M. J. Mumma, E. J. Stone and E. C. Zipf, "Excitation of the CO Fourth Positive Band System by Electron Impact on Carbon Monoxide and Carbon Dioxide", J. Chem. Phys., 54, No. 6, 2627, 1971.
64. M. J. Mumma and E. C. Zipf, "Calibration of Vacuum Ultraviolet Monochromators by the Molecular Branching-Ratio Technique", J. Opt. Soc., 61, 83 (1971).
65. W. L. Borst and E. C. Zipf, "Energy Spectra of Metastable Oxygen Atoms Produced by Electron Impact Dissociation of O_2 ", Phys. Rev. A 4, 153 (1971).
66. W. L. Borst and E. C. Zipf, "Lifetimes of Metastable CO and N_2 Molecules", in press, Phys. Rev. A, 3, 979 (1971).
67. E. J. Stone and E. C. Zipf, "Electron-Impact Excitation of Atomic Nitrogen in the Terrestrial Airglow", Trans. AGU 52, 876, 1971.
68. W. C. Wells, W. L. Borst and E. C. Zipf, "Translational Spectroscopy of Fast Metastable Fragments Produced by Electron Impact Dissociation of Atmospheric Gases", Trans. AGU, 52, 878, 1971.
69. E. C. Zipf, W. L. Borst and T. M. Donahue, "Positive Ion Composition of the Dawn Ionosphere During a Solar Flare", Trans. AGU, 52, 885, 1971.
70. R. Schaeffer, P. Feldman, and E. C. Zipf, "The Excitation of $O(^1D)$ and $O(^1S)$ Atoms in the Dawn Ionosphere", Trans. AGU, 52, 885, 1971.

71. E. J. Stone and E. C. Zipf, "Excitation of the Werner Bands of H_2 by Electron Impact", J. Chem. Phys., 56, 4646, 1972.
72. E. J. Stone and E. C. Zipf, "Electron-Impact Excitation of Nitric Oxide", J. Chem. Phys., 56, 2870, 1972.
73. W. C. Wells, W. L. Borst and E. C. Zipf, "Production of $CO(a^1\Pi)$ and Other Metastable Fragments by Electron Impact Dissociation of CO_2 ", J. Geophys. Res., 77, 69, 1972.
74. E. C. Zipf, R. Gutcheck and G. Unger, "Transport Properties of the Metastable 1S State of Oxygen", submitted for publication J. Chem. Phys., 1974.
75. D. E. Shemansky, T. M. Donahue and E. C. Zipf, " N_2 Positive and N_2^+ Band Systems and the Energy Spectra of Auroral Electrons", Planet. Space Sci., 20, 905, 1972.
76. W. C. Wells, W. L. Borst and E. C. Zipf, "Production of $CO(a^3\Pi)$ and Other Metastable Fragments by Electron Impact Dissociation of CO_2 ", J. Geophys. Res., 77, 69, 1972.
77. E. J. Stone and E. C. Zipf, "Excitation of the Werner Bands of H_2 by Electron Impact", J. Chem. Phys., 56, 4646, 1972.
78. E. J. Stone and E. C. Zipf, "Electron-Impact Excitation of Nitric Oxide", J. Chem. Phys., 56, 2870, 1972.
79. W. C. Wells, W. L. Borst and E. C. Zipf, "Translational Spectroscopy of Fast Metastables Produced by Electron Impact Dissociation of Atmospheric Gases", Bull. Am. Phys. Soc., 17, 397, 1972.
80. E. C. Zipf and E. J. Stone, "Electron Impact Excitation of Atomic Nitrogen", Bull. Am. Phys. Soc., 17, 397, 1972.
81. R. A. Gutcheck and E. C. Zipf, "Collisional Deactivation of Metastable $Ar(^3P_{0,2})$ Atoms and $N_2(^3\Sigma_u^+)$ Molecules", Bull. Am. Phys. Soc., 17, 395, (1972).
82. E. J. Stone and E. C. Zipf, "Electron Impact Excitation of the Werner Bands of H_2 ", Bull. Am. Phys. Soc., 17, 397, 1972.
83. W. C. Wells and E. C. Zipf, "Excitation of Vacuum Ultraviolet Radiation by Electron Impact on Carbon Monoxide: Some Unresolved Questions Near Threshold", J. Chem. Phys., 3583, 57, 1972.
84. R. C. Schaeffer, P. D. Feldman, and E. C. Zipf, "The Dayglow [OI] $\lambda\lambda$ 6300 and 5577 \AA Lines in the Early Morning Ionosphere", J. Geophys. Res., 77, 6828, 1972.
85. R. A. Gutcheck and E. C. Zipf, "Metastable Studies in an Afterglow: I. Quenching of $N_2(A^3\Sigma_u^+)$ ", J. Chem. Phys., 1974.

86. W. C. Wells and E. C. Zipf, "Absolute Cross Sections for the Dissociative Excitation of $\text{OI}(^3\text{S}^0)$ and its Radiative Lifetime", Trans. AGU, 53, 459, 1972.
87. M. J. Mumma, E. J. Stone, and E. C. Zipf, "Dissociative Excitation of $\text{CO}(A^1\Pi)$ by Electron Impact on CO and Implications for the Mars Upper Atmosphere", Trans. AGU, 53, 460, 1972.
88. E. C. Zipf and T. M. Donahue, "A Confirmation of Large Concentrations of Nitric Oxide in an Auroral Arc", Trans. AGU, 53, 729, 1972.
89. M. J. Mumma, E. J. Stone, W. L. Borst and E. C. Zipf, "Dissociative Excitation of Vacuum Ultraviolet Emission Features by Electron Impact on Molecular Gases III. CO_2 ", J. Chem. Phys., 57, 68, 1972.
90. W. L. Borst, W. C. Wells and E. C. Zipf, "Excitation of the Metastable $E^3\Sigma_g^+$ State of N_2 by Electron Impact", The Phys. Rev. A, 5, 1744, 1972.
91. E. J. Stone and E. C. Zipf, "Excitation of Atomic Nitrogen by Electron Impact", J. Chem. Phys., 58, 4278, 1973.
92. W. C. Wells and E. C. Zipf, "Emission Line-Shapes Produced by Electron Impact Dissociative Excitation of Atmospheric Gases", Trans. AGU, 53, 1068, 1972.
93. R. A. Gutcheck and E. C. Zipf, "Excitation of $\text{CO}(A^1\Pi)$ Molecules by the Dissociative Recombination of CO_2^+ Ions", Trans. AGU, 53, 1075, 1972.
94. E. C. Zipf, "Comment on the Role of Photoionization in Auroral Arcs", Trans. Am. Geophys. Union 54, 403, 1973.
95. E. J. Stone and E. C. Zipf, "Excitation of the $\text{OI}(^3\text{S}^0)$ and $\text{OI}(^5\text{S}^0)$ States by Electron Impact on Atomic Oxygen", submitted to J. Chem. Phys., 1973.
96. R. A. Gutcheck and E. C. Zipf, "The Excitation of the CO Fourth Positive System by the Dissociative Recombination of CO_2^+ Ions", J. Geophys. Res., 78, 5429, 1973.
97. W. C. Wells, W. L. Borst, and E. C. Zipf, "Electron Impact Excitation of Metastable States in Carbon Monoxide", Proceedings Eight International Conference on the Physics of Electronic and Atomic Collisions, Belgrad, 1973.
98. W. C. Wells, W. L. Borst, and E. C. Zipf, "Excitation of Higher Lying Metastable States in Carbon Monoxide by Electron Impact: Cross Section and Lifetime Measurements", Phys. Rev. A, 8, 2463, 1973.

99. E. C. Zipf, "Photodissociation of O_2 : A Major Source of the Auroral Green Line", Trans. AGU 54, 1155, 1973.
100. E. C. Zipf, P. W. Erdman and T. G. Finn, "A New Instrument for Mesospheric Composition Measurements", Trans. AGU 54, 1151, 1973.
101. E. C. Zipf, "On the Role of Photoionization in Auroral Arcs", submitted to J. Geophys. Res., 1973.
102. W. C. Wells and E. C. Zipf, "Lifetime of the Metastable $^5S^0$ State of Atomic Oxygen", accepted to Phys. Rev. A, 9, 568, 1974.
103. E. C. Zipf, "Comment on the Production of $N(^2D)$ Atoms by Photoabsorption Processes and by Electron-Impact Excitation of N_2 ", Trans. AGU 55, xxxx, 1974.
104. B. L. Carnahan, T. G. Finn and E. C. Zipf, "Dissociation of CH_4 by Electron Impact: Production of Metastable Hydrogen and Carbon Fragments", submitted to J. Chem. Phys., 1974.
105. G. Unger and E. C. Zipf, "Evidence for Vibrationally Excited CO_2^+ Ions in the Ionosphere of Venus", submitted to Geophys. Res. Letters, June 1974.
106. E. C. Zipf and R. McLaughlin, "On the Dissociation of N_2 by Electron Impact", submitted to Geophys. Res. Letters, June 1974.