provided by NASA Technical Reports Serve

CR-138269 -sc543.12PR

3

"Made available under NASA sponsorship in the interest of early and wide dissemination of Earth Resources Survey Program information and without liability for any use made thereor."

IDENTIFICATION AND INTERPRETATION OF TECTONIC FEATURES

ERTS-A IMAGERY

Monem Abdel-Gawad Science Center, Rockwell International Corp. 1049 Camino Dos Rios (POB 1085) Thousand Oaks, CA 91360

June 4, 1974

Type 1 Progress Report for Period April 1 through May 31, 1974

Prepared for GODDARD SPACE FLIGHT CENTER Greenbelt, Maryland 20771

by

M. addl Down

Monem Abdel-Gawad Member Technical Staff Science Center

1044A

RECEIVED JUN 0 5 1974 SIS/902.6 INTERPRETATION OF TECTONIC FEATURES FROM ERTS-1 IMAGERY Progress Report, 1 Apr.
- 31 May 1974 (Rockwell International
- 37 Conter) 2 p HC \$4.00 CSCL 086

Uncla: 3 00520

N74-26859

TYPE I PROGRESS REPORT FOR PERIOD APRIL 1 THROUGH MAY 31, 1974

TITLE: Identification and Interpretation of Tectonic Features from ERTS-1 Imagery

NASA Contract No. NAS5-21767

GSFC ID No.: PROO1, Dr. Monem Abdel-Gawad
Principal Investigator

Accomplishments

We developed a tectonic model for the development of the Central Coast Ranges in California.

Significant Results

The transverse faults observed in the central Coast Ranges of California are believed to represent the remnants of a major system of shear faults older than the San Andreas system. The transverse shear system is believed to have developed in the Mesozoic when the Pacific Plate was advancing under the North American Plate. Shear faults thus developed due to unequal rates of underthrusting. Our tectonic model indicates that the intrusive belt of the proto-Sierra Nevada and the belt of eugeosynclinal sedimentary belt (Franciscan group) which lay to the west were both subjected to regional lefthanded shear. Later development of the San Andreas system as transform faults of the East Pacific Rise changed the tectonic style to right-lateral tangential. The model explains the peculiar distribution of the Franciscan rocks in the Diablo Range east of the San Andreas fault and in Santa Lucia Range west of Nacimiento fault and the presence of Sierra Nevada type granitic blocks in between the two faults in the Salinia block.

This model is also consistent with our analysis of the Texas and Parras shears which indicates that the southwestern part of North America has been subjected to a major left-lateral regional shear before the development of the San Andreas fault system.

Published Work:

M. Abdel-Gawad and Linda Tubbesing, 1974, Regmatic transverse shear in Mexico and U. S. southwest-tectonic analysis (Abstract): First International Conference on the New Basement Tectonics: June 3-7, 1974, Salt Lake City, Utah.

Plans for Next Period

We plan to integrate the analysis of data and tectonic models obtained so far.

Practical Applications: None