

Technical University of Denmark



EFSA NDA Panel (EFSA Panel on Dietetic Products, Nutrition and Allergies), 2014. Scientific Opinion on Dietary Reference Values for zinc

EFSA Publication; Tetens, Inge

Link to article, DOI:

[10.2903/j.efsa.2014.3844](https://doi.org/10.2903/j.efsa.2014.3844)

Publication date:

2014

Document Version

Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):

EFSA Publication (2014). EFSA NDA Panel (EFSA Panel on Dietetic Products, Nutrition and Allergies), 2014. Scientific Opinion on Dietary Reference Values for zinc. Parma, Italy: European Food Safety Authority. (The EFSA Journal; No. 3844, Vol. 12(10)). DOI: 10.2903/j.efsa.2014.3844

DTU Library

Technical Information Center of Denmark

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

SCIENTIFIC OPINION

Scientific Opinion on Dietary Reference Values for zinc¹

EFSA Panel on Dietetic Products, Nutrition and Allergies (NDA)^{2,3}

European Food Safety Authority (EFSA), Parma, Italy

ABSTRACT

Following a request from the European Commission, the Panel on Dietetic Products, Nutrition and Allergies (NDA) derived Dietary Reference Values for zinc, using a two-stage factorial approach and reference values for body weight. The first stage of estimating physiological requirements used studies that had physiologically plausible data, specifically related to faecal excretion of endogenous zinc. Adult physiological requirements were closely related to body size, and sex differences were not detectable after adjustment for body weight. Average Requirements (ARs) for dietary zinc necessary to meet physiological requirements were estimated using saturation response modelling, taking into account the inhibitory effect of dietary phytate on zinc absorption. Estimated ARs and Population Reference Intakes (PRIs) are provided for phytate intake levels of 300, 600, 900 and 1 200 mg/day, which cover the range of mean/median intakes observed in European populations. ARs range from 6.2 to 10.2 mg/day for women with a reference weight of 58.5 kg and from 7.5 to 12.7 mg/day for men with a reference weight of 68.1 kg. PRIs were derived from the zinc requirement of individuals with a body weight at the 97.5th percentile for reference weights for men and women and range from 7.5 to 12.7 mg/day for women and from 9.4 to 16.3 mg/day for men. ARs for infants from seven months of age and for children were estimated factorially, based on extrapolation from estimates of adult losses plus zinc needs for growth, and range from 2.4 to 11.8 mg/day. PRIs for infants and children were derived by assuming a coefficient of variation of 10 %, and range from 2.9 to 14.2 mg/day. For pregnancy and lactation, additional zinc requirements related to fetal and maternal tissues and transfer of zinc into breast milk, respectively, were considered and additional PRIs of 1.6 and 2.9 mg/day, respectively, were estimated.

© European Food Safety Authority, 2014

KEY WORDS

zinc, Dietary Reference Value, Population Reference Intake, phytate

¹ On request from the European Commission, Question No EFSA-Q-2011-01233, adopted on 19 September 2014.

² Panel members: Carlo Agostoni, Roberto Berni Canani, Susan Fairweather-Tait, Marina Heinonen, Hannu Korhonen, Sébastien La Vieille, Rosangela Marchelli, Ambroise Martin, Androniki Naska, Monika Neuhäuser-Berthold, Grażyna Nowicka, Yolanda Sanz, Alfonso Siani, Anders Sjödin, Martin Stern, Sean (J.J.) Strain, Inge Tetens, Daniel Tomé, Dominique Turck and Hans Verhagen. Correspondence: nda@efsa.europa.eu

³ Acknowledgement: The Panel wishes to thank the members of the Working Group on Dietary Reference Values for minerals: Peter Aggett, Carlo Agostoni, Susan Fairweather-Tait, Marianne Geleijnse, Michael Hambidge, Ambroise Martin, Androniki Naska, Hildegard Przyrembel and Alfonso Siani for the preparatory work on this scientific opinion, the hearing expert: Leland Miller, and EFSA staff: Anja Brönstrup, Fanny Héraud, Sofia Ioannidou, Laura Martino and Liisa Valsta for the support provided to this scientific opinion.

Suggested citation: EFSA NDA Panel (EFSA Panel on Dietetic Products, Nutrition and Allergies), 2014. Scientific Opinion on Dietary Reference Values for zinc. EFSA Journal 2014;12(10):3844, 76 pp. doi:10.2903/j.efsa.2014.3844

Available online: www.efsa.europa.eu/efsajournal

SUMMARY

Following a request from the European Commission, the EFSA Panel on Dietetic Products, Nutrition and Allergies (NDA) was asked to deliver a scientific opinion on Dietary Reference Values (DRVs) for the European population, including zinc.

Zinc has a wide array of vital physiological functions. It has a catalytic role in each of the six classes of enzymes. The human transcriptome has 2 500 zinc finger proteins, which have a broad intracellular distribution and the activities of which include binding of RNA molecules and involvement in protein–protein interactions. Thus, their biological roles include transcriptional and translational control/modulation and signal transduction.

The majority of dietary zinc is absorbed in the upper small intestine. The luminal contents of the duodenum and jejunum, notably phytate, can have a major impact on the percentage of zinc that is available for absorption. Absorption of zinc by the enterocyte is regulated in response to the quantity of bioavailable zinc ingested. Albumin is the major transporter of zinc in both portal and systemic circulation. Virtually no zinc circulates in a free ionised form, and the majority of total body zinc is in muscle and bone; zinc does not have an identified major storage site. The quantity of zinc secreted into and excreted from the intestinal tract depends on body zinc concentrations, and the quantities of endogenous zinc in the faeces and exogenous zinc absorbed in normal adults are related. The kidneys and integument are minor routes of loss of endogenous zinc.

Plasma/serum zinc concentration and other putative biomarkers of zinc adequacy, deficiency and excess are not useful for estimating DRVs for zinc. Zinc requirements have been estimated by the factorial approach involving two stages. The first is the estimation of physiological requirements, defined as the minimum quantity of absorbed zinc needed to match losses of endogenous zinc and to meet any additional requirements for absorbed zinc that may be necessary for growth in healthy well-nourished infants and children, and in pregnancy and lactation. The second stage is the determination of the quantity of dietary zinc available for absorption that is needed to meet these physiological requirements. From the published literature, 15 studies were identified that included data on endogenous faecal zinc and total absorbed zinc that enabled an estimation to be made of the physiological zinc requirements of adults. Individual's data from these studies were supplied by the authors. Data were assessed for physiological plausibility and, after careful evaluation, some data were excluded from further calculations. The final numbers of subjects contributing data to the estimate of physiological zinc requirements were 31 males and 54 females, from a total of 10 studies. Dietary phytate intakes were available for some of the included studies, either as mean study values or as individual's data. The range of dietary phytate intakes in the available data was 0–2 080 mg/day. Multiple regression analysis was used to evaluate the possible relationships between physiological requirements and sex, zinc balance (difference between absorbed zinc and total losses of endogenous zinc) and body size. The coefficient of determination (R^2) values for the models with body weight, height, body mass index and body surface area variables were 0.46, 0.42, 0.37 and 0.47, respectively. It was decided to use the equation relating physiological requirement to body weight in further analyses, for reasons of convenience and accuracy of measurement. The equation for physiological requirement was calculated on the basis that physiological requirement is equivalent to total absorbed zinc when absorbed zinc minus total endogenous zinc losses equals zero at a given body weight. For deriving the dietary zinc requirement, a trivariate saturation response model of the relationship between zinc absorption, and dietary zinc and phytate was established using 72 mean datasets (reflecting 650 individual measurements) reported in 18 publications. The R^2 of the fit of this model was 0.81. From this model, the Average Requirement (AR) was determined as the intercept of the total absorbed zinc needed to meet physiological requirements. Estimated ARs and Population Reference Intakes (PRIs) for zinc are provided for phytate intake levels of 300, 600, 900 and 1 200 mg/day, which cover the range of mean/median phytate intakes observed in European populations. ARs range from 6.2 to 10.2 mg/day for women with a reference body weight of 58.5 kg and from 7.5 to 12.7 mg/day for men with a reference body weight of 68.1 kg. PRIs for adults were estimated as the zinc requirement of individuals with a body weight at the 97.5th percentile for reference body weights

for men and women, respectively, and range from 7.5 to 12.7 mg/day for women and from 9.4 to 16.3 mg/day for men.

For infants from seven months of age and children, DRVs for zinc were derived using the factorial approach, taking into account endogenous zinc losses via urine, sweat and integument, faeces and, in adolescent boys and girls, semen and menses, respectively, as well as zinc required for synthesis of new tissue for growth. Urinary and integumental losses were extrapolated based on estimates of adult losses, whereas endogenous faecal zinc losses were estimated by linear regression analysis of endogenous faecal zinc losses versus body weight for the subjects contributing data to the adult estimates, and for infants and young children from two studies from China and the USA. Zinc requirements for growth were taken into account based on the zinc content of new tissue, and by estimating daily weight gains for each age group. Absorption efficiency of zinc from mixed diets was assumed to be 30%. Estimated ARs range from 2.4 mg/day in infants aged 7–11 months to 11.8 mg/day in adolescent boys. Owing to the absence of reference body weights for infants and children at the 97.5th percentile, and in the absence of knowledge about the variation in requirements, PRIs for infants and children were estimated based on a coefficient of variation (CV) of 10%, and range from 2.9 to 14.2 mg/day.

The physiological requirements for pregnancy and lactation can be calculated by adding the increases in physiological requirements that are predicted to meet the demands for new tissue primarily of the conceptus, and the replacement of zinc that is secreted in breast milk. For pregnancy, an additional requirement for zinc for the four quarters of pregnancy of about 0.4 mg/day was assumed because of zinc accumulation in the fetus; placental, uterine and mammary tissue; amniotic fluid and maternal blood. The Panel decided not to use the trivariate model to estimate the dietary zinc intake required to meet the additional physiological requirement. Instead, the Panel applied a mean fractional absorption of zinc of 0.3 that has been observed in healthy adults to the physiological requirement of 0.4 mg/day. The additional requirement for pregnant women was calculated to be 1.3 mg/day and the additional PRI for pregnancy was estimated based on a CV of 10% and was 1.6 mg/day.

For lactation, taking into account breast milk zinc concentration, the breast milk volume transferred and the postnatal redistribution of zinc owing to involution of the uterus and reduction of maternal blood volume, the additional physiological requirement calculated over six months of lactation was estimated to be 1.1 mg/day. Assuming that fractional absorption of zinc is increased 1.5-fold in lactation, and applying a fractional absorption of zinc of 0.45 to the additional physiological requirement of 1.1 mg/day, resulted in an additional dietary requirement for lactating women of 2.4 mg/day. The additional PRI for lactation, based on a CV of 10%, was 2.9 mg/day.

Meat, legumes, eggs, fish, and grains and grain-based products are rich dietary zinc sources. On the basis of data from 12 dietary surveys in nine European Union (EU) countries, zinc intake was assessed using food consumption data from the EFSA Comprehensive Food Consumption Database and zinc composition data from the EFSA nutrient composition database. Average zinc intake ranged from 4.6 to 6.2 mg/day in children aged one to less than three years, from 5.5 to 9.3 mg/day in children aged 3 to < 10 years, from 6.8 to 14.5 mg/day in adolescents (10 to < 18 years) and from 8.0 and 14.0 mg/day in adults. The main food groups contributing to zinc intake were meat and meat products, grains and grain-based products, and milk and dairy products. Published data on phytate intake in the EU are limited and indicate a wide range of dietary phytate intakes.

TABLE OF CONTENTS

Abstract	1
Summary	2
Table of contents	4
Background as provided by the European Commission.....	6
Terms of reference as provided by the European Commission.....	6
Assessment	8
1. Introduction	8
2. Definition/category	8
2.1. Chemistry	8
2.2. Functions of zinc	8
2.2.1. Biochemical functions	8
2.2.2. Health consequences of deficiency and excess	9
2.2.2.1. Deficiency	9
2.2.2.2. Excess	9
2.3. Physiology and metabolism	9
2.3.1. Intestinal absorption	10
2.3.2. Transport in blood	11
2.3.3. Distribution to tissues	11
2.3.4. Storage	11
2.3.5. Metabolism	11
2.3.6. Elimination	11
2.3.6.1. Faeces.....	11
2.3.6.2. Urine and sweat	11
2.3.6.3. Breast milk.....	11
2.3.7. Interaction with other nutrients.....	12
2.3.8. Biomarkers	12
2.3.9. Plasma zinc concentration	12
2.3.10. Hair zinc concentration.....	12
2.3.11. Urinary zinc concentration	12
2.3.12. Other biomarkers	13
2.3.13. Conclusion on biomarkers	13
2.4. Effects of genotype	13
3. Dietary sources and intake data	13
3.1. Dietary sources.....	13
3.2. Dietary zinc intake	13
3.3. Dietary phytate intake	15
4. Overview of Dietary Reference Values and recommendations.....	15
4.1. Adults.....	15
4.2. Infants and children.....	17
4.3. Pregnancy.....	19
4.4. Lactation	20
5. Criteria (endpoints) on which to base Dietary Reference Values.....	22
5.1. Indicators of adult zinc requirement	22
5.1.1. Physiological requirement	22
5.1.1.1. Identification of studies, data extraction and assessment of methodological quality.....	22
5.1.1.2. Inclusion of studies	22
5.1.1.3. Inclusion of individual data	23
5.1.1.4. Estimation of endogenous zinc losses.....	23
5.1.1.5. Modelling of zinc requirements.....	23
5.1.2. Estimation of dietary zinc intake to meet physiological requirement	25
5.1.2.1. Effect of dietary zinc and phytate on absorbed zinc	25
5.1.2.2. Identification of studies, data extraction and assessment of methodological quality.....	25
5.1.2.3. Modelling of the saturation response model	26

5.2.	Indicators of zinc requirements of children	28
5.3.	Indicators of zinc requirements in pregnancy and lactation.....	28
5.4.	Zinc intake and long-term health consequences	29
6.	Data on which to base Dietary Reference Values.....	29
6.1.	Adults.....	29
6.2.	Infants and children.....	30
6.2.1.	Methodology.....	30
6.2.1.1.	Urinary and integumental zinc losses	30
6.2.1.2.	Endogenous faecal zinc losses	31
6.2.1.3.	Zinc losses in menses and semen.....	31
6.2.1.4.	Zinc requirement for growth.....	31
6.2.1.5.	Fractional absorption of zinc	31
6.2.2.	Infants aged 7 to 11 months.....	31
6.2.3.	Children	32
6.3.	Pregnancy and lactation	34
	Conclusions	35
	References	36
	Appendices	48
Appendix A.	Concentrations of zinc in breast milk from mothers of term infants in Europe	48
Appendix B.	Dietary surveys in the EFSA Comprehensive database update dataset included in the nutrient intake calculation and number of subjects in the different age classes.....	53
Appendix C.	Zinc intake in males in different surveys according to age classes and country	54
Appendix D.	Zinc intake in females in different surveys according to age classes and country	56
Appendix E.	Minimum and maximum percentage contribution of different FoodEx2 level 1 food groups to zinc intake in males	58
Appendix F.	Minimum and maximum percentage contribution of different FoodEx2 level 1 food groups to zinc intake in females.....	59
Appendix G.	Phytate/Phytic acid intake in various European countries.....	60
Appendix H.	Evaluating data when endogenous faecal zinc was estimated using the zinc absorption–intestinal balance method	63
Appendix I.	Data extracted from the selected studies for estimating physiological zinc requirement of adults	67
Appendix J.	Data regression analysis diagnostic results	69
Appendix K.	Data extracted from the selected studies for the trivariate saturation response model.....	72
Appendix L.	Three-dimensional representation of Figure 1.....	74
	Abbreviations	75

BACKGROUND AS PROVIDED BY THE EUROPEAN COMMISSION

Scientific advice on nutrient intakes is important as the basis of Community action in the field of nutrition, for example such advice has in the past been used as the basis of nutrition labelling. The Scientific Committee for Food (SCF) report on nutrient and energy intakes for the European Community dates from 1993. There is a need to review and if necessary to update these earlier recommendations to ensure that the Community action in the area of nutrition is underpinned by the latest scientific advice.

In 1993, the SCF adopted an opinion on nutrient and energy intakes for the European Community.⁴ The report provided Reference Intakes for energy, certain macronutrients and micronutrients, but it did not include certain substances of physiological importance, for example dietary fibre.

Since then, new scientific data have become available for some of the nutrients, and scientific advisory bodies in many European Union Member States and in the United States have reported on recommended dietary intakes. For a number of nutrients these newly established (national) recommendations differ from the reference intakes in the SCF (1993) report. Although there is considerable consensus between these newly derived (national) recommendations, differing opinions remain on some of the recommendations. Therefore, there is a need to review the existing EU Reference Intakes in the light of new scientific evidence, and taking into account the more recently reported national recommendations. There is also a need to include dietary components that were not covered in the SCF opinion of 1993, such as dietary fibre, and to consider whether it might be appropriate to establish reference intakes for other (essential) substances with a physiological effect.

In this context, EFSA is requested to consider the existing Population Reference Intakes for energy, micro- and macronutrients and certain other dietary components, to review and complete the SCF recommendations, in the light of new evidence, and in addition advise on a Population Reference Intake for dietary fibre.

For communication of nutrition and healthy eating messages to the public it is generally more appropriate to express recommendations for the intake of individual nutrients or substances in food-based terms. In this context, EFSA is asked to provide assistance on the translation of nutrient based recommendations for a healthy diet into food based recommendations intended for the population as a whole.

TERMS OF REFERENCE AS PROVIDED BY THE EUROPEAN COMMISSION

In accordance with Article 29 (1)(a) and Article 31 of Regulation (EC) No. 178/2002,⁵ the Commission requests EFSA to review the existing advice of the Scientific Committee for Food on population reference intakes for energy, nutrients and other substances with a nutritional or physiological effect in the context of a balanced diet which, when part of an overall healthy lifestyle, contribute to good health through optimal nutrition.

In the first instance, EFSA is asked to provide advice on energy, macronutrients and dietary fibre. Specifically, advice is requested on the following dietary components:

- Carbohydrates, including sugars;
- Fats, including saturated fatty acids, polyunsaturated fatty acids and monounsaturated fatty acids, *trans* fatty acids;

⁴ Scientific Committee for Food. Nutrient and energy intakes for the European Community. Reports of the Scientific Committee for Food, 31st series. Office for Official Publication of the European Communities, Luxembourg, 1993.

⁵ Regulation (EC) No 178/2002 of the European Parliament and of the Council of 28 January 2002 laying down the general principles and requirements of food law, establishing the European Food Safety Authority and laying down procedures in matters of food safety. OJ L 31, 1.2.2002, p. 1–24.

- Protein;
- Dietary fibre.

Following on from the first part of the task, EFSA is asked to advise on population reference intakes of micronutrients in the diet and, if considered appropriate, other essential substances with a nutritional or physiological effect in the context of a balanced diet which, when part of an overall healthy lifestyle, contribute to good health through optimal nutrition.

Finally, EFSA is asked to provide guidance on the translation of nutrient based dietary advice into guidance, intended for the European population as a whole, on the contribution of different foods or categories of foods to an overall diet that would help to maintain good health through optimal nutrition (food-based dietary guidelines).

ASSESSMENT

1. Introduction

In 1993, the Scientific Committee for Food (SCF) published an opinion on nutrient and energy intakes for the European Community (SCF, 1993). For zinc (Zn), Population Reference Intakes (PRIs) were proposed for all population groups from seven months of age onwards, based on zinc requirements to replace basal losses and losses via breast milk in lactating women, or an increment to supply zinc for growth in children and pregnant women. In addition, a Lowest Threshold Intake was derived for men and women (see Section 4).

2. Definition/category

2.1. Chemistry

Zinc has an atomic mass of 65.39 Da and is the 24th most abundant element in the Earth's crust. It exists as a stable divalent cation. Considered to be of fundamental importance to the far-ranging biology of zinc is its ability for fast exchange coupled with strong binding to organic molecules, especially to thiolate and amine electron donors. Zinc does not exhibit direct redox activity, a feature which facilitates its safe transport within the body (Krezel et al., 2007). There are five naturally occurring stable isotopes of zinc; the most abundant is ⁶⁴Zn (48.63 % natural abundance).

2.2. Functions of zinc

2.2.1. Biochemical functions

Zinc has a wide array of vital physiological functions and is ubiquitous within every cell in the body. It is this very abundance that is thought to be the reason why it has proved so challenging to link zinc deficiency with specific phenotypic features. However, three general functional classes (catalytic, structural and regulatory) define zinc's role in biology (King and Cousins, 2014). Zinc has a structural or catalytic role, or both, in each of the six classes of enzymes, although unequivocal evidence of a direct link between signs of zinc deficiency and a deficiency of a specific metallo-enzyme has not yet been confirmed in humans.

The structural role of zinc is exemplified by transcription factors having zinc motifs (zinc fingers) which link with cysteine and histidine to form a tetrahedral Zn²⁺ coordination complex. The presence of zinc is necessary for the activity of these zinc fingers. The human transcriptome has 2 500 zinc finger proteins, which represent 8 % of the genome and account for a significant portion of the zinc requirement (King and Cousins, 2014). Zinc fingers have a range of binding affinities, suggesting that some zinc finger-dependent transcription may be especially vulnerable to low zinc absorption. Zinc finger proteins have a broad intracellular distribution and their activities include binding of RNA molecules and involvement in protein-protein interactions. Thus, their biological roles include transcriptional and translational control/modulation and signal transduction. A combination of structural and regulatory functions are involved in the large quantities of zinc movement involved in the release of insulin, the secretion of zinc-containing digestive enzymes and acid secretion by parietal cells in the stomach (Guo et al., 2010).

Regulation of gene expression is a key biochemical role of zinc. The metal-response element (MRE)-binding transcription factor (MTF1) is thought to provide zinc responsiveness to many genes (King and Cousins, 2014), including a master regulatory role for micro RNA genes involved in gene expression.

A second regulatory role of zinc is as a regulator of intracellular signalling, analogous to calcium but at a finer level of control, in particular through the regulation of kinase and phosphorylase activity (King and Cousins, 2014). The control of phosphorylation/dephosphorylation may explain the effects of zinc on phosphorylated transcription factors, the effects of zinc on cell surface receptor binding of

growth factors and cytokine receptors, and the major effects of zinc on virtually all aspects of the immune system.

2.2.2. Health consequences of deficiency and excess

2.2.2.1. Deficiency

There is a lack of specific health effects of zinc deficiency, apart from those observed in infants with acrodermatitis enteropathica (see below), and this is the consequence of its essentiality for many core biochemical processes. There is protection by homeostatic mechanisms at both a whole body and a tissue level, which, during periods of rapid growth, include slowing of linear growth (i.e. bone growth). Although by no means unique to zinc deficiency, slowing of linear growth is one of the most clearly defined effects of chronic zinc deficiency. The particular vulnerability of the immune system to zinc deficiency results in part from its high rate of cell proliferation. However, the immune system also epitomises the dependence of many cellular biochemical processes on zinc. These may include atypical regulation of cytokine gene expression and signalling pathways, which can disrupt the balance of cell-mediated versus humoral immunity. The failure of zinc-dependent structural factors needed for antigen presentation may enhance the risk of microbial and parasitic infections (King and Cousins, 2014), of which enteric infections have been the principal foci of interest (Black, 2003).

Acute severe zinc deficiency results from genetic defects in zinc transporters involved in the intestinal absorption of zinc and in the transfer of zinc by the mammary gland into human milk, collectively termed acrodermatitis enteropathica. The onset of clinical features after birth is rapid. The most superficially apparent are skin lesions, which are, characteristically, most prominent around the body orifices and on the extremities. Diarrhoea is prominent in most but not all cases. Growth failure is progressive and these infants are susceptible to a range of immune defects and infections. Loss of appetite and of taste perception are notable, and alterations in affect and mood are early phenomena of incipient zinc deficiency in children with acrodermatitis enteropathica when their supplemental zinc becomes inadequate. Response to treatment with zinc is profound, but without treatment there is typically a fatal outcome in acrodermatitis enteropathica by later infancy. Similar acute acquired zinc deficiency states have been extensively documented, primarily in patients dependent on intravenous nutrition lacking zinc (Younoszai, 1983).

2.2.2.2. Excess

Chronic high zinc intake can result in severe neurological diseases attributable to copper deficiency (Hedera et al., 2009). The SCF (2002) has set a Tolerable Upper Intake Level (UL) of 25 mg/day for adults, including pregnant and lactating women, based on studies of zinc supplementation for up to 14 weeks. A No Observed Adverse Effect Level of 50 mg/day was based on the absence of any adverse effect on a wide range of relevant indicators of copper status in controlled metabolic studies. An Uncertainty Factor of 2 was applied. The UL for children was extrapolated from the UL for adults using body weight to the power of 0.75 and reference body weights for European children (SCF, 1993).

2.3. Physiology and metabolism

Zinc transporter gene regulation currently dominates all aspects of cellular zinc metabolism. The ZnT family (SLC30a) facilitates the efflux of zinc across cell membranes and into vesicles. The ZIP transporters do the reverse. Up- or down-regulation of these genes in response to zinc intake contributes to the tight homeostatic control of zinc by the small intestine. Diet is among the factors that regulate transporter gene expression. These same families of transporters have the major role of regulating uptake, excretion and metabolism of zinc by all cells in the body. Metallothionein also has a supportive role in zinc metabolism. Polymorphisms in these genes can affect phenotypic expression.

2.3.1. Intestinal absorption

Small quantities of zinc may be absorbed throughout the entire gastro-intestinal tract, but the majority is absorbed in the upper small intestine. When ingested from food, it will be firmly bound, particularly to protein thiols and nitrogen ligands. The phytate–zinc ligand is weakened at low pH (Cheryan, 1980) and the results of stable isotope studies of zinc absorption are consistent with the zinc being released from these ligands and entering a common pool in the acidic environment of the stomach and, subsequently, being bound to a variety of other organic ligands, including phytate in the alkaline medium of the distal duodenum. The form in which bioavailable zinc is presented to the apical surface of the enterocyte and the zinc transporters, particularly Zip 4, has not been fully elucidated. The luminal contents of the duodenum and jejunum in particular, especially phytate, can have a major impact on the percentage of zinc available for absorption. With diets low in phytate and low in zinc, for example less than 4 mg/day, the fraction of zinc absorbed may be as high as 60 % or more. The fraction of absorbed zinc then decreases progressively with increasing dietary zinc (Hambidge et al., 2005). The uptake of zinc and its transfer into the body by the enterocyte is regulated in response to the quantity of bioavailable zinc ingested (Chung et al., 2008); this relationship between the quantity of zinc absorbed and that ingested is best fit with saturation response modelling (Hambidge et al., 2010).

WHO/FAO (2004) categorised diets with regard to the fact that their impact on zinc absorption is mainly influenced by the phytate–zinc molar ratio and the amount and source of dietary protein. In most European countries, the main contributors to dietary protein intake of adults are meat and meat products, followed by grains and grain-based products, and milk and dairy products. The mean protein intake of European adults is generally above the Average Requirement (AR) (EFSA NDA Panel, 2012). Thus, for the majority of the European population that consumes mixed diets, the Panel considers that the phytate content of the diet has a more profound effect on zinc availability than the protein content, and that at zinc intake adequate to meet the requirement the absorption efficiency of zinc from the diet is moderate or high (Table 1).

Table 1: Criteria for categorising diets according to their potential absorption efficiency of zinc (adapted from WHO/FAO (2004))

Absorption efficiency	Principal dietary characteristics
High	Refined diets low in cereal fibre, low in phytic acid content and with a phytate–zinc molar ratio below 5; adequate protein content principally from non-vegetable sources, such as meat and fish. At a zinc intake of 10 mg/day, a phytate–zinc molar ratio of below 5 is equivalent to a phytate intake of below about 500 mg/day
Moderate	Mixed diets containing animal or fish protein (Lacto-)ovo-vegetarian or vegan diets not based primarily on unrefined cereal grains or high-extraction-rate flours A phytate–zinc molar ratio of total diet within the range 5–15, or not exceeding 10 if more than 50 % of the energy intake is accounted for by unfermented, unrefined cereal grains and flours At a zinc intake of 10 mg/day, a phytate–zinc molar ratio of 5–15 is equivalent to a phytate intake of about 500–1 500 mg/day
Low	Diets high in unrefined, unfermented and ungerminated cereal grain ^(a) , especially when intake of animal protein is negligible Phytate–zinc molar ratio of total diet exceeds 15; high-phytate soya-protein products constitute the primary protein source Diets in which, singly or collectively, approximately 50 % of the energy intake is accounted for by the following high-phytate foods: high-extraction-rate (≥ 90 %) wheat, rice, maize, grains and flours, oatmeal, and millet; sorghum, cowpeas, pigeon peas, grams, kidney beans, black-eyed beans and groundnut flours At a zinc intake of 10 mg/day, a phytate–zinc molar ratio exceeding 15 is equivalent to a phytate intake higher than about 1 500 mg/day

(a): Germination of cereal grains or fermentation (e.g. leavening) of many flours can reduce antagonistic potency of phytates; if done, the diet should then be classified as having a moderate absorption efficiency of zinc.

2.3.2. Transport in blood

Albumin is the major transporter of zinc in both portal and systemic circulation. Virtually no zinc circulates unbound. Zinc in the plasma compartment is turned over more than 130 times per day and 80 % of circulating zinc is in the cellular components of the blood.

2.3.3. Distribution to tissues

Total body zinc in adult males is approximately 2.5 g in men and 1.5 g in women. The majority of total body zinc, i.e. about 85 %, is in muscle and bone. There are metabolic pools with both short- and long-term turnover. The exchangeable zinc pool exchanges with plasma zinc in approximately two days and is thought to represent the most metabolically active portion of total body zinc.

Zinc uptake capacity by the human placenta is inversely related to maternal plasma zinc concentrations and increases with increasing gestational age. There are no recent data on the metabolism of zinc by the placenta and fetus at the molecular level.

2.3.4. Storage

Zinc does not have an identified major storage site. The liver provides a limited short-term store of zinc, which is readily released as needed. Twenty per cent of bone zinc, which accounts for about 30 % of total body zinc, has been reported to be released into the circulation in times of depletion at a slower rate than liver zinc. At times of increased bone turnover and tissue catabolism, zinc is released adventitiously from these depots. Although muscle has the largest quantity of zinc, release of this zinc in response to zinc depletion has not been documented. Within all cells, vesicles provide sites for temporary storage.

2.3.5. Metabolism

The rapid turnover of plasma zinc reflects its exchange with all tissues and organs in the body. There is a rapidly exchanging pool of zinc that fully exchanges with zinc in plasma and accounts for about 10 % of total body zinc. This zinc is found in soft tissues other than muscle, particularly in the liver (Wastney et al., 1986; Miller et al., 2000).

2.3.6. Elimination

2.3.6.1. Faeces

The quantity of zinc secreted into and excreted from the gastro-intestinal tract depends on zinc intake and status. The amount of endogenous zinc in the faeces and the quantity of exogenous zinc absorbed in normal adults is positively related.

2.3.6.2. Urine and sweat

The kidneys and integument are relatively minor routes of excretion of endogenous zinc. There is a weak positive relationship between absorbed zinc and urinary zinc. However, the latter declines markedly when dietary zinc is severely reduced. For subjects with a normal zinc status, urinary zinc losses of 0.5 mg/day for men and 0.3 mg/day for women have been calculated based on individual data from studies by Jackson et al. (1984); Turnlund et al. (1984); Lowe et al. (1997); Miller et al. (2000); King et al. (2001); Pinna et al. (2001); Sheng et al. (2009) (see Section 5.1.1). Studies of whole body surface zinc losses in men have indicated combined integumental and sweat zinc losses of 0.5 mg/day for men (Jacob et al., 1981; Milne et al., 1983; Johnson et al., 1993).

2.3.6.3. Breast milk

During lactation, the quantity of zinc transferred from the mammary gland to the exclusively (or partially) breast-fed infant decreases, and this physiological decline is quite notable. Milk zinc concentrations do not appear to be associated with maternal zinc status or dietary zinc intake (Mills, 1989), and long-term ingestion of supplementary zinc (15 mg/day from two weeks post partum until

seven months) did not affect the rate of decline of milk zinc concentration in supplemented women (Krebs et al., 1995). A comprehensive review of breast milk zinc concentrations and zinc transferred from mother to child covered 63 studies globally, including 12 studies from European countries (Brown et al., 2009). Zinc concentrations (mean \pm SD) were 4.11 ± 1.50 mg/L below one month ($n = 74$ observations), 1.91 ± 0.53 mg/L at one to two months ($n = 42$), 0.98 ± 0.35 mg/L at three to five months ($n = 24$) and 0.77 ± 0.22 mg/L at 6–11 months post partum ($n = 24$). Taking into account breast milk volume, Brown et al. (2009) estimated a milk zinc transfer of 2.52 mg/day for the first month, 1.37 mg/day for months 1 to 2 and 0.86 mg/day for months 3 to < 6.

Additional data on breast milk zinc concentrations in mothers of term infants in Europe are given in Appendix A.

2.3.7. Interaction with other nutrients

High-dose iron supplements can interfere with zinc absorption when provided simultaneously with zinc supplements. There is no interference with zinc absorption from iron added to foods.

High doses of zinc can interfere with copper absorption (see Section 2.2.2.2).

2.3.8. Biomarkers

A systematic review and meta-analysis of the literature examining the efficacy of potential biomarkers of zinc status was undertaken by Lowe et al. (2009). This review presented an analysis of data from more than 32 potential biomarkers; however, for many biomarkers, there was insufficient evidence to assess their reliability.

2.3.9. Plasma zinc concentration

In apparently healthy subjects, plasma and serum zinc concentration is affected by intake, both inadequate and excessive. Lowe et al. (2009) concluded that plasma zinc concentration responds to an increase in intake over short periods, but that the homeostatic mechanisms that act to maintain plasma zinc concentration within the physiological range may prevent high plasma concentrations from being sustained over a prolonged period.

Plasma zinc concentrations are reduced in severe inherited and acquired zinc deficiency states (Wessells et al., 2014). However, as a biomarker, sensitivity is poor and, with more moderate zinc deficiency states, lacks specificity (King, 2011). Plasma zinc concentration has been recommended as a biomarker of zinc status and of the population's risk of zinc deficiency by the World Health Organization (WHO), the United Nations Children's Fund (UNICEF), the International Atomic Energy Agency (IAEA) and the International Zinc Nutrition Consultative Group (IZiNCG) (de Benoist et al., 2007).

2.3.10. Hair zinc concentration

Low hair zinc concentrations have been associated with retarded growth (Gibson et al., 1989; Gibson et al., 1991). However, there are potential and actual confounders which may have a role in apparent age-related differences in hair zinc concentrations (Hambidge et al., 1972). Based on three randomised controlled trials with a zinc intake between 15 and 100 mg/day, Lowe et al. (2009) concluded that hair zinc concentration increases in response to an increase in zinc intake, but that the effect of zinc depletion is inconclusive.

2.3.11. Urinary zinc concentration

Urinary zinc concentration has been found to increase in response to increases in zinc intake resulting from zinc supplementation; however, the response to zinc depletion has been reported to be inconclusive (Lowe et al., 2009).

2.3.12. Other biomarkers

It was previously hypothesised that there may be biomarkers based on zinc transporters or metallothionein. However, this has not been confirmed. Candidates based on proteomic and metabolomic techniques are of current research interest (Kettunen et al., 2012; Ryu et al., 2012); however, the Panel considers that they are not yet useful for deriving Dietary Reference Values (DRVs).

2.3.13. Conclusion on biomarkers

The Panel considers that neither plasma/serum zinc concentration nor any other putative biomarker is useful for estimating DRVs for zinc.

2.4. Effects of genotype

The most well-documented and severe polymorphisms for zinc result in the clinical syndrome of acrodermatitis enteropathica (Zip 4) and in the “lethal mouse syndrome (ZT4)”. A similar defect in cattle (Adema disease) is less well characterised. There are no known genotypes that would affect the estimation of DRVs for zinc. Significant results for altered putative zinc biomarkers in groups of people with differing gene variants have been documented by Lowe et al. (2013).

3. Dietary sources and intake data

3.1. Dietary sources

Meat, legumes, eggs, fish, and grains and grain-based products are rich dietary zinc sources.

Currently, zinc acetate, zinc bisglycinate, zinc chloride, zinc citrate, zinc gluconate, zinc lactate, zinc oxide, zinc carbonate and zinc sulphate may be added to both foods⁶ and food supplements.⁷ Zinc L-ascorbate, zinc L-aspartate, zinc L-lysinate, zinc malate, zinc mono-L-methionine sulphate, zinc L-pidolate and zinc picolinate may be added to food supplements only. The zinc content content of infant and follow-on formulae⁸ and processed cereal-based foods and baby foods for infants and young children⁹ is regulated.

3.2. Dietary zinc intake

Dietary intake of zinc was estimated by the Evidence Management Unit (DATA) of EFSA using the EFSA Comprehensive Food Consumption Database (EFSA, 2011b) and the EFSA Food Composition Database. This assessment includes food consumption data from 12 dietary surveys (Appendix B) from nine countries (Finland, France, Germany, Ireland, Italy, Latvia, the Netherlands, Sweden and the UK), which were either already classified or re-classified (French and Italian data), in accordance with the FoodEx2 food classification system (EFSA, 2011a). The data covered all age groups from infants to adults aged 75 years or older. The EFSA Food Composition Database was compiled during a procurement project (Roe et al., 2013) involving 14 national food database compiler organisations, who were allowed to borrow compatible data from other countries if no original composition data were available. This assessment includes food composition information from Finland, France, Germany, Italy, the Netherlands, Sweden and the UK. The amount of borrowed zinc values in these datasets varied between 14 and 88 %. For Ireland and Latvia, UK and German food composition data were used, respectively, because no composition data from these countries were available. Zinc concentration was directly available for 2 063 food terms of the food consumption data used in this

⁶ Regulation (EC) No 1925/2006 of the European Parliament and of the Council of 20 December 2006 on the addition of vitamins and minerals and of certain other substances to foods. OJ L 404, 30.12.2006, p. 26.

⁷ Directive 2002/46/EC of the European Parliament and of the Council of 10 June 2002 on the approximation of the laws of the Member States relating to food supplements. OJ L 183, 12.7.2002, p. 51.

⁸ Commission Directive 2006/141/EC of 22 December 2006 on infant formulae and follow-on formulae and amending Directive 1999/21/EC, OJ L 401, 30.12.2006, p. 1.

⁹ Commission Directive 2006/125/EC of 5 December 2006 on processed cereal-based foods and baby foods for infants and young children, OJ L 339, 06.12.2006, p. 16.

assessment, and was missing for all included countries for 599 consumed food items, to which either a value from another food (if the food with the missing value was consumed frequently or in high quantities or belonged to a food group with a high zinc concentration) or a zero value (otherwise) was attributed.

After consistency checks and replacement of missing values for zinc in the EFSA Food Composition Database, zinc intake was calculated in mg/day and mg/MJ for males (Appendix C) and females (Appendix D). Zinc intake calculations were performed only on subjects with at least two reporting days. Food consumption data in children were provided by nine studies, and data on adults were provided by eight studies, including one on pregnant women and adolescents. EFSA's estimates are based on consumption of foods, either fortified or not (i.e. without dietary supplements).

Average zinc intake ranged from 4.6 to 6.2 mg/day (1.1–1.5 mg/MJ) in children aged one to less than three years, from 5.5 to 9.3 mg/day (0.9–1.5 mg/MJ) in children aged 3 to < 10 years, from 6.8 to 14.5 mg/day (1.0–1.5 mg/MJ) in adolescents (10 to < 18 years) and from 8.0 to 14.0 mg/day (1.1–1.7 mg/MJ) in adults. Average daily intake (but not energy-adjusted intake) was, in most cases, slightly higher in males (see Appendix C) than in females (see Appendix D), mainly because of the larger quantities of food consumed per day.

The main food groups contributing to zinc intake were meat and meat products, grains and grain-based products, and milk and dairy products. Other food groups contributing to zinc intake were composite dishes in the Netherlands, Sweden and the UK; vegetables and vegetable products in Italy; and fish and fish products in Italy and Sweden (see Appendices D and E). The differences in the main contributors to zinc intake between males and females were small.

EFSA's zinc intake estimates in mg/day were compared with published intake values from the same survey and dataset and the same age class using the German EsKiMo and VELs surveys in children (Kersting and Clausen, 2003; Mensink et al., 2007), the DIPP study in Finnish children (Kyttälä et al., 2008; Kyttälä et al., 2010), the study in Finnish adolescents (Hoppu et al., 2010), the French national INCA2 survey (Afssa, 2009), the Irish NANS Survey (IUNA, 2011), the FINDIET 2012 Survey (Helldán et al., 2013), the Italian INRAN-SCAI Survey (Sette et al., 2011), the Dutch National Dietary Survey (van Rossum et al., 2011), the Swedish national survey Riksmaten (Amcoff et al., 2012) and the UK NDNS Survey (Bates et al., 2012) (Table 2).

Table 2: EFSA's average daily zinc intake estimates, expressed as percentages of intakes reported in the literature

Country	% of published intake (% range over different age classes in a specific survey)
Finland	88–104 (DIPP, for ages \geq 1 year), 102–103 (Finnish adolescents), 91–96 (FINDIET 2012)
France	93–112 (INCA2)
Germany	95–104 (VELS children), 110–111 (EsKiMo)
Ireland	112–120 (NANS)
Italy	89–93 (INRAN-SCAI)
NL	101–105 (Dutch National Dietary Survey)
Sweden	105–109 (Riksmaten)
UK	99–108 (NDNS–Rolling Programme, Years 1–3, for ages \geq 3 years)

Comparisons had inherent limitations in the case of the UK survey, as published intake values covered only the first two years of the survey, whereas EFSA data from the UK cover the first three years. In the survey in Finnish adolescents, published values were for two consecutive days of dietary recall, whereas EFSA data comprised two 48-hour dietary recalls. Likewise, comparisons were not optimal for the EsKiMo study and the DIPP study, because the published intake values included supplement consumption, whereas the EFSA estimates are based on food consumption only. However, according to these publications (Mensink et al., 2007; Kyttälä et al., 2010), zinc supplements were not among the major contributors to zinc intakes in these age classes. A comparison could not be undertaken for the

Latvian survey, as no matching publication was available. The EFSA estimates differed by up to about 10 % from the published values in Finland, France, Germany, Italy, the Netherlands, Sweden and the UK. The estimated Irish intakes were shown to be an overestimation of 12–20 %, which may partly be caused by the fact that data provided on composite dishes were almost completely disaggregated to the ingredient level, thereby not capturing possible zinc losses due to processing. Uncertainties in the estimates of all countries may be caused by inaccuracies in mapping food consumption data according to the FoodEx2 classification; by inaccuracies in analysing or estimating zinc composition for the food composition table, due to the use of borrowed zinc values from other countries in the food composition database; and by replacing missing zinc values with values of similar foods or food groups in the zinc intake estimation process. These uncertainties may, in principle, cause estimates of zinc intake that are either too high or too low.

3.3. Dietary phytate intake

The range of dietary phytate intake in the few European countries for which English-language data are available varies widely (Schlemmer et al., 2009; Amirabdollahian and Ash, 2010; Prynne et al., 2010). For example, median phytate intake reported in the UK based on the representative National Diet and Nutrition Survey ranged from 692 to 948 mg/day in men and from 538 to 807 mg/day in women of various age groups (Amirabdollahian and Ash, 2010), whereas lower intakes have been reported from studies in Scandinavian countries (Brune et al., 1989; Plaami and Kumpulainen, 1996) and in Italy (Carnovale et al., 1987) (see Appendix G).

The wide variation in phytate intake can partially be explained by differences in dietary patterns within and between countries; for example, dietary patterns dominated by plant foods are accompanied by a higher phytate intake. Besides dietary patterns, differences in food processing that can affect the phytate content of foods consumed, as well as methodological problems associated with phytate intake assessment, also contribute to variation among surveys. It has been estimated that adults ingest about 300 to 800 mg/day of phytate with a mixed diet and that the phytate intake increases to 700 to 1 400 mg/day for mixed diets with a high proportion of unrefined cereal grain products and legumes (Ingelmann et al., 1993; Schlemmer, 1995), whereas dietary phytate intake may be as high as 1 600 to 2 500 mg/day in adults on vegetarian diets (Bindra and Gibson, 1986; Ellis et al., 1987; Khokhar and Pushpanjali, 1994).

4. Overview of Dietary Reference Values and recommendations

4.1. Adults

The Nordic countries (NNR, 2004) estimated zinc requirements using the factorial method. For the estimate of endogenous faecal losses and losses via other routes, the figures of the US Institute of Medicine (IOM, 2001) were used. Endogenous intestinal losses were estimated to be 1.4 mg/day for both sexes based on observed losses at low intake (1–5 mg/day). Thus, it was assumed that 2.67 and 2.4 mg/day for men and women, respectively, have to be absorbed in order to replace all losses. Absorption efficiency of zinc from a mixed animal and vegetable protein diet, which is usually consumed in the Nordic countries, was assumed to be 40 %. The AR of zinc was therefore set at 6.4 and 5.7 mg/day, respectively, for men and women. The inter-individual variation in requirement was set at 15 %, resulting in recommended intakes of 9 mg/day for men and 7 mg/day for women. It was noted that this recommended intake probably has a high safety margin, as the ability to adapt to lower intake appears to be substantial. For NNR 2012, recommended intakes from NNR 2004 were maintained since no strong evidence has emerged to justify a change (Nordic Council of Ministers, 2014).

The German-speaking countries (D-A-CH, 2013) estimated obligatory daily zinc losses to be 2.2 mg in men and 1.6 mg in women based on data from King and Turnlund (1989). To replace these losses, an AR of 7.5 mg/day for men and 5.5 mg/day for women was calculated, assuming a mean zinc absorption efficiency of 30 % from mixed diets (Milne et al., 1983; Taylor et al., 1991). When twice

the coefficient of variation (CV) of 15 % was added to the AR, the recommended intakes were 10 mg/day for men and 7 mg/day for women.

IOM (2001) applied a factorial approach to calculate the minimal quantity of absorbed zinc necessary to replace the daily excretion of endogenous zinc. Losses via routes other than the intestine were regarded as unrelated to dietary zinc intake over a wide range encompassing zinc requirements. They were calculated to be 1.27 mg/day for men and 1.0 mg/day for women, considering data on average urinary excretion, integumental losses, and losses in semen or menstrual losses, respectively. IOM determined the correlation between the losses through excretion of endogenous zinc via the intestine and the quantity of zinc absorbed based on balance studies (Jackson et al., 1984; Turnlund et al., 1984; Wada et al., 1985; Turnlund et al., 1986; Taylor et al., 1991; Hunt J et al., 1992; Lee et al., 1993) and, taking into account a constant for non-faecal endogenous losses, calculated the average total minimal quantity of absorbed zinc required to offset losses as 3.84 mg/day for men and 3.3 mg/day for women. Considering the asymptotic regression of absorbed zinc on zinc intake observed in the balance studies, Estimated Average Requirements (EARs) of 9.4 mg/day for men and 6.8 mg/day for women were determined, corresponding to average fractional absorption of zinc (FAZ) of 0.41 and 0.48 for men and women, respectively. IOM noted that such EARs are supported by data from zinc depletion studies considering changes in functional endpoints (Wada and King, 1986; Grider et al., 1990; Beck et al., 1997b; Beck et al., 1997a) and a study on biochemical zinc status in healthy women (Gibson et al., 2000). Recommended Dietary Allowances (RDAs) of 11 mg/day for men and 8 mg/day for women were set by adding twice the CV of 10 % to the EARs.

WHO/FAO (2004) applied a factorial approach, which involved totalling the requirements for tissue maintenance, metabolism and endogenous losses. The body's ability to adapt to different levels of zinc intake was taken into consideration by defining the normative requirement for absorbed zinc as the obligatory loss during the early phase of zinc depletion before adaptive reductions in excretion take place. The normative requirements for absorbed zinc were estimated to be 1.4 mg/day for men and 1.0 mg/day for women by adding estimations of faecal, urinary and skin losses (data derived from Milne et al. (1983); Milne et al. (1987); Taylor et al. (1991)). To translate these estimates into requirements for dietary zinc, the influence of the nature of the diet (i.e. its content of promoters and inhibitors of zinc absorption) and the efficiency of absorption of potentially available zinc were considered. Overall, three categories of diets were distinguished, characterised by high, moderate and low zinc bioavailability, and the absorption efficiency figures that were estimated to be adequate to meet the normative requirements for absorbed zinc were 50 %, 30 % and 15 %, respectively. Corresponding average individual dietary requirements were estimated to be 36, 59 and 119 µg/kg body weight per day for women and 43, 72 and 144 µg/kg body weight per day for men. Assuming an inter-individual variation of zinc requirements of 25 %, the recommended nutrient intakes are 3.0, 4.9 and 9.8 mg/day for women and 4.2, 7.0 and 14.0 mg/day for men for diets of high, moderate and low zinc bioavailability, respectively.

Afssa (2001) set two levels of recommended intakes, depending on the dietary content of products of animal origin. A daily intake of 7 mg/day for women and 9 mg/day for men was recommended if the diet contains relatively high amounts of products of animal origin (estimated intestinal zinc absorption of 30 %). An increased daily intake of 12 mg/day for women and 14 mg/day for men was proposed if the diet contained relatively low amounts of products of animal origin (estimated intestinal absorption of 20 %).

The Netherlands Food and Nutrition Council (1992) applied a factorial approach. Total zinc losses were estimated to be 1.3–1.9 mg/day for men and 1.1–1.7 mg/day for women, considering data on average urinary excretion, integumental losses and additional losses in semen for men. Menstrual zinc losses were considered negligible. Minimum requirements were estimated to be 5.2–7.6 mg/day for men and 4.4–6.8 mg/day for women, applying an estimated average absorption efficiency of 25 %. The council proposed adequate ranges of intakes of 7.0–10.0 mg/day for men and 6.0–9.0 mg/day for women, assuming an inter-individual variation in zinc losses of 20 %.

The UK Committee on Medical Aspects of Food Policy (COMA) (DH, 1991) assessed the zinc requirement on the basis of factorial analyses using measurements of basal losses during metabolic studies of deprivation, the turnover time of radiolabelled endogenous zinc pools, and deduction from metabolic studies of patients receiving total parenteral nutrition. Minimal zinc losses in the order of 2.2 and 1.6 mg/day in men and women, respectively, were estimated, considering data on basal faecal and urinary losses, and on losses via skin, hair, semen and menstruation, where appropriate (Hambidge et al., 1986; King and Turnlund, 1989; Taylor et al., 1991). Assuming an absorption efficiency of 30 %, these figures translate into EARs of 7.3 mg/day in men and 5.5 mg/day in women. Reference Nutrient Intakes (RNIs) of 9.5 and 7.0 mg/day were set for men and women, respectively. Based on the same considerations, the SCF also proposed PRIs of 9.5 mg/day for men and 7.0 mg/day for women (SCF, 1993).

Table 3: Overview of Dietary Reference Values for zinc for adults

	NNR (2012)	D-A-CH (2013)	WHO/FAO (2004)			Afssa (2001) ^(a)		IOM (2001)	SCF (1993)	NL (1992) ^(b)	DH (1991)
			High BA (50 %)	Moderate BA (30 %)	Low BA (15 %)	IA of 20 %	IA of 30 %				
Age (years)	≥ 18	≥ 19	≥ 19	≥ 19	≥ 19	≥ 20	≥ 20	≥ 19	≥ 18	≥ 19	≥ 19
PRI men (mg/day)	9	10	4.2	7.0	14.0	14	9	11	9.5	7–10	9.5
PRI women (mg/day)	7	7	3.0	4.9	9.8	12	7	8	7	6–9	7.0

NL, Netherlands Food and Nutrition Council; BA, bioavailability; IA, intestinal absorption.

(a): The values vary according to the bioavailability of zinc from the diet: for predominantly vegetarian diets, a bioavailability of 20 % is assumed, and, for balanced diets rich in animal products, including meat products, a bioavailability of 30 % is assumed.

(b): Adequate range of daily intake.

4.2. Infants and children

The Nordic countries (NNR, 2004) noted that data on endogenous losses of zinc at different levels of intake are almost completely lacking for children. It was also noted that, in relation to body weight, children appear to have larger losses of zinc than adults. The need for growth was estimated to be 175 µg/kg body weight per day during the first month of life, decreasing to approximately 30 µg/kg body weight per day at 9–12 months of age (Krebs and Hambidge, 1986). For growing children, the need for zinc was based on basal losses of 0.1 mg/kg body weight per day and a zinc content in new tissue of 30 µg/g. For adolescents, growth was assumed to result in an average zinc content in new tissue of 23 µg/g, due to an increase in fat tissue with a lower zinc content than children. The physiological needs for rapidly growing adolescents were considered to be increased by 0.3–0.4 mg/day. Applying the same principles as for adults, the recommended zinc intakes vary from 2 mg/day in the youngest age group to 12 mg/day for adolescent boys. For NNR 2012, the recommended intakes from 2004 were kept unchanged (Nordic Council of Ministers, 2014).

WHO/FAO (2004) considered evidence that the maintenance requirement in infants is influenced by the nature of the diet (Krebs and Hambidge, 1986; Krebs, 1993) and assumed endogenous losses of zinc to be 20 µg/kg body weight per day for human milk-fed infants and about 30–40 µg/kg body weight per day for infants fed formula or weaning foods. The estimated zinc requirements for infant growth were set at 120 and 140 µg/kg body weight per day for female and male infants, respectively, for the first three months of life. These values decreased to 33 µg/kg body weight per day for infants aged 6–12 months. A bioavailability of 80 % was assumed for exclusively breast-fed infants, and a bioavailability of 15 or 30 % was assumed for formula-fed infants, depending on the type of formula. For infants up to six months of age, it was assumed that the inter-individual variation of zinc

requirements is 12.5 % and is the same for breast-fed (derived from Vuori (1979b)) and formula-fed infants. After that age, a CV of 25 % was assumed. For other age groups, an average loss of 0.57 µg/kcal of resting energy expenditure (REE) was derived by extrapolating from the adult REE values. For children aged 1–10 years, the requirements for growth were based on the assumption that new tissue contains 30 µg zinc/g. For adolescent growth, a tissue zinc content of 23 µg/g was assumed. Taking into account that pubertal growth spurts increase physiological zinc requirements substantially, growth of adolescent males was assumed to correspond to an increase in body zinc requirement of about 0.5 mg/day.

For infants aged 0–6 months, IOM (2001) set an Adequate Intake (AI) of 2.0 mg/day, which reflects the observed mean zinc intake of infants exclusively fed human milk. Human milk alone was considered an inadequate source of zinc after the first six months of life, and EARs for older infants and children were based on the factorial approach. Excretion of endogenous zinc was estimated by extrapolation from measured values for either adults or younger infants. Requirements for growth were derived from chemical analyses of zinc concentrations of infant and adult tissues (Widdowson and Dickerson, 1964) and average daily accretion of new tissue (Kuczmarski et al., 2000). For pre-adolescent children (seven months to 13 years), a conservative FAZ of 0.3 was applied (Fairweather-Tait et al., 1995; Davidsson et al., 1996), whereas a FAZ of 0.4 was used for adolescents (14 to 18 years). IOM noted that growth data from supplementation studies with zinc in children aged 7 to 12 months (Walravens et al., 1989) and four to eight years (Walravens et al., 1983; Gibson et al., 1989) were consistent with the EARs derived from the factorial approach. Corresponding RDAs were set by adding twice the CV of 10 % to the EARs.

The Netherlands Food and Nutrition Council (1992) applied a factorial approach. Total zinc losses were extrapolated from adults on the basis of metabolic weight ($\text{kg}^{0.75}$). For the first half year of life, the requirement for growth was estimated at 400 µg/day (Widdowson and Dickerson, 1964; Sandstead, 1973; WHO, 1973), on the basis of the increase in fat-free body mass and the zinc content per kg of fat-free body mass. An estimated average absorption efficiency of 25 % was applied to derive the minimum dietary requirements. Corresponding adequate ranges of intakes were set, assuming inter-individual variations of 20 % in zinc loss and 15 % in zinc requirement for growth.

The UK COMA (DH, 1991) used a factorial approach to calculate daily zinc requirements for infants and children. Growth increments were estimated on the basis of growth progressing along the 50th percentile and on a lean tissue zinc content of 30 µg/g. Urine and sweat zinc losses were taken as 10 and 20 µg/kg body weight per day, respectively, and faecal losses as 77 µg/kg body weight per day. This led to a daily requirement of absorbed zinc of 1.0 mg. Taking into account an absorption efficiency of 30 % from infant formula, an EAR of 3.3 mg/day was derived. The RNI was set at 4 mg/day by adding twice the CV of 10 % to the EAR. For children over one year of age, RNIs were based on interpolated basal losses from adults and calculated increments for growth, assuming an absorption efficiency of 30 %. The SCF set the PRIs for infants and children based on the same approaches (SCF, 1993).

Table 4: Overview of Dietary Reference Values for zinc for children from four months of age

	NNR (2012)	D-A-CH (2013)	WHO/FAO (2004)			Afssa (2001) ^(a)		IOM (2001)	SCF (1993)	NL (1992) ^(b)	DH (1991)
			High BA (50 %)	Moderate BA (30 %)	Low BA (15 %)	IA of 20 %	IA of 30 %				
Age (months)	6–11	4– < 12	7–12	7–12	7–12			7–12	6–11	6–12	7–12
PRI (mg/day)	5	2	0.8 ^(c) , 2.5 ^(d)	4.1	8.4			3	4	3–4	5
Age (years)	1– < 2	1– < 4	1–3	1–3	1–3	1–3	1–3	1–3	1–3	1–4	1–3
PRI (mg/day)	5	3	2.4	4.1	8.3	8	5	3	4	3–4	5
Age (years)	2–5	4– < 7	4–6	4–6	4–6	4–9	4–9	4–8	4–6	4–7	4–6
PRI (mg/day)	6	5	2.9	4.8	9.6	11	6	5	6	4–5	6.5
Age (years)	6–9	7– < 10	7–9	7–9	7–9			9–13	7–10	7–10	7–10
PRI (mg/day)	7	7	3.3	5.6	11.2			8	7	4–6	7
Age (years)	10–13	10– < 13	10–18	10–18	10–18	10–12	10–12	14–18	11–14	10–13	11–14
PRI: boys (mg/day)	11	9	5.1	8.6	17.1	14	9	11	9	5–7	9
PRI: girls (mg/day)	8	7	4.3	7.2	14.4	13	9	9	9	5–7	9
Age (years)	14–17	13– < 15				13–19	13–19		15–17	13–16	15–18
PRI: boys (mg/day)	12	9.5				14	11		9	7–10	9.5
PRI: girls (mg/day)	9	7				11	9		7	7–10	7.0
Age (years)		15– < 19								16–19	
PRI: boys (mg/day)		10								8–11	
PRI: girls (mg/day)		7								6–9	

NL, Netherlands Food and Nutrition Council; BA, bioavailability; IA, intestinal absorption.

(a): The values vary according to the bioavailability of zinc from the diet: for predominantly vegetarian diets, a bioavailability of 20 % is assumed, and, for balanced diets rich in animal products, including meat products, a bioavailability of 30 % is assumed.

(b): Adequate range of daily intake.

(c): Exclusively human milk-fed infants.

(d): Not applicable to infants consuming human milk only.

4.3. Pregnancy

The Nordic countries (Nordic Council of Ministers, 2014) considered that the total need for zinc during pregnancy for the fetus, placenta and other tissues is approximately 100 mg (King, 2000), and that studies on whether or not homeostatic adjustments occur during pregnancy are inconclusive (Swanson and King, 1982; Fung et al., 1997). The recommended intakes were based on an assumed increase of the physiological requirement by 0.7 mg/day. With adjustment for absorption, the additional recommended intake was set at 2 mg/day.

The German-speaking countries (D-A-CH, 2013) considered that the average additional requirement of absorbed zinc during the second half of pregnancy is 0.8 mg/day and recommended an additional zinc intake of 3 mg/day from the fourth month of pregnancy onwards.

WHO/FAO (2004) considered an estimated amount of zinc retained during pregnancy of 100 mg (Lentner, 1984; Swanson and King, 1987). During the third trimester, the physiological requirement of zinc was assumed to be approximately twice as high as that of non-pregnant women.

Applying a factorial approach, IOM (2001) determined an additional requirement of 2.7 mg/day, considering the highest average daily rate of zinc accumulation by maternal and fetal tissues of 0.73 mg observed during the fourth quarter of pregnancy (Swanson and King, 1987), and an estimated average FAZ of 0.27 (Turnlund et al., 1991; Hunt J et al., 1992; Sian et al., 1996; Fung et al., 1997; Hunt et al., 1998; Miller et al., 1998). EARs of 10 mg/day for pregnant adolescents aged 14–18 years and of 9.5 mg/day for pregnant women were derived, and RDAs were set at 12 mg/day and 11 mg/day, respectively, by adding twice the CV of 10 % to the EARs and rounding to the nearest 1 mg.

The Netherlands Food and Nutrition Council (1992) considered extra zinc requirements of 0.6, 0.9 and 1.0 mg/day during the first, second and third trimesters of pregnancy, respectively, according to WHO (1973), and set adequate ranges of intakes of 9–12 mg/day during the first trimester and of 11–15 mg/day during the second and third trimesters.

The UK COMA (DH, 1991) noted that, although there was evidence that extra zinc is required during pregnancy, studies have shown no increase in customary daily zinc intake by pregnant women and no benefit from zinc supplements (Mahomed et al., 1989). The committee considered it probable that, in healthy women, metabolic adaptation ensures an adequate transfer of zinc to the fetus, and no increment was proposed for pregnant women. The SCF (1993) adopted the same approach.

Table 5: Overview of Dietary Reference Values for zinc for pregnant women

	NNR (2012)	D-A-CH (2013)	WHO/FAO (2004)			Afssa (2001) ^(a)		IOM (1998)	SCF (1993)	NL (1992) ^(b)	DH (1991)
			High BA (50 %)	Moderate BA (30 %)	Low BA (15 %)	IA of 20 %	IA of 30 %				
Age (years)								14–18			
PRI (mg/day)	9	10 ^(c)	3.4 ^(d) 4.2 ^(f) 6.0 ^(g)	5.5 ^(d) 7.0 ^(f) 10.0 ^(g)	11.0 ^(d) 14.0 ^(f) 20.0 ^(g)	16 ^(e)	11 ^(e)	12	7	9–12 ^(d) 11–15 ^(f) 11–15 ^(g)	7
Age (years)								19–50			
PRI (mg/day)								11			

NL, Netherlands Food and Nutrition Council; BA, bioavailability; IA, intestinal absorption.

(a): The values vary according to the bioavailability of zinc from the diet: for predominantly vegetarian diets, a bioavailability of 20 % is assumed, and, for balanced diets rich in animal products, including meat products, a bioavailability of 30 % is assumed.

(b): Adequate range of daily intake.

(c): From four months of age.

(d): First trimester.

(e): Increases during gestation; value is for the third trimester.

(f): Second trimester.

(g): Third trimester.

4.4. Lactation

The Nordic countries (Nordic Council of Ministers, 2014) considered milk zinc concentration to be 2.5 mg/L in the first month of lactation and to fall to approximately 0.7 mg/L after four months (Krebs and Hambidge, 1986). An additional requirement of 1.7 mg/day for the replacement of zinc losses with human milk was assumed. Taking into account absorption efficiency, an additional dietary intake of 4 mg/day was recommended.

Assuming that fully breast-fed infants receive 1 mg zinc/day with 0.75 L of human milk, the German-speaking countries (D-A-CH, 2013) considered that the average additional requirement of absorbed zinc during lactation is 1 mg/day and recommended an additional zinc intake of 4 mg/day.

From data on maternal milk volume and milk zinc concentration, WHO/FAO (2004) estimated the daily output of zinc in milk to be 1.4 mg/day during the first three months of lactation, 0.8 mg/day from three to six months of lactation and 0.5 mg/day thereafter (Vuori, 1979b; Krebs and Hambidge, 1986; Casey et al., 1989). In setting the requirements for early lactation (zero to three months post partum), it was assumed that around 0.5 mg/day is covered by postnatal involution of the uterus and from skeletal resorption.

IOM (2001) estimated the losses of zinc in human milk to be 3 mg/L at four weeks and 1.2 mg/L at 24 weeks post partum on the basis of observed average zinc concentrations in human milk (Moser-Veillon and Reynolds, 1990; Krebs et al., 1995) and an average secretion of 0.78 L milk/day. IOM also took into account that zinc is released from the post partum involution of the uterus and the decreased maternal blood volume (King and Turnlund, 1989), and assumed that it is available for re-utilisation. Overall, the average calculated increased requirement of absorbed zinc was 1.35 mg/day. Applying a FAZ of 0.377 during lactation (Fung et al., 1997), the additional zinc requirement was estimated to be 3.6 mg/day.

To compensate for zinc transfer into breast milk, the Netherlands Food and Nutrition Council (1992) estimated an additional requirement of 2.4 mg/day during the first month of lactation, of 2.0 mg/day during the second and third months of lactation and of 1.2 mg/day thereafter (Vuori, 1979a; Ruz, 1984; Casey et al., 1985).

During lactation, the UK COMA (DH, 1991) proposed an increment of 6 mg/day during the initial four months of lactation and 2.5 mg/day thereafter, on the basis of a daily milk volume of 0.85 L and zinc losses of 2.13 and 0.94 mg/day, respectively. The SCF (1993) proposed an additional dietary intake of 5 mg/day during lactation to cover the amount of zinc transferred into milk.

Table 6: Overview of Dietary Reference Values for zinc for lactating women

	NNR (2012)	D-A-CH (2013)	WHO/FAO (2004)			Afssa (2001) ^(a)		IOM (1998)	SCF (1993)	NL (1992) ^(b)	DH (1991)
			High BA (50 %)	Moderate BA (30 %)	Low BA (15 %)	IA of 20 %	IA of 30 %				
Age (years)								14–18			
PRI (mg/day)	11	11	5.8 ^(c) 5.3 ^(f) 4.3 ^(h)	9.5 ^(c) 8.8 ^(f) 7.2 ^(h)	19.0 ^(c) 17.5 ^(f) 14.4 ^(h)	23 ^(d)	15 ^(d)	13	12	16–20 ^(c) 13–16 ^(g)	+ 6 ^(e) + 2.5 ^(g)
Age (years)								19–50			
PRI (mg/day)								12			

NL, Netherlands Food and Nutrition Council; BA, bioavailability; IA, intestinal absorption.

(a): The values vary according to the bioavailability of zinc from the diet: for predominantly vegetarian diets, a bioavailability of 20 % is assumed, and, for balanced diets rich in animal products, including meat products, a bioavailability of 30 % is assumed.

(b): Adequate range of daily intake.

(c): Zero to three months post partum.

(d): Decreases during lactation; values for the first month.

(e): Zero to four months post partum.

(f): Three to six months post partum.

(g): After three (Netherlands)/four (UK) months post partum.

(h): 6–12 months post partum.

5. Criteria (endpoints) on which to base Dietary Reference Values

5.1. Indicators of adult zinc requirement

The lack of sensitive specific biomarkers or clinical features of “mild” zinc deficiency precludes the possibility of using a dose–response approach for estimating zinc requirements. Theoretically, the traditional balance technique combined with urine and integumental zinc losses has the potential to provide information on zinc requirements. In practice, despite a long history of such measurements, this approach has not provided satisfactory results. The small difference obtained from subtracting total faecal excretion from total ingested zinc to derive net absorption detracts from the accuracy and reliability of this approach, and it does not provide information on true zinc absorption. The advent and progressive improvement of equipment and techniques for the application of zinc stable isotopes to studies of zinc homeostasis has progressively facilitated the application of a factorial approach in the estimation of zinc requirements.

The estimation of zinc requirements by the factorial approach requires two stages. The first is the estimation of physiological requirements, defined as the minimum quantity of absorbed zinc necessary to match losses of endogenous zinc and to meet any additional requirements for absorbed zinc that may be necessary in lactation as well as for growth in healthy well-nourished infants and children and in pregnancy. The second stage is the determination of the quantity of dietary zinc available for absorption that is necessary to meet the physiological requirement.

5.1.1. Physiological requirement

5.1.1.1. Identification of studies, data extraction and assessment of methodological quality

A total of 15 studies were identified from the published literature that included data on endogenous faecal zinc (EFZ) and total absorbed zinc (TAZ) for the estimation of physiological zinc requirement. Fourteen studies were identified by comprehensive literature searches in PubMed up to mid-February 2014 using the following search string: zinc[TI] AND ((endogenous f*ecal) OR (intestinal excretion endogenous) OR (intestinal endogenous losses) OR isotope* OR compartmental OR extrinsic* OR balance) AND ((total absorbed) OR absorption OR retention OR depletion OR pool* OR metabolism), with a limit to human studies. One study was identified by hand-searching the reference list of studies retrieved by the comprehensive literature search.

Inclusion criteria were the following: studies of healthy adults, whole-day isotope studies of true zinc absorption, studies with information on body weight of participants, and retrieval of individual data at time of final data analyses. Second stage exclusion criteria included physiologically implausible data for EFZ and evidence of clinical disease.

After detailed review of all potential data and elimination of studies that had significant methodological limitations, the methodologies used in the studies included in the final analyses are considered to be reliable. For example, only studies that employed isotope tracer methods for determining zinc absorption were considered acceptable.

5.1.1.2. Inclusion of studies

For 15 studies, data from the individual study participants were supplied by the authors. Data quality was assessed initially based on whether or not data were physiologically plausible. Initial evaluation identified two study designs (compartmental modelling and faecal isotope dilution) that included intravenous administration of a zinc stable isotope tracer and its dilution in the faeces, i.e. isotopic labelling of the endogenous zinc appearing in the faeces, and which consistently provided physiologically plausible data. The quality of a third method involving isotopic measurement of absorption coupled with gastro-intestinal balance of non-labelled zinc was judged on the physiologically plausible results (see Section 5.1.1.3 and Appendix H). Information extracted from the studies were total dietary zinc ($\mu\text{mol/day}$ or mg/day), total dietary phytate ($\mu\text{mol/day}$ or mg/day), total absorbed zinc ($\mu\text{mol/day}$ or mg/day), faecal excretion of endogenous zinc ($\mu\text{mol/day}$ or mg/day), daily

urinary zinc excretion ($\mu\text{mol/day}$ or mg/day), subject body weight (kg) and subject height (m) (see also Appendix I).

5.1.1.3. Inclusion of individual data

Thirteen individual data points, all from studies utilising the zinc absorption–intestinal balance technique, had physiologically impossible negative values for EFZ, and accordingly were omitted from subsequent calculations. After omitting these individual data points, the remaining data from these studies (Wada et al., 1985; Hunt J et al., 1992; Hunt et al., 1995; Hunt et al., 1998) and the data from the study of Sandstrom et al. (2000) were evaluated in comparison to the data from studies which had no negative EFZ values, the majority of which were data from studies using isotope tracer methods (isotope dilution, compartmental modelling) to directly measure EFZ and were, therefore, the most reliable. The EFZ data in question were found to differ from the standard data in distribution and relationships to other variables, bringing into question their accuracy (see Appendix H for details).

In view of the uncertainty about the accuracy of the EFZ results in particular studies (Wada et al., 1985; Hunt J et al., 1992; Hunt et al., 1995; Hunt et al., 1998; Sandstrom et al., 2000), the Panel decided to exclude data from these studies from subsequent consideration. Also excluded was one participant in the study of Taylor et al. (1991), who had biochemical and haematological indices of hepatitis possibly caused by alcohol abuse. Exclusion involved a total of 103 data points (more details on the preliminary data analysis are given in Appendix H).

The final numbers of subjects contributing data to the estimate of physiological zinc requirements were 31 males and 54 females from a total of 10 studies. These included data from all available published studies that contained the data required, including a study in China (Sian et al., 1996) that had results that fit well with data from studies in the USA and Europe. Although several studies used semipurified formula diets, in most studies, the diets were composed of conventional foods, sometimes based on the habitual diets of the subjects. The period of time during which the subjects consumed these constant diets prior to the isotope studies varied from five days to five weeks, although seven days was the most common duration. Relevant individual data for the 85 subjects included in the final estimates are given in Appendix I. Dietary zinc intake of subjects ranged from 0.8 to 29 mg/day (see Appendix I). Dietary phytate intake was available for some of the studies using conventional food. Mean phytate intake reported in 4 of the 10 studies ranged from 585 to 835 mg/day (see Appendix I).

5.1.1.4. Estimation of endogenous zinc losses

Total endogenous zinc losses were calculated as the sum of losses via faeces, urine, combined integument and sweat (0.5 mg and 0.3 mg/day for men and women, respectively; see explanation below); semen (0.1 mg/day (Hunt CD et al., 1992; Johnson et al., 1993)); and menses (0.01 mg/day (Hess et al., 1977)), of which endogenous faecal zinc is the major component. Urinary zinc losses were reported for 57 of the 85 subjects. For the remaining 28 individuals, estimated urinary zinc losses based on sex were used. The estimated mean urinary zinc losses were 0.5 mg/day for men and 0.3 mg/day for women. These were the averages of the 53 reported values (22 men, 31 women) for subjects in normal zinc status. Integumental and sweat zinc losses were estimated from published studies. An estimate of 0.5 mg/day for men was obtained from studies of whole body surface zinc losses in men (Jacob et al., 1981; Milne et al., 1983; Johnson et al., 1993). The estimate of 0.3 mg/day for women was calculated by multiplying the value for men by the female to male ratio of sweat zinc losses observed in studies of whole body sweat zinc losses and whole body sweat rates in men and women (Cohn and Emmett, 1978; Avellini et al., 1980; Frye and Kamon, 1983; Tipton et al., 1993; DeRuisseau et al., 2002; Hazelhurst and Claassen, 2006). These studies reported female to male ratios for sweat zinc loss rates between 0.5 and 0.7, while sweat zinc concentrations were similar.

5.1.1.5. Modelling of zinc requirements

The assumptions that the regression errors were normally distributed and exhibited constant variance, and that the model was valid, were checked primarily by visual examination of plots of the residuals.

Both the raw residuals and the externally studentised residuals were examined. Normality of the residuals was also tested with the D'Agostino–Pearson and Shapiro–Wilk tests. Externally studentised residuals were also examined for outliers. The variance inflation factor was used to evaluate multicollinearity. Details of the regression diagnostics are provided in Appendix J.

A multiple regression analysis was used to evaluate the relationship of TAZ with (TAZ – total endogenous zinc losses), the predictor variable of interest, and with sex and body size as covariates. A major finding from this analysis was that TAZ varies significantly with body size expressed as weight, height, body mass index (BMI) or surface area. Each of the body size variables was made a covariate along with (TAZ – total endogenous zinc losses) in four separate regression models. In each case, the body size variable was significant, with p-values < 0.001, except for BMI, which had a p-value of 0.013. The R² values for the models with body weight, height, BMI and body surface area variables were 0.46, 0.42, 0.37 and 0.47, respectively. A variety of other unmeasured/unmeasurable variables presumably also contributed, ranging from inter-/intra-research facility variation to possible biological factors, for example the extent of up- or down-regulation of zinc transporters and other proteins involved in the absorption of zinc by the enterocyte, or variations in body zinc stores. The variable for sex was entered in each model. With the exception of BMI, none of the models demonstrated a significant sex effect, as sex differences were apparently accounted for by the body size covariate. In the BMI model, sex was a significant predictor (p-value: 0.011). The equation with the covariate body weight will be used for reasons of convenience and accuracy of measurement. The equation resulting from a least-squares fit to the body weight data, and which therefore links TAZ to body weight and the difference of TAZ minus total endogenous zinc losses, is the following Equation 1:

$$\text{TAZ [mg/day]} = 0.642 + 0.038 \times \text{body weight [kg]} + 0.716 \times (\text{TAZ} - \text{total endogenous zinc losses [mg/day]})$$

Details of parameter estimates for the model corresponding to Equation 1 are shown in Table 7.

Table 7: Details of parameter estimates for the model describing the relationship between TAZ and the difference between TAZ and total endogenous zinc losses, as well as body size, corresponding to Equation 1 above

Parameter	Estimate	95 % confidence limits	p-value
Intercept	0.642	–0.403, 1.687	0.23
Body weight	0.038	0.022, 0.054	< 0.0001
TAZ – total endogenous zinc losses	0.716	0.512, 0.919	< 0.0001

TAZ, total absorbed zinc.

Examination of the residuals indicated that errors were normally distributed and exhibited constant variance, and that there was no deviation of the model from the data. One outlier with an externally studentised residual of 3.7 was observed. Re-examination of the source of the outlier indicated no basis for its removal; therefore, the outlier was retained. There was no evidence of collinearity of predictors.

As mentioned below, the residuals did not indicate any deviation from the linear model. Nonetheless, polynomial terms were added to the model to explore the possibility of non-linear relationships. Only the second order polynomial of (TAZ – total endogenous zinc losses) was significant, but this was due to the presence of the outlying point (see above). When this point was momentarily removed, no significant polynomial terms were observed. It was therefore concluded that there was no evidence of non-linear relationships.

The physiological requirement is equivalent to TAZ when the difference between absorbed zinc and total endogenous zinc losses equals zero at a given body weight. Therefore, the equation for estimating the physiological requirement (Equation 2) was derived from Equation 1 by removal of the (TAZ – total endogenous zinc losses) term:

Physiological zinc requirement [mg/day] = $0.642 + 0.038 \times \text{body weight [kg]}$ (Equation 2)

This equation is valid over a body weight range of roughly 40 to 100 kg. The size of the 95 % confidence interval (CI) for the estimation of the physiological requirements in Table 9 varies between ± 0.23 and ± 0.25 .

Recent developments which have facilitated the estimation of physiological requirements include a simple model for estimating physiological requirements; the use of individual rather than mean data; recognition of the inaccuracies associated with the zinc absorption–intestinal balance technique in some studies, which were omitted from final estimates; recognition of the extent of the impact of body weight on the estimation of intestinal excretion of endogenous zinc and, therefore, of physiological requirements; and recognition of the absence of a sex effect on endogenous faecal zinc losses beyond that accounted for by differences in body weight.

5.1.2. Estimation of dietary zinc intake to meet physiological requirement

Although experimental data are still limited (Hambidge et al., 2010), there are also theoretical reasons for supporting the conclusion that the relationship between TAZ and total dietary zinc (TDZ) is most appropriately fitted with saturation response modelling. Therefore, the intercept of the TAZ necessary to meet physiological requirements with the saturation response model (Morgan et al., 1975) for the population studied should give the AR for that population. Saturation response modelling is based on the assumption that zinc absorption is a carrier-mediated, saturable process, and this is used to characterise the relationship between the quantity of zinc absorbed and the quantity of zinc ingested. It is accomplished by fitting one of several appropriate models to data from isotope tracer studies of zinc absorption using non-linear regression analysis.

5.1.2.1. Effect of dietary zinc and phytate on absorbed zinc

Although quantities vary greatly, diets containing plant foods, i.e. virtually all non-synthetic diets, contain phytate. The luminal contents of the duodenum and jejunum, especially phytate, can have a major impact on the percentage of zinc available for absorption (see Section 2.3.1). A trivariate model of TAZ as a function of dietary zinc and dietary phytate, based on saturation response modelling, has been found to account for more than 80 % of the variance in TAZ (Miller et al., 2007).

5.1.2.2. Identification of studies, data extraction and assessment of methodological quality

The data used in this trivariate model of the relationship between zinc absorption, and dietary zinc and phytate were 72 mean data points (reflecting 650 individual measurements) reported in 18 publications. These are the data used in the development and early application of the model (Miller et al., 2007; Hambidge et al., 2010). The eligibility criteria were the following: studies of healthy adults, whole-day isotope studies of true zinc absorption, reporting measurements of TDZ, and total dietary phytate (TDP) and TAZ. No extensive literature search was performed in addition to that performed by the EUROpean micronutrient RECommendations Aligned (EURRECA) network on factors affecting zinc bioavailability (Lowe et al., 2013); relevant publications were identified through knowledge of the existing work of the small number of investigators in this field of research and ongoing monitoring of the new literature. All the data came from research groups having extensive experience with the application of isotope tracer methods to the study of zinc absorption. A formal assessment of the quality of the data was not performed. The data are summarised in Appendix K.

In all studies, participants ate controlled diets which contained known quantities of zinc and phytate (in many cases dietary calcium, iron and protein were also measured) in free-living and metabolic study environments. After varying lengths of time on the study diets, zinc stable or radio isotope tracers were administered, and enrichment was measured in body tissues and/or excretions to determine absorption. TDZ, TDP and TAZ data from these studies were used to develop the saturation response zinc absorption model (Miller et al., 2007).

5.1.2.3. Modelling of the saturation response model

The assumptions that the regression errors were normally distributed and exhibited constant variance, and that the model was valid, were checked primarily by visual examination of plots of the raw and standardised residuals. Normality of the residuals was also tested with the Shapiro–Wilk test. Residuals were also examined for outliers.

The trivariate saturation response model is described by the following Equation 3:

$$TAZ = 0.5 * \left(0.033 * \left(1 + \frac{TDP}{0.68} \right) + 0.091 + TDZ - \sqrt{\left(0.033 * \left(1 + \frac{TDP}{0.68} \right) + 0.091 + TDZ \right)^2 - 4 * 0.091 * TDZ} \right)$$

where TAZ, TDP and TDZ are all in mmol/day. Units are converted to mg/day for plots and values reported in this Opinion. The range of TDP and TDZ of the data are 0–3 730 and 4–21 mg/day, respectively. The R² of the fit was 0.81. The TAZ predicted by this model for the range of dietary zinc intake and selected dietary phytate levels is shown in Figure 1.

Table 8: Details of parameter estimates in the model on the relationship between zinc absorption, and dietary zinc and phytate corresponding to Equation 3 above

Parameter	Estimate	95 % confidence limits	p-value
Amax	0.091	0.079, 0.108	< 0.0001
KP	0.678	0.290, 1.230	0.0029
KT	0.033	0.014, 0.062	0.0038

Amax, maximum possible absorbed zinc; KP, zinc–phytate binding equilibrium dissociation constant, rounded to 0.68 in Equation 3; KT, zinc–transporter binding equilibrium dissociation constant.

Examination of the residuals indicated that errors were normally distributed and exhibited constant variance, and that there was no deviation of the model from the data. No outliers were detected. Details of the regression diagnostics are provided in Appendix J.




Figure 1: Saturation response model predictions of total absorbed zinc (TAZ) for selected levels of dietary phytate. Portions of the curves between total dietary zinc (TDZ) values of 0 and 4 mg/day are extrapolated, as there were no zinc intake data within that range. A three-dimensional plot giving a complete range of TAZ as a function of TDZ and dietary phytate is given in Appendix L

The dietary zinc intake required to meet the AR associated with different body weights (as predicted by the model described in Section 5.1.1.5) can be derived from the intersection of the respective physiological zinc requirements (identified on the axes of the absorbed zinc) with the saturation response model curve back-predicting the dietary zinc intake conditional to an expected level of phytate intake.

This is illustrated in Figure 2, derived from the established model of phytate effect (Miller et al., 2007; Hambidge et al., 2010). The curves show the relationships of absorbed zinc to dietary zinc for dietary phytate levels of 0 and 900 mg/day, as predicted by the saturation response model. The horizontal dashed lines indicate the physiological requirements for males and females based on measured body weights of the subjects in the study (i.e. 59.1 kg for females and 72.7 kg for males, see Appendix I). Average dietary zinc requirements of these subjects are the corresponding dietary zinc intakes for the intersections of physiological zinc requirement values with the model curves. In Table 9, dietary zinc requirements, depending on the level of phytate intake, are shown for the subjects who contributed data to establish the physiological requirement model.

Table 9: Average dietary zinc requirements depending on phytate intake and body weight

Body weight (kg)	Physiological requirement (mg Zn/day)	Average Requirement (mg Zn/day) for			
		300 mg/day of dietary phytate	600 mg/day of dietary phytate	900 mg/day of dietary phytate	1 200 mg/day of dietary phytate
72.7 ^(a)	3.4	8.2	10.2	12.1	14.0
59.1 ^(b)	2.9	6.3	7.7	9.1	10.4

(a): Mean of the body weight data for men used to establish the physiological requirement (Equations 1 and 2) as described above (see Appendix I).

(b): Mean of the body weight data for women used to establish the physiological requirement (Equations 1 and 2) as described above (see Appendix I).




Figure 2: Relationships between absorbed zinc and dietary zinc for dietary phytate levels of 0 and 900 mg/day, as predicted by the saturation response model

No data are available for subjects older than 52 years. Although muscle mass decreases with increasing age, the turnover of zinc in muscle is slow. Without the relevant experimental data, the Panel considers that the basis for setting DRVs for older adults should be the same as for younger adults.

5.2. Indicators of zinc requirements of children

No specific indicators of zinc requirements are available for older infants and children. Linear growth is affected by zinc deficiency, but is far from being a specific indicator.

5.3. Indicators of zinc requirements in pregnancy and lactation

Although a variety of clinical features have been linked to zinc deficiency in pregnancy, these features are non-specific and have not been adequately substantiated.

The additional need for zinc during pregnancy can be calculated from the weight of tissues gained during gestation and the concentration of zinc in those tissues. Widdowson and Dickerson (1964) measured the concentration of zinc in 24 human fetuses and full-term infants ranging in weight from 0.75 to about 4 400 g. Using a mean measured zinc concentration of 18.4 µg/g fat-free tissue and measured fetal growth rates, Shaw (1979) calculated the rate of zinc accumulation by a human fetus growing along the 10th, 50th or 90th percentiles. The zinc accumulation rate for a fetus growing along the 50th percentile increased progressively from 0.21 mg/day at the 24th week of gestation to 0.67 mg/day at the 36th week. In addition to the fetus, placental, uterine and mammary tissue, amniotic fluid and maternal blood are also gained during gestation. Hytten (1980) calculated the total weight of the pregnancy tissues at term. Based on the total weight of those tissues and their zinc concentrations, the total zinc requirement for pregnancy has been calculated to be about 100 mg (Swanson and King, 1987). Approximately 60 % of the gain in zinc is associated with the fetus. The daily requirement for zinc in pregnancy above that of non-pregnant women can be calculated from the rate of tissue gain and the tissue zinc concentrations. The daily rates of zinc accumulation for the four quarters of pregnancy have been estimated to be 0.08, 0.24, 0.53 and 0.73 mg (Swanson and King, 1987). Taking into account the cessation of zinc losses with menstruation (equivalent to about 0.01 mg/day in menstruating women), Swanson and King (1987) estimated an additional physiological requirement for zinc in the second half of pregnancy of about 0.6 mg/day. When the average daily rates of zinc

accumulation for the four quarters of pregnancy (Swanson and King, 1987) are calculated, an additional physiological requirement of about 0.4 mg/day for the whole pregnancy is determined.

In lactation, additional zinc may be needed to replace zinc secreted in breast milk. Losses of zinc in breast milk have been estimated, taking into account milk zinc concentrations and the amount of milk transferred, and are 2.52 mg/day for the first month, 1.37 mg/day for months 1 to 2 and 0.86 mg/day for months 3 to < 6 post partum (Brown et al., 2009) (see Section 2.3.6.3). Estimations for the additional zinc requirement in lactation also need to take into account redistribution of tissue zinc during postnatal re-adaptation to the non-pregnant state. Post partum involution of the uterus and decreased maternal blood volume have been estimated to release about 30 mg of zinc that has accumulated during pregnancy (King and Turnlund, 1989). The Panel assumed that this endogenous zinc is available for re-utilisation and decreases the additional amount of zinc required during the first month of lactation by 1 mg/day. This would reduce the additional physiological requirement to about 1.5 mg/day for the first month of lactation. It has also been postulated that bone resorption in early lactation contributes to the amount of endogenous zinc available for secretion in breast milk (Moser-Veillon, 1995; WHO, 1996), although the amount of zinc released from maternal bone during lactation has not been quantified (Donangelo and King, 2012).

In pregnancy and, notably, in early lactation, up-regulation of zinc absorption has been reported (Fung et al., 1997; Harvey et al., 2007; Donangelo and King, 2012). For example, in two longitudinal studies of zinc homeostasis during pregnancy and lactation, FAZ increased 1.3-fold ($p > 0.05$) from pre-conception to late pregnancy in 13 US women with a zinc intake of about 12 mg/day (Fung et al., 1997) and 1.5-fold ($p < 0.05$) from early (10–12 weeks) to late (34–36 weeks) pregnancy in 10 Brazilian women ingesting about 9 mg/day (Donangelo et al., 2005). FAZ increased 1.7-fold ($p = 0.023$) from pre-conception to lactation in the US women (Fung et al., 1997) and 1.4-fold ($p < 0.05$) from early pregnancy to lactation in the Brazilian women (Donangelo et al., 2005). There is some evidence to indicate that this up-regulation of zinc absorption may be sufficient to match increased requirements (Hambidge KM et al., unpublished).

5.4. Zinc intake and long-term health consequences

Mild to moderate dietary zinc depletion is a cause of several non-specific features including growth retardation, depressed immune function with susceptibility to infections, delayed wound healing, loss of appetite and loss of cognitive function. Severe restriction of dietary zinc is a cause of other clinical features including skin rashes. However, clinical features are non-specific and cannot be used for estimating DRVs. A systematic literature search and review for studies addressing zinc intake and health relationships was done by EURRECA; many studies were retrieved that addressed the relationship between zinc intake and outcomes such as cognitive and immune function, depression, anorexia, diabetes mellitus, ischaemic heart disease and cancer in adults. The authors concluded that studies were heterogeneous in their methodological approaches and outcomes assessed (Lowe et al., 2013). The Panel concludes that the available evidence on zinc intake and health outcomes cannot be used for setting DRVs for zinc.

6. Data on which to base Dietary Reference Values

The data required to derive ARs and PRIs in different population groups are the zinc intake needed to replace endogenous losses and the quantity needed for growth and lactation, where appropriate. The factorial approach for deriving DRVs for zinc is used for all age groups.

6.1. Adults

As dietary zinc requirement depends on body weight and dietary phytate intake (Sections 5.1.1 and 5.1.2), the Panel considers it appropriate to estimate ARs and PRIs for the range of mean/median dietary phytate intake observed in Europe. Thus, estimated ARs and PRIs are provided for phytate intake levels of 300, 600, 900 and 1 200 mg/day, which cover the range of mean/median phytate intake observed in European populations (see Section 3.3 and Appendix G) and thus reflect the variety

of European dietary patterns. Where population data on phytate intakes are available, ARs and PRIs could subsequently be adjusted using well-validated statistical models, an example of which has been used in this Opinion.

Table 10 contains estimates on ARs and PRIs for zinc based on reference body weights for a BMI of 22 kg/m² (see Appendix 11 in EFSA NDA Panel (2013)). PRIs for adults were estimated as the zinc requirement of individuals with a body weight at the 97.5th percentile for reference body weights for men and women, as body weight is a strong determinant of the requirement for zinc and as this approach is considered to have less uncertainty than the mathematical application of a CV of between 10 and 20 %. The PRIs based on the 97.5th percentile for reference body weights are equivalent to CVs for the ARs of between 10 and 14 %.

Table 10: Estimations of Average Requirements and Population Reference Intakes for zinc according to phytate intake and body weight

Level of phytate intake (mg/day)	Body weight (kg)	Average Requirement (mg/day)	Population Reference Intake (mg/day) ^(a)
300	58.5 ^(b)	6.2	7.5
	68.1 ^(c)	7.5	9.4
600	58.5 ^(b)	7.6	9.3
	68.1 ^(c)	9.3	11.7
900	58.5 ^(b)	8.9	11.0
	68.1 ^(c)	11.0	14.0
1 200	58.5 ^(b)	10.2	12.7
	68.1 ^(c)	12.7	16.3

(a): Dietary zinc intake of subjects with a body weight at the 97.5th percentile of the reference body weights (i.e. 79.4 kg for men and 68.1 kg for women).

(b): Median body weight of 18- to 79-year-old women based on measured body heights of 19 969 women in 13 European Union Member States and assuming a body mass index of 22 kg/m² (see Appendix 11 in EFSA NDA Panel (2013)). At this body weight, the physiological zinc requirement is 2.9 mg/day.

(c): Median body weight of 18- to 79-year-old men based on measured body heights of 16 500 men in 13 European Union Member States and assuming a body mass index of 22 kg/m² (see Appendix 11 in EFSA NDA Panel (2013)). At this body weight, the physiological zinc requirement is 3.2 mg/day.

6.2. Infants and children

Estimation of DRVs for zinc uses a factorial approach taking into account endogenous zinc losses via urine, sweat and integument, faeces and, in adolescent boys and girls, semen and menses, respectively, as well as zinc required for the synthesis of new tissue for growth.

6.2.1. Methodology

6.2.1.1. Urinary and integumental zinc losses

After early infancy, urinary excretion rates for children on a body weight basis seem to differ very little from adult values (Krebs and Hambidge, 1986). Thus, for infants aged 7 to 11 months and children from one year of age, data for urinary losses were extrapolated from adult values (see Section 2.3.6.2) using isometric scaling, i.e. linear with body weight. For this, reference body weights based on the WHO Multicentre Growth Reference Study Group (2006) were used for infants and young children and based on van Buuren et al. (2012) for older children. For the age classes shown in Table 10, median body weights at mid-point ages were chosen, i.e. at age nine months and 2, 5, 8.5, 12.5 and 16 years.

Integumental zinc losses were estimated from adult values (see Section 2.3.6.2) using allometric scaling, i.e. body weight to the power of 0.67 as a proxy for body surface area.

6.2.1.2. Endogenous faecal zinc losses

In infants aged two to four months, the average intestinal excretion of endogenous zinc in exclusively breast-fed infants was approximately 50 µg/kg body weight per day (Krebs et al., 1996). For infants receiving complementary foods in addition to infant formula or human milk, endogenous faecal losses of 40 µg/kg body weight per day were assumed by WHO (1996) and WHO/FAO (2004). This figure was thus used to calculate daily EFZ by multiplication with infants' reference body weights.

Linear regression analysis of EFZ versus body weight (kg) for the subjects contributing data to the adult estimates (see Section 5.1.1), for 43 young children aged 19 to 25 months from China (Sheng et al., 2006) and from a study in 45 infants aged 9 to 10 months in the USA (Krebs N et al., unpublished), gives the following equation:

$$\text{EFZ [mg Zn/day]} = 0.0318 \times \text{body weight [kg]} + 0.362, \text{ with } R^2 = 0.75 \quad (\text{Equation 4})$$

This equation was used to estimate EFZ for children from one year of age.

6.2.1.3. Zinc losses in menses and semen

Zinc losses in menses and semen of 0.01 and 0.1 mg/day, respectively, have been assumed (see Section 5.1.1.4). The mean age of menarche in the European Union (EU) has been reported to be 12.7 years (van Buuren et al., 2012); thus, menstrual zinc losses have been taken into account for the 11- to 14-year-old group of girls, whereas zinc losses in semen have been assumed for boys from 15 years of age and older.

6.2.1.4. Zinc requirement for growth

For the estimation of zinc requirement for tissue gain, a figure of 30 µg/g of new tissue has previously been used for infants and children, whereas, for adolescents, a zinc content of 23 µg/g wet weight has been assumed due to an increase in fat tissue with a lower zinc content than that in younger children (WHO, 1996). Analyses of whole fetuses of various gestational ages have shown a constant zinc content of about 20 µg/g of fat-free tissue (Widdowson and Spray, 1951), and this value has also been used in factorial estimates (IOM, 2001). The Panel considers that, in the absence of direct and precise data on body composition of infants and children at various postnatal ages, a figure of 20 µg zinc/g tissue gained appears to be a reasonable estimate. This value was multiplied with daily weight gains of the age groups. Daily weight gains of infants in the second half-year of life were assumed to be 11.5 g/day, based on observed weight increments of infants in the Euro-Growth Study, where median weight gain of boys and girls was 13 g/day from month 6 to 9 and 10 g/day from month 9 to 12 of age (van't Hof et al., 2000). For children, daily weight gains were calculated by subtracting the median weight at the lower boundary of the age group from that at the higher boundary of the age group and dividing this by the number of days in that age interval, assuming that one year equals 365 days.

6.2.1.5. Fractional absorption of zinc

The IOM (2001) used a figure of 0.30 for the FAZ for older infants and children based on the literature available at that time. The Panel considers that subsequent data (Manary et al., 2000; Griffin et al., 2004; Lopez de Romana et al., 2005; Mazariegos et al., 2006; Sheng et al., 2006; Griffin et al., 2007) provide no reason to modify this figure. As this figure is based on mixed diets that probably contain variable quantities of phytate, no adjustment for phytate intake has been made.

6.2.2. Infants aged 7 to 11 months

Using a mean weight-for-age of 8.6 kg for boys and girls aged nine months at the 50th percentile (WHO Multicentre Growth Reference Study Group, 2006), endogenous faecal zinc losses were estimated as 0.343 mg/day (see Section 6.2.1.2) and urine losses were estimated as 0.054 mg/day, extrapolated from adult values (see Section 6.2.1.1). No data are available on integumental zinc losses in infants which have, therefore, been extrapolated from integumental losses in adults, as described in

Section 6.2.1.1, giving a figure of 0.105 mg/day. Estimated total endogenous zinc losses are 0.502 mg/day.

Based on an average weight gain of 11.5 g/day for infants in the second half-year of life, the estimated zinc requirement for growth is 0.230 mg/day.

Therefore, the estimated physiological zinc requirement for infants aged 7 to 11 months is 0.732 mg/day.

Assuming a FAZ of 0.3 (see Section 6.2.1.5), the AR for infants aged 7 to 11 months is 2.4 mg/day. Owing to the absence of reference body weights for infants at the 97.5th percentile, and with no knowledge about the variation in requirement, the PRI for infants was estimated based on a CV of 10 %, and is 2.9 mg/day.

6.2.3. Children

Components that were considered for the factorial approach for the various age groups are listed in Table 11. The number of digits used for the calculations has been retained in the table, with the exception of the AR, for which an erroneous impression of accuracy would be given.

Table 11: Estimates used in the factorial approach to derive the Average Requirements for zinc for children

Age	Reference weight (kg)	Zinc losses (mg/day) ^(a) via					Estimated daily weight gain (g/day) ^(b)	Zinc need for growth (mg/day)	Physiological requirement (mg/day) ^(c)	Average Requirement (mg/day) ^(d)
		Faeces	Urine	Sweat	Semen	Menses				
1–3 years	11.9 ^(e)	0.738	0.075	0.130	–	–	6.57	0.131	1.074	3.6
4–6 years	19.0 ^(f)	0.965	0.120	0.178	–	–	6.35	0.127	1.390	4.6
7–10 years	28.7 ^(g)	1.275	0.181	0.236	–	–	8.82	0.176	1.869	6.2
11–14 years (M)	44.0 ^(h)	1.762	0.278	0.314	–	–	14.1	0.282	2.635	8.8
11–14 years (F)	45.1 ⁽ⁱ⁾	1.797	0.285	0.319	–	0.01	12.6	0.252	2.663	8.9
15–17 years (M)	64.1 ^(j)	2.401	0.405	0.403	0.1	–	11.7	0.235	3.544	11.8
15–17 years (F)	56.4 ^(k)	2.157	0.357	0.370	–	0.01	3.78	0.076	2.969	9.9

M, males; F, females.

(a): See Sections 6.2.1.1 and 6.2.1.2.

(b): See Section 6.2.1.3.

(c): Sum of losses and need for growth.

(d): Estimated from the physiological requirement and assuming an absorption efficiency of 30 % from a mixed diet (see Section 6.2.1.5); values were rounded to the nearest 0.1.

(e): Mean body weight-for-age at 50th percentile of boys and girls aged 24 months (WHO Multicentre Growth Reference Study Group, 2006).

(f): Mean body weight at 50th percentile of boys and girls aged five years (van Buuren et al., 2012).

(g): Mean body weight at 50th percentile of boys and girls aged 8.5 years (van Buuren et al., 2012).

(h): Body weight at 50th percentile of boys aged 12.5 years (van Buuren et al., 2012).

(i): Body weight at 50th percentile of girls aged 12.5 years (van Buuren et al., 2012).

(j): Body weight at 50th percentile of boys aged 16 years (van Buuren et al., 2012).

(k): Body weight at 50th percentile of girls aged 16 years (van Buuren et al., 2012).

Due to the absence of reference body weights for infants and children at the 97.5th percentile, and with no knowledge about the variation in requirement, PRIs for infants and children were estimated based on a CV of 10 %. Table 12 contains estimates on ARs and PRIs for zinc for infants and children.

Table 12: Summary of Average Requirements and Population Reference Intakes for zinc for infants and children

Age	Average Requirement (mg/day)	Population Reference Intake (mg/day)
7–11 months	2.4	2.9
1–3 years	3.6	4.3
4–6 years	4.6	5.5
7–10 years	6.2	7.4
11–14 years	8.9	10.7
15–17 years (M)	11.8	14.2
15–17 years (F)	9.9	11.9

M, males; F, females.

6.3. Pregnancy and lactation

Despite some evidence of up-regulation of zinc absorption during pregnancy and notably during early lactation (see Section 5.3) and evidence from one unpublished study that this may be sufficient to meet increased requirements, the Panel considers that data are insufficient to modify estimated additional physiological requirements. The most reliable indicators of zinc requirements at present are the addition of the estimated daily increment for pregnancy and the quantity of zinc secreted in milk over the first six months of lactation, adjusted for re-absorption of zinc owing to redistribution of tissue zinc during postnatal re-adaptation to the non-pregnant state.

The additional requirements for pregnancy and lactation may be calculated by estimating the additional physiological requirement for synthesis of new tissue, primarily the conceptus, and for replacement of zinc secreted in breast milk (see Section 5.3).

For pregnancy, an additional physiological requirement of about 0.4 mg/day may be calculated for the whole pregnancy (see Section 5.3). This combined estimate probably overestimates the requirement in the first half of pregnancy and underestimates the requirement in the second half of pregnancy. It is unknown if the trivariate model used to estimate dietary zinc requirements of non-pregnant non-lactating women is also suitable in pregnancy and lactation, and up-regulation of zinc absorption is likely to modify the inhibitory effect of phytate on zinc absorption. Thus, the Panel decided not to use the trivariate model to estimate the dietary zinc intake required to meet the additional physiological requirement. Instead, the Panel applied a mean FAZ of 0.30 observed in healthy adults (see Appendix I) to the physiological requirement of 0.4 mg/day and estimated the additional average dietary zinc requirement in pregnancy to be 1.3 mg/day. In the absence of knowledge about the variation in requirement, the additional PRI for pregnancy was estimated based on a CV of 10 %, and was 1.6 mg/day.

For lactation, the Panel assumed that the mean increases in physiological requirement are 1.5 mg/day for the first month, 1.37 mg/day for months 1 to 2 and 0.86 mg/day for months 3 to 6 post partum (see Section 5.3). When the average over six months of lactation is calculated, an additional physiological requirement of 1.1 mg/day is determined. Assuming that FAZ is increased 1.5-fold in lactation (see Section 5.3), and applying a FAZ of 0.45 to the additional physiological requirement of 1.1 mg/day, an additional average dietary zinc requirement in lactation of 2.4 mg/day is determined. In the absence of knowledge about the variation in requirement, the additional PRI for lactation was estimated based on a CV of 10 %, and was 2.9 mg/day.

CONCLUSIONS

The Panel concludes that ARs and PRIs for zinc can be derived for adults based on a two-stage factorial approach. The first stage comprised the estimation of physiological requirement, defined as the minimum quantity of absorbed zinc necessary to match losses of endogenous zinc, and its relationship with body weight. In the second stage, the quantity of dietary zinc available for absorption that is necessary to meet the physiological requirement was determined, taking into account the inhibitory effect of phytate on zinc absorption. ARs for adults were estimated as the zinc requirement at the 50th percentile of reference body weights for men and women in the EU and for levels of phytate intake of 300, 600, 900 and 1 200 mg/day, and PRIs for adults were estimated as the zinc requirement of individuals with a body weight at the 97.5th percentile for reference body weights for men and women, and for the same range of phytate intake. For infants and children, ARs were estimated based on factorial calculation of losses and an estimation of the need for growth. For pregnant and lactating women, the increase in physiological requirement was estimated based on the demand for new tissue, primarily by the conceptus, and on the provision of zinc secreted in breast milk, respectively. ARs were derived taking into account the FAZ. In the absence of knowledge about the variation in requirement, PRIs for infants and children and PRIs to cover the additional requirement of pregnant and lactating women were estimated based on a CV of 10 %.

Table 13: Summary of Population Reference Intakes for zinc

	Level of phytate intake (mg/day)	Population Reference Intake (mg/day)
Age		
7–11 months		2.9
1–3 years		4.3
4–6 years		5.5
7–10 years		7.4
11–14 years		9.4
15–17 years (M)		12.5
15–17 years (F)		10.4
≥ 18 years (M)	300	9.4
	600	11.7
	900	14.0
	1 200	16.3
≥ 18 years (F)	300	7.5
	600	9.3
	900	11.0
	1 200	12.7
Pregnancy		+ 1.6
Lactation		+ 2.9

M, males; F, females.

RECOMMENDATIONS FOR RESEARCH

The Panel suggests that studies of zinc homeostasis in European populations should be undertaken using state of the art techniques, and should target the more vulnerable populations such as young children, adolescents, pregnant and lactating women and the elderly.

The Panel recommends that additional reliable data on phytate intake in the EU be collected.

The Panel recommends that studies be undertaken to identify suitable biomarkers of zinc status and also recommends that methods to derive zinc requirements be further refined.

REFERENCES

- Adams CL, Hambidge M, Raboy V, Dorsch JA, Sian L, Westcott JL and Krebs NF, 2002. Zinc absorption from a low-phytic acid maize. *American Journal of Clinical Nutrition*, 76, 556-559.
- Afssa (Agence française de sécurité sanitaire des aliments), 2001. Apports nutritionnels conseillés pour la population française. Editions Tec&Doc, Paris, France, 605 pp.
- Amcoff E, Edberg A, Enghardt Barbieri H, Lindroos A, Nälsén C, Pearson M and Warensjö Lemming E (Livsmedelsverket), 2012. Riksmaten – vuxna 2010–11. Livsmedels- och näringsintag bland vuxna i Sverige. Resultat från matvaneundersökning utförd 2010–11. 180 pp.
- Amirabdollahian F and Ash R, 2010. An estimate of phytate intake and molar ratio of phytate to zinc in the diet of the people in the United Kingdom. *Public Health Nutrition*, 13, 1380-1388.
- Avellini BA, Kamon E and Krajewski JT, 1980. Physiological responses of physically fit men and women to acclimation to humid heat. *Journal of Applied Physiology: Respiratory, Environmental and Exercise Physiology*, 49, 254-261.
- Bates B, Lennox A, Prentice A, Bates C and Swan G, 2012. National Diet and Nutrition Survey. Headline results from Years 1, 2 and 3 (combined) of the Rolling Programme (2008/2009 – 2010/11). A survey carried out on behalf of the Department of Health and the Food Standards Agency. 79 pp.
- Bates CJ and Tsuchiya H, 1990. Zinc in breast milk during prolonged lactation: comparison between the UK and the Gambia. *European Journal of Clinical Nutrition*, 44, 61-69.
- Beck FW, Prasad AS, Kaplan J, Fitzgerald JT and Brewer GJ, 1997a. Changes in cytokine production and T cell subpopulations in experimentally induced zinc-deficient humans. *American Journal of Physiology*, 272, E1002-1007.
- Beck FW, Kaplan J, Fine N, Handschu W and Prasad AS, 1997b. Decreased expression of CD73 (ecto-5'-nucleotidase) in the CD8+ subset is associated with zinc deficiency in human patients. *Journal of Laboratory and Clinical Medicine*, 130, 147-156.
- Bieri A, Petkovic V, Miletta MC, Fluck CE and Mullis PE, 2013. Short stature and GH deficiency caused by zinc deficiency in a 6-month-old breast-fed infant due to a dominant negative heterozygous G87R mutation in the Zinc Transporter ZnT-2 (SLC30A2) in the mother. *Hormone Research in Paediatrics*, 80, 154-155.
- Bindra GS and Gibson RS, 1986. Mineral intakes of predominantly lacto-ovo vegetarian East Indian adults. *Biological Trace Element Research*, 10, 223-234.
- Bjorklund KL, Vahter M, Palm B, Grander M, Lignell S and Berglund M, 2012. Metals and trace element concentrations in breast milk of first time healthy mothers: A biological monitoring study. *Environmental Health: A Global Access Science Source*, 11.
- Black RE, 2003. Zinc deficiency, infectious disease and mortality in the developing world. *Journal of Nutrition*, 133, 1485S-1489S.
- Brown KH, Engle-Stone R, Krebs NF and Peerson JM, 2009. Dietary intervention strategies to enhance zinc nutrition: promotion and support of breastfeeding for infants and young children. *Food and Nutrition Bulletin*, 30, S144-171.
- Brune M, Rossander L and Hallberg L, 1989. Iron absorption: no intestinal adaptation to a high-phytate diet. *American Journal of Clinical Nutrition*, 49, 542-545.
- Carnovale E, Lombardi-Boccia G and Lugaro E, 1987. Phytate and zinc content of Italian diets. *Human Nutrition. Applied Nutrition*, 41, 180-186.
- Casey CE, Hambidge KM and Neville MC, 1985. Studies in human lactation: zinc, copper, manganese and chromium in human milk in the first month of lactation. *American Journal of Clinical Nutrition*, 41, 1193-1200.

- Casey CE, Neville MC and Hambidge KM, 1989. Studies in human lactation: secretion of zinc, copper, and manganese in human milk. *American Journal of Clinical Nutrition*, 49, 773-785.
- Cheryan M, 1980. Phytic acid interactions in food systems. *Critical Reviews in Food Science and Nutrition*, 13, 297-335.
- Chierici R, Saccomandi D and Vigi V, 1999. Dietary supplements for the lactating mother: influence on the trace element content of milk. *Acta Paediatrica. Supplement*, 88, 7-13.
- Chowanadisai W, Lonnerdal B and Kelleher SL, 2006. Identification of a mutation in SLC30A2 (ZnT-2) in women with low milk zinc concentration that results in transient neonatal zinc deficiency. *Journal of Biological Chemistry*, 281, 39699-39707.
- Chung CS, Stookey J, Dare D, Welch R, Nguyen TQ, Roehl R, Peerson JM, King JC and Brown KH, 2008. Current dietary zinc intake has a greater effect on fractional zinc absorption than does longer term zinc consumption in healthy adult men. *American Journal of Clinical Nutrition*, 87, 1224-1229.
- Coelho S, Fernandes B, Rodrigues F, Reis JP, Moreno A and Figueiredo A, 2006. Transient zinc deficiency in a breast fed, premature infant. *European Journal of Dermatology*, 16, 193-195.
- Cohn JR and Emmett EA, 1978. The excretion of trace metals in human sweat. *Annals of Clinical and Laboratory Science*, 8, 270-275.
- D-A-CH (Deutsche Gesellschaft für Ernährung, Österreichische Gesellschaft für Ernährung, Schweizerische Gesellschaft für Ernährungsforschung, Schweizerische Vereinigung für Ernährung), 2013. Referenzwerte für die Nährstoffzufuhr. Neuer Umschau Buchverlag, Neustadt an der Weinstraße, Germany, 292 pp.
- Davidsson L, Mackenzie J, Kastenmayer P, Aggett PJ and Hurrell RF, 1996. Zinc and calcium apparent absorption from an infant cereal: a stable isotope study in healthy infants. *British Journal of Nutrition*, 75, 291-300.
- de Benoist B, Darnton-Hill I, Davidsson L, Fontaine O and Hotz C, 2007. Conclusions of the Joint WHO/UNICEF/IAEA/IZiNCG Interagency Meeting on Zinc Status Indicators. *Food and Nutrition Bulletin*, 28, S480-484.
- DeRuisseau KC, Chevront SN, Haymes EM and Sharp RG, 2002. Sweat iron and zinc losses during prolonged exercise. *International Journal of Sport Nutrition and Exercise Metabolism*, 12, 428-437.
- DH (Department of Health), 1991. Dietary reference values for food energy and nutrients for the United Kingdom. Report of the Panel on Dietary Reference Values of the Committee on Medical Aspects of Food Policy. HM Stationary Office, London, UK, 212 pp.
- Domellof M, Hernell O, Dewey KG, Cohen RJ and Lonnerdal B, 2004. Factors influencing concentrations of iron, zinc, and copper in human milk. *Advances in Experimental Medicine and Biology*, 554, 355-358.
- Donangelo CM, Zapata CL, Woodhouse LR, Shames DM, Mukherjea R and King JC, 2005. Zinc absorption and kinetics during pregnancy and lactation in Brazilian women. *American Journal of Clinical Nutrition*, 82, 118-124.
- Donangelo CM and King JC, 2012. Maternal zinc intakes and homeostatic adjustments during pregnancy and lactation. *Nutrients*, 4, 782-798.
- EFSA (European Food Safety Authority), 2011a. Report on the development of a food classification and description system for exposure assessment and guidance on its implementation and use. *EFSA Journal* 2011;9(12):2489, 84 pp. doi:10.2903/j.efsa.2011.2489
- EFSA (European Food Safety Authority), 2011b. Use of the EFSA Comprehensive European Food Consumption Database in Exposure Assessment. *EFSA Journal* 2011;9(3):2097, 34 pp. doi:10.2903/j.efsa.2011.2097

- EFSA NDA Panel (EFSA Panel on Dietetic Products Nutrition and Allergies), 2012. Scientific Opinion on Dietary Reference Values for protein. *EFSA Journal* 2012;10(2):2557, 66 pp. doi:10.2903/j.efsa.2012.2557
- EFSA NDA Panel (EFSA Panel on Dietetic Products, Nutrition and Allergies), 2013. Scientific Opinion on Dietary Reference Values for energy. *EFSA Journal* 2013;11(1):3005, 112 pp. doi:10.2903/j.efsa.2013.3005
- Ellis R, Kelsay JL, Reynolds RD, Morris ER, Moser PB and Frazier CW, 1987. Phytate:zinc and phytate X calcium:zinc millimolar ratios in self-selected diets of Americans, Asian Indians, and Nepalese. *Journal of the American Dietetic Association*, 87, 1043-1047.
- Elmastas M, Can M, Uzun S and Aboul-Enein HY, 2005. Determinations of copper, zinc, cadmium, and nickel in cows', goats', ewes', and human milk samples using flame atomic absorption spectrometry (FAAS) microwave digestion. *Analytical Letters*, 38, 157-165.
- Fairweather-Tait SJ, Wharf SG and Fox TE, 1995. Zinc absorption in infants fed iron-fortified weaning food. *American Journal of Clinical Nutrition*, 62, 785-789.
- Frye AJ and Kamon E, 1983. Sweating efficiency in acclimated men and women exercising in humid and dry heat. *Journal of Applied Physiology: Respiratory, Environmental and Exercise Physiology*, 54, 972-977.
- Fung EB, Ritchie LD, Woodhouse LR, Roehl R and King JC, 1997. Zinc absorption in women during pregnancy and lactation: a longitudinal study. *American Journal of Clinical Nutrition*, 66, 80-88.
- Gass JK, Hafiji J and Burrows NP, 2010. Acrodermatitis due to zinc-deficient breast milk. *British Journal of Dermatology*, 162, 1164-1165.
- Gehan EA and George SL, 1970. Estimation of human body surface area from height and weight. *Cancer Chemotherapy Reports. Part 1*, 54, 225-235.
- Gibson RS, Vanderkooy PD, MacDonald AC, Goldman A, Ryan BA and Berry M, 1989. A growth-limiting, mild zinc-deficiency syndrome in some southern Ontario boys with low height percentiles. *American Journal of Clinical Nutrition*, 49, 1266-1273.
- Gibson RS, Heywood A, Yaman C, Sohlstrom A, Thompson LU and Heywood P, 1991. Growth in children from the Wosera subdistrict, Papua New Guinea, in relation to energy and protein intakes and zinc status. *American Journal of Clinical Nutrition*, 53, 782-789.
- Gibson RS, Heath AL, Prosser N, Parnell W, Donovan UM, Green T, McLaughlin KE, O'Connor DL, Bettger W and Skeaff CM, 2000. Are young women with low iron stores at risk of zinc as well as iron deficiency? In: *Trace Elements in Man and Animals*. Eds Roussel AM, Anderson RA and Favrier A. Kluwer Academic, New York, USA, 323-328.
- Grider A, Bailey LB and Cousins RJ, 1990. Erythrocyte metallothionein as an index of zinc status in humans. *Proceedings of the National Academy of Sciences of the United States of America*, 87, 1259-1262.
- Griffin IJ, Hicks PD, Liang LK and Abrams SA, 2004. Metabolic adaptations to low zinc intakes in premenarcheal girls. *American Journal of Clinical Nutrition*, 80, 385-390.
- Griffin IJ, Lynch MF, Hawthorne KM, Chen Z, Hamzo MG and Abrams SA, 2007. Zinc homeostasis in 1-4 year olds consuming diets typical of US children. *British Journal of Nutrition*, 98, 358-363.
- Guo L, Lichten LA, Ryu MS, Liuzzi JP, Wang F and Cousins RJ, 2010. STAT5-glucocorticoid receptor interaction and MTF-1 regulate the expression of ZnT2 (Slc30a2) in pancreatic acinar cells. *Proceedings of the National Academy of Sciences of the United States of America*, 107, 2818-2823.
- Hambidge K, Casey C and Krebs N, 1986. Zinc. In: *Human and animal nutrition*. Ed Mertz W. Academic Press, New York, USA, 1-137.

- Hambidge KM, Hambidge C, Jacobs M and Baum JD, 1972. Low levels of zinc in hair, anorexia, poor growth, and hypogeusia in children. *Pediatric Research*, 6, 868-874.
- Hambidge KM, Huffer JW, Raboy V, Grunwald GK, Westcott JL, Sian L, Miller LV, Dorsch JA and Krebs NF, 2004. Zinc absorption from low-phytate hybrids of maize and their wild-type isohybrids. *American Journal of Clinical Nutrition*, 79, 1053-1059.
- Hambidge KM, Miller LV, Tran CD and Krebs NF, 2005. Measurements of zinc absorption: application and interpretation in research designed to improve human zinc nutriture. *International Journal for Vitamin and Nutrition Research*, 75, 385-393.
- Hambidge KM, Miller LV, Westcott JE, Sheng X and Krebs NF, 2010. Zinc bioavailability and homeostasis. *American Journal of Clinical Nutrition*, 91, 1478S-1483S.
- Harland BF and Oberleas D, 1977. A modified method for phytate analysis using an ion exchange procedure: application to textured vegetable proteins. *Cereal Chem*, 54, 827-832.
- Harland BF and Oberleas D, 1986. Anion-exchange method for determination of phytate in foods: collaborative study. *Journal of the Association of Official Analytical Chemists*, 69, 667-670.
- Harvey LJ, Dainty JR, Hollands WJ, Bull VJ, Hoogewerff JA, Foxall RJ, McAnena L, Strain JJ and Fairweather-Tait SJ, 2007. Effect of high-dose iron supplements on fractional zinc absorption and status in pregnant women. *American Journal of Clinical Nutrition*, 85, 131-136.
- Hazelhurst LT and Claassen N, 2006. Gender differences in the sweat response during spinning exercise. *Journal of Strength and Conditioning Research*, 20, 723-724.
- Heath AL, Roe MA, Oyston SL and Fairweather-Tait SJ, 2005. Meal-based intake assessment tool: relative validity when determining dietary intake of Fe and Zn and selected absorption modifiers in UK men. *British Journal of Nutrition*, 93, 403-416.
- Hedera P, Peltier A, Fink JK, Wilcock S, London Z and Brewer GJ, 2009. Myelopolyneuropathy and pancytopenia due to copper deficiency and high zinc levels of unknown origin II. The denture cream is a primary source of excessive zinc. *Neurotoxicology*, 30, 996-999.
- Helldán A, Raulio S, Kosola M, Tapanainen H, Ovaskainen ML and Virtanen S, 2013. Finravinto 2012 -tutkimus - The National FINDIET 2012 Survey. THL. Raportti 16/2013, 217 pp.
- Hess FM, King JC and Margen S, 1977. Zinc excretion in young women on low zinc intakes and oral contraceptive agents. *Journal of Nutrition*, 107, 1610-1620.
- Hoppu U, Lehtisalo J, Tapanainen H and Pietinen P, 2010. Dietary habits and nutrient intake of Finnish adolescents. *Public Health Nutrition*, 13, 965-972.
- Hunt CD, Johnson PE, Herbel J and Mullen LK, 1992. Effects of dietary zinc depletion on seminal volume and zinc loss, serum testosterone concentrations, and sperm morphology in young men. *American Journal of Clinical Nutrition*, 56, 148-157.
- Hunt J, Mullen L and Lykken G, 1992. Zinc retention from an experimental diet based on the FDA Total Diet Study. *Nutrition Research*, 1335-1344.
- Hunt JR, Gallagher SK, Johnson LK and Lykken GI, 1995. High- versus low-meat diets: effects on zinc absorption, iron status, and calcium, copper, iron, magnesium, manganese, nitrogen, phosphorus, and zinc balance in postmenopausal women. *American Journal of Clinical Nutrition*, 62, 621-632.
- Hunt JR, Matthys LA and Johnson LK, 1998. Zinc absorption, mineral balance, and blood lipids in women consuming controlled lactoovo-vegetarian and omnivorous diets for 8 wk. *American Journal of Clinical Nutrition*, 67, 421-430.
- Hunt JR, Beiseigel JM and Johnson LK, 2008. Adaptation in human zinc absorption as influenced by dietary zinc and bioavailability. *American Journal of Clinical Nutrition*, 87, 1336-1345.

- Hunt JR and Beiseigel JM, 2009. Dietary calcium does not exacerbate phytate inhibition of zinc absorption by women from conventional diets. *American Journal of Clinical Nutrition*, 89, 839-843.
- Hytten FE, 1980. Weight gain in pregnancy. In: *Clinical Physiology in Obstetrics. Part 2. Nutrition and Metabolism*. Eds Hytten FE and Chamberlain G. Blackwell Scientific Publications, Oxford, UK, 193-233.
- Ingelmann H-J, Rimbach G and Pallauf J, 1993. Phytinsäure - ein antinutritiver Faktor? *Ernährungs-Umschau*, 40, 400-404.
- IOM (Institute of Medicine), 2001. *Dietary Reference Intakes for vitamin A, vitamin K, arsenic, boron, chromium, copper, iodine, iron, manganese, molybdenum, nickel, silicon, vanadium and zinc*. National Academies Press, Washington DC, USA, 797 pp.
- IUNA (Irish Universities Nutrition Alliance), 2011. *National Adult Nutrition Survey*. 40 pp.
- Jackson MJ, Jones DA, Edwards RH, Swainbank IG and Coleman ML, 1984. Zinc homeostasis in man: studies using a new stable isotope-dilution technique. *British Journal of Nutrition*, 51, 199-208.
- Jacob RA, Sandstead HH, Munoz JM, Klevay LM and Milne DB, 1981. Whole body surface loss of trace metals in normal males. *American Journal of Clinical Nutrition*, 34, 1379-1383.
- Johnson PE, Hunt CD, Milne DB and Mullen LK, 1993. Homeostatic control of zinc metabolism in men: zinc excretion and balance in men fed diets low in zinc. *American Journal of Clinical Nutrition*, 57, 557-565.
- Kantola M and Vartiainen T, 2001. Changes in selenium, zinc, copper and cadmium contents in human milk during the time when selenium has been supplemented to fertilizers in Finland. *Journal of Trace Elements in Medicine and Biology*, 15, 11-17.
- Kersting M and Clausen K, 2003. *Ernährungsphysiologische Auswertung einer repräsentativen Verzehrsstudie bei Säuglingen und Kleinkindern VELS mit dem Instrumentarium der DONALD Studie*. Forschungsinstitut für Kinderernährung, Dortmund, Germany, 103 pp.
- Kettunen J, Tukiainen T, Sarin AP, Ortega-Alonso A, Tikkanen E, Lyytikäinen LP, Kangas AJ, Soininen P, Wurtz P, Silander K, Dick DM, Rose RJ, Savolainen MJ, Viikari J, Kahonen M, Lehtimäki T, Pietiläinen KH, Inouye M, McCarthy MI, Jula A, Eriksson J, Raitakari OT, Salomaa V, Kaprio J, Jarvelin MR, Peltonen L, Perola M, Freimer NB, Ala-Korpela M, Palotie A and Ripatti S, 2012. Genome-wide association study identifies multiple loci influencing human serum metabolite levels. *Nature Genetics*, 44, 269-276.
- Kharfi M, Zaraa I, Kury S, Moisan JP and Kamoun MR, 2005. [Acrodermatitis enteropathica in full-term breast-fed infant]. *Annales de Dermatologie et de Vénérologie*, 132, 246-248.
- Khokhar S and Pushpanjali, 1994. Phytate content of Indian foods and intakes by vegetarian Indians of Hisar Region, Haryana State. *Journal of Agricultural and Food Chemistry*, 42, 2440-2444.
- Kim J, Paik HY, Joung H, Woodhouse LR, Li S and King JC, 2007. Effect of dietary phytate on zinc homeostasis in young and elderly Korean women. *Journal of the American College of Nutrition*, 26, 1-9.
- King JC and Turnlund JR, 1989. Human zinc requirements. In: *Zinc in Human Biology*. Ed Mills C. Springer Verlag, London, UK, 335-350.
- King JC, 2000. Determinants of maternal zinc status during pregnancy. *American Journal of Clinical Nutrition*, 71, 1334S-1343S.
- King JC, Shames DM, Lowe NM, Woodhouse LR, Sutherland B, Abrams SA, Turnlund JR and Jackson MJ, 2001. Effect of acute zinc depletion on zinc homeostasis and plasma zinc kinetics in men. *American Journal of Clinical Nutrition*, 74, 116-124.

- King JC, 2011. Zinc: an essential but elusive nutrient. *American Journal of Clinical Nutrition*, 94, 679S-684S.
- King JC and Cousins R, 2014. Zinc. In: *Modern Nutrition in Health and Disease*. Eds Ross AC, Caballero B, Cousins RJ, Tucker KL and Ziegler TR. Lippincott Williams & Wilkins, Philadelphia, USA, 189-205.
- Kirchgessner M, Spörl R and Roth-Maier DA, 1980. [Fecal excretion and apparent absorption of copper, zinc nickel and manganese in nonpregnant and pregnant sows with differing dietary supplies of trace elements]. *Zeitschrift für Tierphysiologie, Tierernährung und Futtermittelkunde*, 44, 98-111.
- Knudsen E, Sandström B and Solgaard P, 1996. Zinc, copper and magnesium absorption from a fibre-rich diet. *Journal of Trace Elements in Medicine and Biology*, 10, 68-76.
- Krebs NF and Hambidge KM, 1986. Zinc requirements and zinc intakes of breast-fed infants. *American Journal of Clinical Nutrition*, 43, 288-292.
- Krebs NF, 1993. Zinc absorption and fecal excretion of endogenous zinc in the breastfed infant. In: *Trace elements in man and animals - Tema 8*. Ed Anke M. Media Verlag Toursistik, Dresden, Germany, 1110-1113.
- Krebs NF, Reidinger CJ, Hartley S, Robertson AD and Hambidge KM, 1995. Zinc supplementation during lactation: effects on maternal status and milk zinc concentrations. *American Journal of Clinical Nutrition*, 61, 1030-1036.
- Krebs NF, Reidinger CJ, Miller LV and Hambidge KM, 1996. Zinc homeostasis in breast-fed infants. *Pediatric Research*, 39, 661-665.
- Krezel A, Hao Q and Maret W, 2007. The zinc/thiolate redox biochemistry of metallothionein and the control of zinc ion fluctuations in cell signaling. *Archives of Biochemistry and Biophysics*, 463, 188-200.
- Kristensen MB, Hels O, Morberg CM, Marving J, Bugel S and Tetens I, 2006. Total zinc absorption in young women, but not fractional zinc absorption, differs between vegetarian and meat-based diets with equal phytic acid content. *British Journal of Nutrition*, 95, 963-967.
- Kuczmariski RJ, Ogden CL, Grummer-Strawn LM, Flegal KM, Guo SS, Wei R, Mei Z, Curtin LR, Roche AF and Johnson CL (National Center for Health Statistics), 2000. *CDC Growth Charts: United States. Advance data from vital health statistics. No. 314. 28 pp.*
- Kyttälä P, Ovaskainen M, Kronberg-Kippilä C, Erkkola M, Tapanainen H, Tuokkola J, Veijola R, Simell O, Knip M and Virtanen S, 2008. *The Diet of Finnish Preschoolers (in Finnish, abstract in English)*. Publications of the National Public Health Institute B32/2008, 154 pp.
- Kyttälä P, Erkkola M, Kronberg-Kippilä C, Tapanainen H, Veijola R, Simell O, Knip M and Virtanen SM, 2010. Food consumption and nutrient intake in Finnish 1-6-year-old children. *Public Health Nutrition*, 13, 947-956.
- Lee DY, Prasad AS, Hydrick-Adair C, Brewer G and Johnson PE, 1993. Homeostasis of zinc in marginal human zinc deficiency: role of absorption and endogenous excretion of zinc. *Journal of Laboratory and Clinical Medicine*, 122, 549-556.
- Lentner C, 1984. *Geigy scientific tables, Vol. 3. Body composition during pregnancy*. CIBA-Geigy, Basel, Switzerland, 359 pp.
- Leotsinidis M, Alexopoulos A and Kostopoulou-Farri E, 2005. Toxic and essential trace elements in human milk from Greek lactating women: Association with dietary habits and other factors. *Chemosphere*, 61, 238-247.
- Leverkus M, Kutt S, Brocker EB, Frank J and Hamm H, 2006. Nutritional zinc deficiency mimicking acrodermatitis enteropathica in a fully breast-fed infant. *Journal of the European Academy of Dermatology and Venereology*, 20, 1380-1381.

- Lopez de Romana D, Salazar M, Hambidge KM, Penny ME, Peerson JM, Krebs NF and Brown KH, 2005. Longitudinal measurements of zinc absorption in Peruvian children consuming wheat products fortified with iron only or iron and 1 of 2 amounts of zinc. *American Journal of Clinical Nutrition*, 81, 637-647.
- Lowe NM, Shames DM, Woodhouse LR, Matel JS, Roehl R, Saccomani MP, Toffolo G, Cobelli C and King JC, 1997. A compartmental model of zinc metabolism in healthy women using oral and intravenous stable isotope tracers. *American Journal of Clinical Nutrition*, 65, 1810-1819.
- Lowe NM, Fekete K and Decsi T, 2009. Methods of assessment of zinc status in humans: a systematic review. *American Journal of Clinical Nutrition*, 89, 2040S-2051S.
- Lowe NM, Dykes FC, Skinner AL, Patel S, Warthon-Medina M, Decsi T, Fekete K, Souverein OW, Dullemeijer C, Cavelaars AE, Serra-Majem L, Nissensohn M, Bel S, Moreno LA, Hermoso M, Vollhardt C, Berti C, Cetin I, Gurinovic M, Novakovic R, Harvey LJ, Collings R and Hall-Moran V, 2013. EURRECA-Estimating zinc requirements for deriving dietary reference values. *Critical Reviews in Food Science and Nutrition*, 53, 1110-1123.
- Mahomed K, James DK, Golding J and McCabe R, 1989. Zinc supplementation during pregnancy: a double blind randomised controlled trial. *British Medical Journal (Clinical Research Edition)*, 299, 826-830.
- Manary MJ, Hotz C, Krebs NF, Gibson RS, Westcott JE, Arnold T, Broadhead RL and Hambidge KM, 2000. Dietary phytate reduction improves zinc absorption in Malawian children recovering from tuberculosis but not in well children. *Journal of Nutrition*, 130, 2959-2964.
- Mandato F, Rubegni P, Buonocore G and Fimiani M, 2009. A case of transient zinc deficiency in a breast-fed preterm infant successfully treated with oral zinc supplementation: Review of zinc metabolism and related diseases. *Giornale Italiano di Dermatologia e Venereologia*, 144, 729-734.
- Matos C, Moutinho C, Balcao V, Almeida C, Ribeiro M, Marques A and Guerra A, 2009. Total antioxidant activity and trace elements in human milk: the first 4 months of breast-feeding. *European Food Research and Technology*, 230, 201-208.
- Mazariegos M, Hambidge KM, Krebs NF, Westcott JE, Lei S, Grunwald GK, Campos R, Barahona B, Raboy V and Solomons NW, 2006. Zinc absorption in Guatemalan schoolchildren fed normal or low-phytate maize. *American Journal of Clinical Nutrition*, 83, 59-64.
- Mensink GB, Heseker H, Richter A, Stahl A and Vohmann C (Robert Koch-Institut & Universität Paderborn), 2007. *Ernährungsstudie als KIGGS-Modul (EsKiMo)*. 143 pp.
- Milacic R, Ajlec D, Zuliani T, Zigon D and Scancar J, 2012. Determination of Zn-citrate in human milk by CIM monolithic chromatography with atomic and mass spectrometry detection. *Talanta*, 101, 203-210.
- Miletta MC, Bieri A, Kernland K, Schoni MH, Petkovic V, Fluck CE, Eble A and Mullis PE, 2013. Transient neonatal zinc deficiency caused by a heterozygous G87R mutation in the zinc transporter ZnT-2 (SLC30A2) gene in the mother highlighting the importance of Zn (2+) for normal growth and development. *International Journal of Endocrinology*, 2013, 259189.
- Miller LV, Krebs NF and Hambidge KM, 1998. Human zinc metabolism: advances in the modeling of stable isotope data. *Advances in Experimental Medicine and Biology*, 445, 253-269.
- Miller LV, Krebs NF and Hambidge KM, 2000. Development of a compartmental model of human zinc metabolism: identifiability and multiple studies analyses. *American Journal of Physiology - Regulatory, Integrative and Comparative Physiology*, 279, R1671-1684.
- Miller LV, Krebs NF and Hambidge KM, 2007. A mathematical model of zinc absorption in humans as a function of dietary zinc and phytate. *Journal of Nutrition*, 137, 135-141.
- Mills C, 1989. *Zinc in Human Biology*. Springer, London, UK, 388 pp.

- Milne DB, Canfield WK, Mahalko JR and Sandstead HH, 1983. Effect of dietary zinc on whole body surface loss of zinc: impact on estimation of zinc retention by balance method. *American Journal of Clinical Nutrition*, 38, 181-186.
- Milne DB, Canfield WK, Gallagher SK, Hunt JR and Klevay LM, 1987. Ethanol metabolism in postmenopausal women fed a diet marginal in zinc. *American Journal of Clinical Nutrition*, 46, 688-693.
- Morgan PH, Mercer LP and Flodin NW, 1975. General model for nutritional responses of higher organisms. *Proceedings of the National Academy of Sciences of the United States of America*, 72, 4327-4331.
- Moser-Veillon PB and Reynolds RD, 1990. A longitudinal study of pyridoxine and zinc supplementation of lactating women. *American Journal of Clinical Nutrition*, 52, 135-141.
- Moser-Veillon PB, 1995. Zinc needs and homeostasis during lactation. *Analyst*, 120, 895-897.
- Netherlands Food and Nutrition Council, 1992. Recommended dietary allowances 1989 in The Netherlands. The Hague, The Netherlands, 115 pp.
- NNR (Nordic Nutrition Recommendations), 2004. Integrating nutrition and physical activity. Nordic Council of Ministers, Copenhagen, Denmark, 435 pp.
- Nordic Council of Ministers, 2014. Nordic Nutrition Recommendations 2012. Integrating nutrition and physical activity. 5th edition. Nordic Council of Ministers, Copenhagen, Denmark, 627 pp.
- Ortega RM, Andres P, Martinez RM, Lopez-Sobaler AM and Quintas ME, 1997. Zinc levels in maternal milk: the influence of nutritional status with respect to zinc during the third trimester of pregnancy. *European Journal of Clinical Nutrition*, 51, 253-258.
- Perrone L, Di Palma L, Di Toro R, Gialanella G and Moro R, 1993. Trace element content of human milk during lactation. *Journal of Trace Elements and Electrolytes in Health and Disease*, 7, 245-247.
- Perrone L, Dipalma L, Ditoro R, Gialanella G and Moro R, 1994. Interaction of trace elements in a longitudinal study of human milk from full-term and preterm mothers. *Biological Trace Element Research*, 41, 321-330.
- Pinna K, 1999. Effect of a low zinc intake on zinc homeostasis and immune function in adult men [doctoral dissertation]. University of California, Berkeley, CA, USA.
- Pinna K, Woodhouse LR, Sutherland B, Shames DM and King JC, 2001. Exchangeable zinc pool masses and turnover are maintained in healthy men with low zinc intakes. *Journal of Nutrition*, 131, 2288-2294.
- Piotrowska-Dept MJ, Piotrowska-Jastrzebska J, Borawska M, Markiewicz R, Socha K and Malinowska A, 2006. Zinc concentration in breast-milk of women living in Podlasie. *Pediatrica Wspolczesna Zawartosc cynku w mleku kobiet karmiaczych piersia zamieszkalych na terenie Podlasia*, 2006, 121-125.
- Plaami S and Kumpulainen J, 1996. Trace elements, natural antioxidants and contaminants in European foods and diets. *Proceedings of a Technical Workshop on Trace Elements, Natural Antioxidants and Contaminants*. Helsinki (Finland), 25-26 Aug 1995. REU Technical Series (FAO). 1020-3737, no. 49, 169-185.
- Prynne CJ, McCarron A, Wadsworth ME and Stephen AM, 2010. Dietary fibre and phytate--a balancing act: results from three time points in a British birth cohort. *British Journal of Nutrition*, 103, 274-280.
- Rodriguez Rodriguez EM, Sanz Alaejos M and Diaz Romero C, 2000. Concentrations of iron, copper and zinc in human milk and powdered infant formula. *International Journal of Food Sciences and Nutrition*, 51, 373-380.

- Roe MA, Bell S, Oseredczuk M, Christensen T, Westenbrink S, Pakkala H, Presser K and Finglas PM, 2013. Updated food composition database for nutrient intake EFSA supporting publication 2013:EN-355, 21 pp.
- Rosado JL, Hambidge KM, Miller LV, Garcia OP, Westcott J, Gonzalez K, Conde J, Hotz C, Pfeiffer W, Ortiz-Monasterio I and Krebs NF, 2009. The quantity of zinc absorbed from wheat in adult women is enhanced by biofortification. *Journal of Nutrition*, 139, 1920-1925.
- Ruz M, 1984. Recommended zinc intake for the first six months of life. *Nutrition Research*, 4, 923-927.
- Ryu MS, Guthrie GJ, Maki AB, Aydemir TB and Cousins RJ, 2012. Proteomic analysis shows the upregulation of erythrocyte dematin in zinc-restricted human subjects. *American Journal of Clinical Nutrition*, 95, 1096-1102.
- Salmenpera L, Perheentupa J, Nanto V and Siimes MA, 1994. Low zinc intake during exclusive breast-feeding does not impair growth. *Journal of Pediatric Gastroenterology and Nutrition*, 18, 361-370.
- Sandstead HH, 1973. Zinc nutrition in the United States. *American Journal of Clinical Nutrition*, 26, 1251-1260.
- Sandstrom B, Bugel S, McGaw BA, Price J and Reid MD, 2000. A high oat-bran intake does not impair zinc absorption in humans when added to a low-fiber animal protein-based diet. *Journal of Nutrition*, 130, 594-599.
- SCF (Scientific Committee for Food), 1993. Nutrient and energy intakes for the European Community. Reports of the Scientific Committee for Food, 31st series. Food - Science and Techniques. Scientific Committee for Food, Luxembourg, European Commission, 248 pp.
- SCF (Scientific Committee on Food), 2002. Opinion of the Scientific Committee on Food on the Tolerable Upper Intake Level of zinc (expressed on 5 March 2002). SCF/CS/NUT/UPPLEV/62 Final, 18 pp.
- Schlemmer U, 1995. Phytinsäuren und Phytasen in der Human- und Tierernährung. In: 5th Symposium Vitamine und Zusatzstoffe in der Ernährung von Mensch und Tier, Friedrich-Schiller-Universität. Eds Schubert R, Flachowsky G and Bitsch R. Jena, Germany, 354-365.
- Schlemmer U, Frolich W, Prieto RM and Grases F, 2009. Phytate in foods and significance for humans: food sources, intake, processing, bioavailability, protective role and analysis. *Molecular Nutrition & Food Research*, 53 Suppl 2, S330-375.
- Sette S, Le Donne C, Piccinelli R, Arcella D, Turrini A and Leclercq C, 2011. The third Italian National Food Consumption Survey, INRAN-SCAI 2005-06 - Part 1: Nutrient intakes in Italy. *Nutrition, Metabolism and Cardiovascular Diseases*, 21, 922-932.
- Shaw JC, 1979. Trace elements in the fetus and young infant. I. Zinc. *American Journal of Diseases of Children*, 133, 1260-1268.
- Sheng XY, Hambidge KM, Zhu XX, Ni JX, Bailey KB, Gibson RS and Krebs NF, 2006. Major variables of zinc homeostasis in Chinese toddlers. *American Journal of Clinical Nutrition*, 84, 389-394.
- Sheng XY, Hambidge KM, Miller LV, Westcott JE, Lei S and Krebs NF, 2009. Measurement of zinc absorption from meals: comparison of extrinsic zinc labeling and independent measurements of dietary zinc absorption. *International Journal for Vitamin and Nutrition Research*, 79, 230-237.
- Sian L, Mingyan X, Miller LV, Tong L, Krebs NF and Hambidge KM, 1996. Zinc absorption and intestinal losses of endogenous zinc in young Chinese women with marginal zinc intakes. *American Journal of Clinical Nutrition*, 63, 348-353.
- Sievers E, Oldigs HD, Dorner K and Schaub J, 1992. Longitudinal zinc balances in breast-fed and formula-fed infants. *Acta Paediatrica*, 81, 1-6.

- Sievers E and Schaub J, 2004. Zinc concentration in human milk: Longitudinal studies in consecutive lactation periods. *Advances in Experimental Medicine and Biology*, 554, 403-405.
- Silvestre D, Martinez-Costa C, Lagarda MJ, Brines J, Farre R and Clemente G, 2001. Copper, iron, and zinc contents in human milk during the first three months of lactation: A longitudinal study. *Biological Trace Element Research*, 80, 1-11.
- Silvestre MD, Lagarda MJ, Farre R, Martinez-Costa C and Brines J, 2000a. Copper, iron and zinc determinations in human milk using FAAS with microwave digestion. *Food Chemistry*, 68, 95-99.
- Silvestre MD, Lagarda MJ, Farre R, Martinez-Costa C, Brines J, Molina A and Clemente G, 2000b. A study of factors that may influence the determination of copper, iron, and zinc in human milk during sampling and in sample individuals. *Biological Trace Element Research*, 76, 217-227.
- Stawarz R, Formicki G and Massanyi P, 2007. Daily fluctuations and distribution of xenobiotics, nutritional and biogenic elements in human milk in Southern Poland. *Journal of Environmental Science and Health - Part A Toxic/Hazardous Substances and Environmental Engineering*, 42, 1169-1175.
- Swanson CA and King JC, 1982. Zinc utilization in pregnant and nonpregnant women fed controlled diets providing the zinc RDA. *Journal of Nutrition*, 112, 697-707.
- Swanson CA and King JC, 1987. Zinc and pregnancy outcome. *American Journal of Clinical Nutrition*, 46, 763-771.
- Taylor CM, Bacon JR, Aggett PJ and Bremner I, 1991. Homeostatic regulation of zinc absorption and endogenous losses in zinc-deprived men. *American Journal of Clinical Nutrition*, 53, 755-763.
- Tipton K, Green NR, Haymes EM and Waller M, 1993. Zinc loss in sweat of athletes exercising in hot and neutral temperatures. *International Journal of Sport Nutrition*, 3, 261-271.
- Torelm I and Bruce A, 1982. Fytinsyra i livesmedel. *Var Foda*, 34, 79-96.
- Turnlund JR, King JC, Keyes WR, Gong B and Michel MC, 1984. A stable isotope study of zinc absorption in young men: effects of phytate and alpha-cellulose. *American Journal of Clinical Nutrition*, 40, 1071-1077.
- Turnlund JR, Durkin N, Costa F and Margen S, 1986. Stable isotope studies of zinc absorption and retention in young and elderly men. *Journal of Nutrition*, 116, 1239-1247.
- Turnlund JR, Keyes WR, Hudson CA, Betschart AA, Kretsch MJ and Sauberlich HE, 1991. A stable-isotope study of zinc, copper, and iron absorption and retention by young women fed vitamin B-6-deficient diets. *American Journal of Clinical Nutrition*, 54, 1059-1064.
- Ustundag B, Yilmaz E, Dogan Y, Akarsu S, Canatan H, Halifeoglu I, Cikim G and Denizmen Aygun A, 2005. Levels of cytokines (IL-1(beta), IL-2, IL-6, IL-8, TNF-(alpha)) and trace elements (Zn, Cu) in breast milk from mothers of preterm and term infants. *Mediators of Inflammation*, 2005, 331-336.
- van't Hof MA, Haschke F and Darvay S, 2000. Euro-Growth references on increments in length, weight, and head and arm circumferences during the first 3 years of life. Euro-Growth Study Group. *Journal of Pediatric Gastroenterology and Nutrition*, 31 Suppl 1, S39-47.
- van Buuren S, Schönbeck Y and van Dommelen P, 2012. Collection, collation and analysis of data in relation to reference heights and reference weights for female and male children and adolescents (0-18 years) in the EU, as well as in relation to the age of onset of puberty and the age at which different stages of puberty are reached in adolescents in the EU. Project developed on the procurement project CT/EFSA/NDA/2010/01. EFSA supporting publication 2012:EN-255, 59 pp.
- van Rossum CTM, Fransen HP, Verkaik-Kloosterman J, Burma-Rethans EJM and Ocké MC, 2011. Dutch National Food Consumption Survey 2007-2010: Diet of children and adults aged 7 to 69 years. RIVM Report number: 350050006/2011, National Institute for Public Health and the Environment, 143 pp.

- Vuori E, 1979a. Copper, iron, manganese and zinc in breast milk. University of Helsinki, Helsinki, Finland, 40 pp.
- Vuori E, 1979b. Intake of copper, iron, manganese and zinc by healthy, exclusively-breast-fed infants during the first 3 months of life. *British Journal of Nutrition*, 42, 407-411.
- Vuori E, Makinen SM, Kara R and Kuitunen P, 1980. The effects of the dietary intakes of copper, iron, manganese, and zinc on the trace element content of human milk. *American Journal of Clinical Nutrition*, 33, 227-231.
- Wada L, Turnlund JR and King JC, 1985. Zinc utilization in young men fed adequate and low zinc intakes. *Journal of Nutrition*, 115, 1345-1354.
- Wada L and King JC, 1986. Effect of low zinc intakes on basal metabolic rate, thyroid hormones and protein utilization in adult men. *Journal of Nutrition*, 116, 1045-1053.
- Walravens PA, Krebs NF and Hambidge KM, 1983. Linear growth of low income preschool children receiving a zinc supplement. *American Journal of Clinical Nutrition*, 38, 195-201.
- Walravens PA, Hambidge KM and Koepfer DM, 1989. Zinc supplementation in infants with a nutritional pattern of failure to thrive: a double-blind, controlled study. *Pediatrics*, 83, 532-538.
- Wasowicz W, Gromadzinska J, Szram K, Rydzynski K, Cieslak J and Pietrzak Z, 2001. Selenium, zinc, and copper concentrations in the blood and milk of lactating women. *Biological Trace Element Research*, 79, 221-233.
- Wastney ME, Aamodt RL, Rumble WF and Henkin RI, 1986. Kinetic analysis of zinc metabolism and its regulation in normal humans. *American Journal of Physiology*, 251, R398-408.
- Weigand E and Kirchgessner M, 1982. Factorial estimation of the zinc requirement of lactating dairy cows. *Zeitschrift für Tierphysiologie, Tierernährung und Futtermittelkunde*, 47, 1-9.
- Weigand E and Kirchgessner M, 1992. Absorption, endogenous excretion, and balance of zinc in growing rats on diets with various sugars replacing starch. *Biological Trace Element Research*, 34, 67-77.
- Wessells KR, King JC and Brown KH, 2014. Development of a plasma zinc concentration cutoff to identify individuals with severe zinc deficiency based on results from adults undergoing experimental severe dietary zinc restriction and individuals with acrodermatitis enteropathica. *Journal of Nutrition*, 144, 1204-1210.
- WHO (World Health Organization), 1973. Trace elements in human nutrition. WHO Technical Report Series 532. 64 pp.
- WHO (World Health Organization), 1996. Trace elements in human nutrition and health. 343 pp.
- WHO Multicentre Growth Reference Study Group (World Health Organization), 2006. WHO Child Growth Standards: Length/height-for-age, weight-for-age, weight-for-length, weight-for-height and body mass index-for-age: Methods and development. 312 pp.
- WHO/FAO (World Health Organization/Food and Agriculture Organization of the United Nations), 2004. Vitamin and mineral requirements in human nutrition. Report of a joint FAO/WHO expert consultation, Bangkok, Thailand, 21-30 September 1998. 341 pp.
- Widdowson EM and Spray CM, 1951. Chemical development in utero. *Archives of Disease in Childhood*, 26, 205-214.
- Widdowson EM and Dickerson JWT, 1964. Chemical composition of the body. In: *Mineral Metabolism. An Advanced Treatise, Vol. II. The elements. Part A.* Eds Comar CL and Bronner F. Academic Press, New York, USA, p 26.
- Yalcin SS, Baykan A, Yurdakok K, Yalcin S and Gucus AI, 2009. The factors that affect milk-to-serum ratio for iron during early lactation. *Journal of Pediatric Hematology/Oncology*, 31, 85-90.

Younoszai HD, 1983. Clinical zinc deficiency in total parenteral nutrition: zinc supplementation. *Journal of Parenteral and Enteral Nutrition*, 7, 72-74.

APPENDICES

Appendix A. Concentrations of zinc in breast milk from mothers of term infants in Europe

Reference	n (number of samples)	Country	Maternal dietary intake (mg/day) mean \pm SD	Stage of lactation	Zinc concentration (mg/L)			Analytical method	Comments				
					Mean \pm SD	Median	Range						
Bates and Tsuchiya (1990)	57	UK	Not reported	2 months post partum	1.34 ^(a)			AAS	Infants assumed to be term infants on the basis of the study design and setting				
				3 months post partum	2.06 ^(a)								
				4 months post partum	0.87 ^(a)								
				5 months post partum	0.73 ^(a)								
				6 months post partum	0.73 ^(a)								
Bjorklund et al. (2012)	60	Sweden	Not reported	14–21 days post partum	3.47 \pm 0.98	3.52	1.24–5.71	ICP-MS					
Chierici et al. (1999)	11	Italy	Non-supplemented women with a dietary zinc intake of about 12 mg/day as estimated by three-day dietary record	3 days post partum	8.16 \pm 2.96			Inorganic mass spectrometry					
				30 days post partum	3.99 \pm 1.01								
				90 days post partum	2.87 \pm 1.23								
				Supplemented women receiving 20 mg/day of supplemental zinc									
				3 days post partum	5.89 \pm 2.65								
				30 days post partum	3.36 \pm 1.40								
				90 days post partum	2.63 \pm 1.35								
				Domellof et al. (2004)	86	Sweden	Not reported	9 months post partum	0.46 \pm 0.26			AAS	Dietary intake of the mothers assessed with a five-day food diary
				Elmastas et al. (2005)	32 (32)	Turkey	Not reported	2 months post partum	1.20 \pm 0.01			FAAS with microwave digestion	
Kantola and Vartiainen (2001)	175 (175)	Finland	Not reported	4 weeks post partum	3.00 \pm 1.00			FAAS with microwave digestion	Two analyses (in 1987 and in 1993–1995)				
	81 (81)				1.40 \pm 0.70								

Reference	n (number of samples)	Country	Maternal dietary intake (mg/day) mean \pm SD	Stage of lactation	Zinc concentration (mg/L)			Analytical method	Comments
					Mean \pm SD	Median	Range		
Leotsinidis et al. (2005)	180 (180) 95 (95)	Greece	Not reported	3 days post partum	4.91 \pm 1.73	5.01	1.32–9.12	FAAS	
				17 days post partum	2.99 \pm 0.92	2.97	0.86–6.55		
Matos et al. (2009)	31 (155)	Portugal	Not reported	7 days post partum	4.13 \pm 1.22 ^(a)	4.04 ^(a)	2.12–6.98 ^(a)	ICP-MS	
				4 weeks post partum	2.22 \pm 0.61 ^(a)	2.10 ^(a)	1.26–3.77 ^(a)		
				8 weeks post partum	1.53 \pm 0.64 ^(a)	1.46 ^(a)	0.33–3.05 ^(a)		
				12 weeks post partum	1.11 \pm 0.56 ^(a)	1.10 ^(a)	0.22–2.27 ^(a)		
				16 weeks post partum	1.04 \pm 0.47 ^(a)	1.00 ^(a)	0.15–2.29 ^(a)		
Ortega et al. (1997)	25	Spain	Women with zinc intake < 50 % RI ^(b) from diet and supplements assessed during the third trimester of pregnancy: 8.3 \pm 1.0 Women with zinc intake > 50 % RI ^(b) from diet and supplements assessed during the third trimester of pregnancy: 12.3 \pm 1.9	13–14 days post partum	3.03 \pm 0.47			Not reported	Maternal intake assessed with five-day dietary record
				40 days post partum	1.86 \pm 0.40				
				13–14 days post partum	3.31 \pm 0.60				
				40 days post partum	2.15 \pm 0.52				
Perrone et al. (1993); Perrone et al. (1994)	(46) (15) (19) (59)	Italy	Not reported	1 week post partum		36.4 \pm 2.8 ^(c, d)		Not reported	
				2 weeks post partum		24.2 \pm 1.6 ^(c, d)			
				3 weeks post partum		28.6 \pm 6.8 ^(c, d)			
				> 3 weeks post partum		21.7 \pm 1.4 ^(c, d)			
Piotrowska-Dept et al. (2006)	27 18 8	Poland	10.7 \pm 3.3 (range 5.7–18.2)	0–30 days post partum	3.42 \pm 1.62	3.29	0.53–7.28	AAS	Dietary zinc intake of the mothers assessed by a 24-hour record
				31–90 days post partum	1.50 \pm 0.87	1.37	0.12–3.58		
				> 90 days post partum	0.86 \pm 0.57	0.64	0.28–1.51		
Rodriguez Rodriguez et al. (2000)	11 (56)	Spain	Not reported	2 weeks to 5 months post partum	2.10 \pm 1.10		0.14–3.99	AAS	

Reference	n (number of samples)	Country	Maternal dietary intake (mg/day) mean \pm SD	Stage of lactation	Zinc concentration (mg/L)			Analytical method	Comments
					Mean \pm SD	Median	Range		
Salmenpera et al. (1994)	75	Finland	Non- supplemented women	4–5 days post partum	4.75		3.27–6.9		
	77			2 months post partum	1.41		1.1–2.19		
	67			4 months post partum	0.9		0.58–1.38		
	56			6 months post partum	0.67		0.4–1.13		
	31			7.5 months post partum	0.61		0.39–0.97		
	14			9 months post partum	0.6		0.38–0.95		
	8			10 months post partum	0.61		0.42–0.87		
	6			11 months post partum	0.43		0.33–0.57		
	5			12 months post partum	0.43		0.33–0.56		
	62			Supplemented women (20 mg/day)	4–5 days post partum	4.94		3.50–6.98	
	58		2 months post partum		1.52		1.10–2.11		
	48		4 months post partum		0.95		0.65–1.39		
	38		6 months post partum		0.67		0.43–1.03		
	26		7.5 months post partum		0.63		0.43–0.92		
	16		9 months post partum		0.6		0.41–0.88		
	4		10 months post partum		0.41		0.36–0.48		
	5		11 months post partum		0.51		0.44–0.60		
	2		12 months post partum		0.46		0.26–0.79		
	22		Supplemented women (40 mg/day)		4–5 days post partum	5.18		3.33–8.04	
	24			2 months post partum	1.38		0.70–2.73		
15	4 months post partum	1.08			0.61–1.94				
13	6 months post partum	0.88			0.50–1.53				
5	7.5 months post partum	0.9			0.62–1.30				
4	9 months post partum	0.94			0.71–1.24				
Sievers et al. (1992)	10	Germany	Not reported	17 days post partum	3.6			AAS	
				35 days post partum	2.6				
				56 days post partum	1.7				
				85 days post partum	1.3				
				117 days post partum	1.2				

Reference	n (number of samples)	Country	Maternal dietary intake (mg/day) mean \pm SD	Stage of lactation	Zinc concentration (mg/L)			Analytical method	Comments
					Mean \pm SD	Median	Range		
Silvestre et al. (2000a)	(10)	Spain	Not reported	Colostrum (number of days not reported)	8.60 \pm 1.82			FAAS with microwave digestion	
				Transitional milk (number of days not reported)	3.45 \pm 0.58				
				30 days post partum	1.97 \pm 0.25				
				60 days post partum	1.24 \pm 0.33				
				90 days post partum	0.89 \pm 0.27				
Silvestre et al. (2000b)	62 (136)	Spain	Not reported	2 days post partum	7.73 \pm 0.86			FAAS with microwave digestion	
				15 days post partum	3.15 \pm 0.86				
Silvestre et al. (2001)	22 (110)	Spain	Not reported	Colostrum (number of days not reported)	7.99 \pm 3.23			FAAS	
				Transitional milk (number of days not reported)	3.31 \pm 1.06				
				30 days post partum	2.41 \pm 0.90				
				60 days post partum	1.40 \pm 0.65				
				90 days post partum	1.05 \pm 0.71				
Stawarz et al. (2007)	5 (210)	Poland	Not reported	12 weeks	17.94 \pm 7.10 ^(c)		4.42–38.61	Volumetric method	
Ustundag et al. (2005)	20	Turkey	Not reported	Colostrum (0–7 days post partum)	3.08 \pm 0.30			AAS	
				7–14 days post partum	2.72 \pm 0.20				
				21 days post partum	2.65 \pm 0.20				
				60 days post partum	2.81 \pm 0.18				

Reference	n (number of samples)	Country	Maternal dietary intake (mg/day) mean ± SD	Stage of lactation	Zinc concentration (mg/L)			Analytical method	Comments
					Mean ± SD	Median	Range		
Vuori et al. (1980)	15 (15)	Finland	13.7 ± 2.7	6–8 weeks post partum	1.89 ± 0.74			FAAS	Two seven-day food records; infants assumed to be term infants on the basis of the study design and setting
			12.8 ± 2.8	17–22 weeks post partum	0.72 ± 0.44				
Wasowicz et al. (2001)	43	Poland	Not reported	0–4 days post partum	8.2 ± 2.8			ICP-AES	
	46			5–9 days post partum	3.7 ± 1.8				
	41			10–30 days post partum	1.4 ± 0.7				
Yalcin et al. (2009)	47	Turkey	Not reported	2 weeks post partum	4.78 ± 1.83	4.5		AAS	

Studies were identified by a comprehensive literature search for publications from the year 2000 onwards, earlier publications were identified from Brown et al. (2009). The following articles based on one or two case reports are not presented in this table: Sievers and Schaub (2004), Kharfi et al. (2005), Chowanadisai et al. (2006), Coelho et al. (2006), Mandato et al. (2009); Milacic et al. (2012), Leverkus et al. (2006); Gass et al. (2010); Bieri et al. (2013); Miletta et al. (2013).

AAS, atomic absorption spectrometry; ICP-MS, inductively coupled plasma mass spectrometry; FAAS, flame atomic absorption spectrometry; RI, recommended intake; ICP-AES, inductively coupled plasma atomic emission spectroscopy.

(a): After conversion from mg/g into mg/L using a conversion factor of 1.03 kg/L of breast milk, as reported in Brown et al. (2009).

(b): Value not reported.

(c): mg/kg dry weight of breast milk.

(d): Median ± SD

Appendix B. Dietary surveys in the EFSA Comprehensive database update dataset included in the nutrient intake calculation and number of subjects in the different age classes

Country	Dietary survey	Year	Method	Days	Age (years)	Number of subjects					
						Children 1 to < 3 y	Children 3 to < 10 y	Adolescents 10 to < 18 y	Adults 18 to < 65 y	Adults 65 to < 75 y	Adults ≥ 75 y
Finland/1	DIPP	2000–2010	Dietary record	3	< 1–6	500	750				
Finland/2	NWSSP	2007–2008	48-hour dietary recall ^(a)	2 x 2 ^(a)	13–15			306			
Finland/3	FINDIET2012	2012	48-hour dietary recall ^(a)	2 ^(a)	25–74				1 295	413	
France	INCA2	2006–2007	Dietary record	7	3–79		482	973	2 276	264	84
Germany/1	EsKiMo	2006	Dietary record	3	6–11		835	393			
Germany/2	VELS	2001–2002	Dietary record	6	< 1–4	347	299				
Ireland	NANS	2008–2010	Dietary record	4	18–90				1 274	149	77
Italy	INRAN-SCAI 2005–06	2005–2006	Dietary record	3	< 1–98	36 ^(b)	193	247	2 313	290	228
Latvia	FC_PREGNANTWOMEN 2011	2011	24-hour dietary recall	2	15–45			12 ^(b)	991 ^(c)		
Netherlands	DNFCS	2007–2010	24-hour dietary recall	2	7–69		447	1 142	2 057	173	
Sweden	RISKMATEN	2010–2011	Dietary record (Web) ^(d)	4	18–80				1 430	295	72
UK	NDNS—Rolling Programme (1–3 years)	2008–2011	Dietary record	4	1–94	185	651	666	1 266	166	139

y, years; DIPP, type 1 Diabetes Prediction and Prevention survey; DNFCS, Dutch National Food Consumption Survey; EsKiMo, Ernährungstudie als KIGGS-Modul; FINDIET, the national dietary survey of Finland; INCA, étude Individuelle Nationale de Consommations Alimentaires; INRAN-SCAI, Istituto Nazionale di Ricerca per gli Alimenti e la Nutrizione - Studio sui Consumi Alimentari in Italia; FC_PREGNANTWOMEN, food consumption of pregnant women in Latvia; NANS, National Adult Nutrition Survey; NDNS, National Diet and Nutrition Survey; NWSSP, Nutrition and Wellbeing of Secondary School Pupils; VELS, Verzehrsstudie zur Ermittlung der Lebensmittelaufnahme von Säuglingen und Kleinkindern für die Abschätzung eines akuten Toxizitätsrisikos durch Rückstände von Pflanzenschutzmitteln.

(a): A 48-hour dietary recall comprises two consecutive days.

(b): 5th or 95th percentile of intake calculated over a number of subjects lower than 60 require cautious interpretation as the results may not be statistically robust (EFSA, 2011b) and therefore for these dietary surveys/age classes the 5th, 95th percentile estimates will not be presented in the intake results.

(c): One subject was excluded from the dataset because only one 24-hour dietary recall day was available, i.e. final n = 990.

(d): The Swedish dietary records were introduced through the internet.

Appendix C. Zinc intake in males in different surveys according to age classes and country

Age class (years)	Country	Survey	Intake expressed in mg/day					Intake expressed in mg/MJ				
			n ^(a)	Average	Median	P5	P95	n ^(a)	Average	Median	P5	P95
1 to < 3	Finland	DIPP_2001_2009	245	5.5	5.4	3.2	8.4	245	1.5	1.5	1.1	2.0
	Germany	VELS	174	5.0	4.8	2.9	7.3	174	1.1	1.0	0.7	1.4
	Italy	INRAN_SCAI_2005_06	20	6.2	6.3	(b)	(b)	20	1.3	1.3	(b)	(b)
	UK	NDNS–RollingProgrammeYears1–3	107	5.8	5.6	3.6	9.0	107	1.2	1.2	0.8	1.6
3 to < 10	Finland	DIPP_2001_2009	381	8.5	8.4	5.3	12.1	381	1.4	1.4	1.1	1.9
	France	INCA2	239	8.8	8.5	4.9	12.9	239	1.4	1.4	0.9	2.0
	Germany	EsKiMo	426	9.3	9.0	5.9	13.1	426	1.2	1.2	0.9	1.6
	Germany	VELS	146	6.1	5.8	3.9	8.7	146	1.1	1.0	0.8	1.4
	Italy	INRAN_SCAI_2005_06	94	9.3	8.9	5.3	13.9	94	1.3	1.3	0.9	1.8
	Netherlands	DNFCS2007_2010	231	8.0	7.7	4.6	12.6	231	0.9	0.9	0.6	1.3
	UK	NDNS–RollingProgrammeYears1–3	326	6.7	6.6	3.9	10.3	326	1.1	1.0	0.7	1.5
10 to < 18	Finland	NWSSP07_08	136	12.2	11.9	7.5	17.5	136	1.5	1.5	1.1	1.9
	France	INCA2	449	10.7	10.3	6.0	16.9	449	1.4	1.3	0.9	2.0
	Germany	EsKiMo	197	10.0	9.6	6.3	14.9	197	1.2	1.2	0.9	1.6
	Italy	INRAN_SCAI_2005_06	108	12.2	11.4	7.2	18.0	108	1.3	1.2	0.9	1.7
	Netherlands	DNFCS2007_2010	566	10.2	9.8	5.6	16.5	566	1.0	0.9	0.6	1.4
	UK	NDNS–RollingProgrammeYears1–3	340	8.7	8.4	4.9	13.5	340	1.1	1.0	0.7	1.6
18 to < 65	Finland	FINDIET2012	585	12.7	12.2	6.6	20.7	585	1.4	1.3	0.9	2.0
	France	INCA2	936	11.6	11.4	6.3	18.2	936	1.3	1.3	0.9	2.0
	Ireland	NANS_2012	634	12.2	11.9	7.0	18.8	634	1.2	1.2	0.8	1.8
	Italy	INRAN_SCAI_2005_06	1 068	11.3	11.0	6.6	17.0	1 068	1.3	1.2	0.9	1.7
	Netherlands	DNFCS2007_2010	1 023	12.0	11.6	6.6	18.9	1 023	1.1	1.0	0.7	1.5
	Sweden	Riksmaten 2010	623	13.7	13.2	7.4	21.7	623	1.4	1.4	1.0	2.0
	UK	NDNS–RollingProgrammeYears1–3	560	9.9	9.5	5.3	15.9	560	1.1	1.1	0.7	1.7
	65 to < 75	Finland	FINDIET2012	210	10.8	10.4	5.9	17.0	210	1.3	1.3	0.9
France		INCA2	111	11.0	10.4	6.4	16.8	111	1.3	1.2	0.9	1.8
Ireland		NANS_2012	72	11.3	11.2	5.5	17.0	72	1.3	1.2	0.9	2.1
Italy		INRAN_SCAI_2005_06	133	11.2	10.8	6.8	16.3	133	1.3	1.3	0.9	1.7
Netherlands		DNFCS2007_2010	91	11.1	10.6	5.9	17.0	91	1.2	1.2	0.8	1.7
Sweden		Riksmaten 2010	127	12.0	11.0	7.4	19.3	127	1.4	1.3	1.0	2.0
UK		NDNS–RollingProgrammeYears1–3	75	10.0	9.8	3.7	16.5	75	1.2	1.2	0.7	1.8

Age class (years)	Country	Survey	Intake expressed in mg/day					Intake expressed in mg/MJ				
			n ^(a)	Average	Median	P5	P95	n ^(a)	Average	Median	P5	P95
≥ 75	France	INCA2	40	9.9	9.4	(b)	(b)	40	1.3	1.3	(b)	(b)
	Ireland	NANS_2012	34	10.0	9.5	(b)	(b)	34	1.3	1.2	(b)	(b)
	Italy	INRAN_SCAI_2005_06	69	10.6	10.3	6.9	15.0	69	1.2	1.2	1.0	1.7
	Sweden	Riksmaten 2010	42	11.0	10.9	(b)	(b)	42	1.3	1.2	(b)	(b)
	UK	NDNS–RollingProgrammeYears1–3	56	8.4	8.0	(b)	(b)	56	1.2	1.1	(b)	(b)

P5, 5th percentile; P95, 95th percentile; DIPP, type 1 Diabetes Prediction and Prevention survey; DNFCS, Dutch National Food Consumption Survey; EsKiMo, Ernährungstudie als KIGGS-Modul; FINDIET, the national dietary survey of Finland; INCA, étude Individuelle Nationale de Consommations Alimentaires; INRAN-SCAI, Istituto Nazionale di Ricerca per gli Alimenti e la Nutrizione - Studio sui Consumi Alimentari in Italia; FC_PREGNANTWOMEN, food consumption of pregnant women in Latvia; NANS, National Adult Nutrition Survey; NDNS, National Diet and Nutrition Survey; NWSSP, Nutrition and Wellbeing of Secondary School Pupils; VELs, Verzehrsstudie zur Ermittlung der Lebensmittelaufnahme von Säuglingen und Kleinkindern für die Abschätzung eines akuten Toxizitätsrisikos durch Rückstände von Pflanzenschutzmitteln.

(a): Number of individuals in the population group.

(b): 5th or 95th percentile of intake calculated from less than 60 subjects requires cautious interpretation, as the results may not be statistically robust (EFSA, 2011b) and, therefore, for these dietary surveys/age classes, the 5th and 95th percentile estimates will not be presented in the intake results.

Appendix D. Zinc intake in females in different surveys according to age classes and country

Age class (years)	Country	Survey	Intake expressed in mg/day					Intake expressed in mg/MJ				
			n ^(a)	Average	Median	P5	P95	n ^(a)	Average	Median	P5	P95
1 to < 3	Finland	DIPP_2001_2009	255	5.3	5.0	2.9	8.3	255	1.5	1.5	1.1	2.0
	Germany	VELS	174	4.6	4.6	2.8	6.7	174	1.1	1.1	0.8	1.4
	Italy	INRAN_SCAI_2005_06	16	5.8	6.1	(b)	(b)	16	1.3	1.1	(b)	(b)
	UK	NDNS–RollingProgrammeYears1–3	78	5.3	5.3	3.1	7.8	78	1.2	1.2	0.9	1.5
3 to < 10	Finland	DIPP_2001_2009	369	7.7	7.6	5.3	10.9	369	1.5	1.4	1.1	1.9
	France	INCA2	243	7.9	7.8	5.0	12.4	243	1.4	1.4	0.9	2.1
	Germany	EsKiMo	409	8.4	8.1	5.5	12.4	409	1.2	1.2	0.9	1.7
	Germany	VELS	147	5.5	5.3	3.5	8.4	147	1.1	1.0	0.7	1.5
	Italy	INRAN_SCAI_2005_06	99	8.9	8.5	5.0	13.1	99	1.2	1.2	0.9	1.7
	Netherlands	DNFCS2007_2010	216	7.5	7.3	4.3	11.7	216	0.9	0.9	0.5	1.4
	UK	NDNS–RollingProgrammeYears1–3	325	6.4	6.3	3.7	9.6	325	1.1	1.0	0.7	1.5
10 to < 18	Finland	NWSSP07_08	170	9.7	9.5	5.7	15.0	170	1.5	1.5	1.1	1.9
	France	INCA2	524	8.4	8.2	4.7	12.9	524	1.3	1.3	0.9	2.0
	Germany	EsKiMo	196	9.3	9.0	6.0	13.5	196	1.2	1.2	0.9	1.7
	Italy	INRAN_SCAI_2005_06	139	9.8	9.6	5.8	14.4	139	1.2	1.2	0.9	1.6
	Latvia	FC_PREGNANTWOMEN_2011 ^(c)	12	14.5	13.2	(b)	(b)	12	1.5	1.5	(b)	(b)
	Netherlands	DNFCS2007_2010	576	8.4	8.2	4.9	12.5	576	1.0	0.9	0.6	1.4
	UK	NDNS–RollingProgrammeYears1–3	326	6.8	6.7	3.4	10.6	326	1.0	1.0	0.6	1.5
18 to < 65	Finland	FINDIET2012	710	9.8	9.5	5.2	15.5	710	1.4	1.3	0.9	2.0
	France	INCA2	1340	8.9	8.6	4.7	14.0	1340	1.4	1.3	0.9	2.2
	Ireland	NANS_2012	640	9.0	8.8	5.0	13.8	640	1.2	1.2	0.8	1.8
	Italy	INRAN_SCAI_2005_06	1245	9.4	9.2	5.4	13.8	1245	1.3	1.2	0.9	1.8
	Latvia	FC_PREGNANTWOMEN_2011 ^(c)	990	14.0	13.3	7.9	22.8	990	1.7	1.6	1.0	2.6
	Netherlands	DNFCS2007_2010	1034	9.5	9.0	5.3	15.0	1034	1.2	1.1	0.7	1.8
	Sweden	Riksmaten 2010	807	10.5	10.1	5.8	16.6	807	1.4	1.3	1.0	2.0
	UK	NDNS–RollingProgrammeYears1–3	706	8.0	7.8	4.2	12.3	706	1.2	1.2	0.7	1.8
65 to < 75	Finland	FINDIET2012	203	8.5	8.3	4.5	13.2	203	1.4	1.4	0.9	1.9
	France	INCA2	153	8.6	8.0	4.5	13.7	153	1.4	1.3	0.9	2.2
	Ireland	NANS_2012	77	9.9	9.7	5.0	14.7	77	1.5	1.4	1.0	2.1
	Italy	INRAN_SCAI_2005_06	157	9.1	8.9	4.7	14.5	157	1.3	1.3	0.9	1.9
	Netherlands	DNFCS2007_2010	82	8.7	8.6	4.2	13.5	82	1.2	1.2	0.7	1.7
	Sweden	Riksmaten 2010	168	9.6	9.3	5.1	14.8	168	1.4	1.3	1.0	1.9
	UK	NDNS–RollingProgrammeYears1–3	91	8.1	8.0	5.2	11.8	91	1.4	1.3	0.9	1.9

Age class (years)	Country	Survey	Intake expressed in mg/day					Intake expressed in mg/MJ				
			n ^(a)	Average	Median	P5	P95	n ^(a)	Average	Median	P5	P95
≥ 75	France	INCA2	44	8.0	7.3	(b)	(b)	44	1.3	1.3	(b)	(b)
	Ireland	NANS_2012	43	9.1	9.2	(b)	(b)	43	1.5	1.4	(b)	(b)
	Italy	INRAN_SCAI_2005_06	159	8.6	8.4	5.1	12.4	159	1.3	1.3	0.9	1.8
	Sweden	Riksmaten 2010	30	9.3	9.4	(b)	(b)	30	1.3	1.3	(b)	(b)
	UK	NDNS–RollingProgrammeYears1–3	83	8.2	7.8	4.9	12.3	83	1.4	1.3	0.9	1.9

P5, 5th percentile; P95, 95th percentile; DIPP, type 1 Diabetes Prediction and Prevention survey; DNFCS, Dutch National Food Consumption Survey; EsKiMo, Ernährungstudie als KIGGS-Modul; FINDIET, the national dietary survey of Finland; INCA, étude Individuelle Nationale de Consommations Alimentaires; INRAN-SCAI, Istituto Nazionale di Ricerca per gli Alimenti e la Nutrizione - Studio sui Consumi Alimentari in Italia; FC_PREGNANTWOMEN, food consumption of pregnant women in Latvia; NANS, National Adult Nutrition Survey; NDNS, National Diet and Nutrition Survey; NWSSP, Nutrition and Wellbeing of Secondary School Pupils; VELS, Verzehrsstudie zur Ermittlung der Lebensmittelaufnahme von Säuglingen und Kleinkindern für die Abschätzung eines akuten Toxizitätsrisikos durch Rückstände von Pflanzenschutzmitteln.

(a): Number of individuals in the population group.

(b): 5th or 95th percentile of intake calculated from less than 60 subjects requires cautious interpretation, as the results may not be statistically robust (EFSA, 2011b) and, therefore, for these dietary surveys/age classes, the 5th and 95th percentile estimates will not be presented in the intake results.

(c): Pregnant women only.

Appendix E. Minimum and maximum percentage contribution of different FoodEx2 level 1 food groups to zinc intake in males

Food groups	Age (years)					
	1 to < 3	3 to < 10	10 to < 18	18 to < 65	65 to < 75	≥ 75
Additives, flavours, baking and processing aids	< 0.1–0.2	0–0.7	< 0.1–1.1	< 0.1–0.2	0–0.1	0
Alcoholic beverages	< 0.1	< 0.1	< 0.1–0.1	0.7–1.6	0.7–2.2	0.3–2.2
Animal and vegetable fats and oils	0.1–0.3	0.1–0.3	0.1–0.3	0.1–0.3	0.1–0.4	0.1–0.4
Coffee, cocoa, tea and infusions	< 0.1–2.2	1–4.1	1.5–2.9	0.9–4.6	0.9–7.1	0.7–7.9
Composite dishes	0.6–12.8	0.1–15.8	0.3–23	0.4–18.6	0.5–12.5	0.2–12.8
Eggs and egg products	0.5–2	0.1–3.4	0.1–3.3	< 0.1–2.6	< 0.1–2.6	0.1–2.4
Fish, seafood, amphibians, reptiles and invertebrates	0.2–7.2	0.2–6.5	0.3–6	0.7–6.4	1.1–7.4	2.1–5.4
Food products for young population	3–15.6	0.2–0.8	< 0.1–0.1	< 0.1	–	–
Fruit and fruit products	1.6–3.2	1.3–6.6	0.8–2.2	0.9–2	1.2–3	1.5–5.3
Fruit and vegetable juices and nectars	0.5–1.9	0.3–2.3	0.3–2.1	0.2–1.4	0.1–1.4	0.1–0.7
Grains and grain-based products	18.8–31.7	18.1–35.4	20–36	18.5–28.9	18.7–31.5	18.9–34.5
Human milk	< 0.1–1.7	–	–	–	–	–
Legumes, nuts, oilseeds and spices	1.1–3	1.2–4	1.1–3.2	1.6–4	1.8–4.4	1.3–2.9
Meat and meat products	9.8–24.1	13.2–36.7	19.7–44.5	22.4–45.8	20.5–40.8	20.1–38.8
Milk and dairy products	27.2–34.2	19.1–34.4	13.4–31.1	12.1–23.9	11.6–24.8	13–20.4
Products for non-standard diets, food imitates and food supplements or fortifying agents	0–0.2	0–1.1	< 0.1–0.6	< 0.1–1	< 0.1–0.6	0–0.2
Seasoning, sauces and condiments	0.1–0.7	0.1–0.6	0.1–0.6	0.2–0.9	0.2–0.8	0.2–0.8
Starchy roots or tubers and products thereof, sugar plants	1–2.4	0.7–4.9	0.7–5.9	1.2–5.1	1.2–5.1	1.2–5.6
Sugar, confectionery and water-based sweet desserts	0.2–2.7	0.4–4.3	0.4–4.2	0.3–1	0.2–0.6	0.1–0.7
Vegetables and vegetable products	2–6.5	2.1–7	2–6.8	1.5–8.8	1.7–9.8	2.1–9.8
Water and water-based beverages	0.5–1.7	0.5–1.6	0.6–2.1	0.6–1.6	0.4–1.5	0.3–1.5

“–” means that there was no consumption event of the food group for the age and sex group considered, whereas “0” means that there were some consumption events, but that the food group does not contribute to the intake of the nutrient considered, for the age and sex group considered.

Appendix F. Minimum and maximum percentage contribution of different FoodEx2 level 1 food groups to zinc intake in females

Food groups	Age (years)					
	1 to < 3	3 to < 10	10 to < 18	18 to < 65	65 to < 75	≥ 75
Additives, flavours, baking and processing aids	0–0.2	0–0.7	< 0.1–1.1	< 0.1–0.2	< 0.1–0.1	0
Alcoholic beverages	< 0.1	< 0.1	< 0.1–0.1	< 0.1–0.8	0.1–1	0.2–0.9
Animal and vegetable fats and oils	0.1–0.3	0.1–0.3	0.1–0.3	0.1–0.3	0.1–0.4	0.1–0.4
Coffee, cocoa, tea and infusions	< 0.1–2.6	0.8–4.4	1.6–4.5	1.2–7.1	1–9.3	1.4–9.5
Composite dishes	0.2–11.7	0.1–15.9	0.5–23.7	0.5–15	0.3–13.2	0.5–12.2
Eggs and egg products	0.4–2.2	0.1–3.3	0.1–3.3	0.1–2.4	0.1–2.3	0.1–2.7
Fish, seafood, amphibians, reptiles and invertebrates	0.1–8.6	0.2–5.2	0.3–7.3	0.6–7	0.9–6.9	1–4.7
Food products for young population	3.1–13	< 0.1–0.4	< 0.1–0.1	< 0.1	–	< 0.1
Fruit and fruit products	1.1–2.7	1.2–8.1	1.2–6	1.2–4.7	2.6–8.7	2–4.9
Fruit and vegetable juices and nectars	0.3–1.7	0.4–2.2	0.4–2	0.2–1.3	0.2–1.3	0.2–1.1
Grains and grain-based products	19.6–31.1	16.9–36.8	20.2–35.8	17.4–39.7	16.3–31.7	15.5–33.4
Human milk	< 0.1	–	–	–	–	–
Legumes, nuts, oilseeds and spices	1–3.1	1.2–3.8	1.1–3.2	1.6–4.5	1.5–3.8	1–3.1
Meat and meat products	10.9–22.3	13.5–35.9	17.4–41.3	19.4–40	18.2–36.4	16–37.4
Milk and dairy products	27.2–30.9	19.4–36.3	13.2–30.3	13.2–26.5	13.8–26.5	14.4–24.1
Products for non-standard diets, food imitates and food supplements or fortifying agents	0–0.7	0–1.4	< 0.1–1	0.1–3.4	0.1–1.7	0–1.7
Seasoning, sauces and condiments	0.1–0.5	0.1–0.6	0.1–0.9	0.2–1	0.2–0.7	0.2–0.9
Starchy roots or tubers and products thereof, sugar plants	1.1–3.5	0.6–5.2	0.6–5.6	1–4.8	1.1–4.5	1.3–4.1
Sugar, confectionery and water-based sweet desserts	0.2–2.5	0.6–4.2	0.5–4.3	0.3–2.4	0.2–0.8	0.2–0.9
Vegetables and vegetable products	2–5.4	2.3–7.2	1.9–6.8	2.2–10.1	2.4–11.2	2.8–10.1
Water and water-based beverages	0.5–1.5	0.5–1.7	0.2–1.9	0.2–2.1	0.5–1.9	0.4–2.5

“–” means that there was no consumption event of the food group for the age and sex group considered, whereas “0” means that there were some consumption events, but that the food group does not contribute to the intake of the nutrient considered, for the age and sex group considered.

Appendix G. Phytate/Phytic acid intake in various European countries

Study	Country	Sex (n)	Age (years)	Phytic acid/phytate intake (mg/day)		Phytate–zinc molar ratio median (IQR)	Comments/methods of assessment	
				Mean ± SD or (range)	Median (IQR)			
Adults								
Amirabdollahian and Ash (2010)	UK	Male (108)	19–24	817	762 (565–940)	8.21 (6.82–10.30)	Phytate intake was assessed based on food consumption data obtained in the National Diet and Nutrition Survey and the content of phytate in food according to published and unpublished data (the phytate content of food in the UK is unavailable)	
		Male (219)	25–34	1 010	904 (659–1 132)	9.11 (7.31–11.47)		
		Male (253)	35–49	993	903 (670–1 262)	8.80 (6.58–11.65)		
		Male (253)	50–64	1 094	948 (679–1 314)	9.27 (7.24–12.23)		
		Male (371)	65–74	891	733 (509–1 112)	8.70 (6.26–11.50)		
		Male (200)	75–84	938	692 (453–1 145)	8.78 (6.32–12.58)		
		Male (62)	> 85	1 059	779 (496–1 419)	8.97 (7.13–17.94)		
		Female (104)	19–24	650	645 (438–790)	9.28 (7.00–12.11)		The authors acknowledged that those data may be inaccurate owing to the use of non-peer-reviewed food composition data, the unavailability of data on the phytate content for many foods and the accuracy of the method to measure phytate
		Female (210)	25–34	756	714 (486–910)	10.50 (8.20–13.23)		
		Female (318)	35–49	868	792 (568–1 071)	10.27 (8.00–14.03)		
		Female (259)	50–64	928	807 (599–1 138)	10.51 (8.26–13.82)		
		Female (434)	65–74	693	630 (426–849)	8.93 (6.43–11.26)		
		Female (638)	75–84	674	549 (392–777)	8.50 (6.25–10.91)		
		Female (251)	> 85	712	538 (416–772)	8.40 (6.90–11.62)		
Prynne et al. (2010)	UK	Male (562)	36	662 (626–698) ^(a)	5.7 (5.5–6.0) ^(a)	Dietary survey following the same individuals over several years (follow-up dietary survey)		
			43	666 (634–698) ^(a)	5.9 (5.6–6.1) ^(a)			
			53	715 (684–747) ^(a)	6.8 (6.5–7.0) ^(a)			
		Female (691)	36	566 (536–597) ^(a)	6.3 (6.0–6.6) ^(a)		Phytate intake was assessed based on food consumption data obtained by five-day dietary records and the phytate content of foods. The original (British) nutrient composition database was updated with phytate data for US foods for principal sources of phytate	
			43	562 (537–587) ^(a)	6.3 (6.1–6.5) ^(a)			
			53	647 (622–671) ^(a)	7.5 (7.3–7.8) ^(a)			
Heath et al. (2005)	UK	Male (49)	> 40	MBIAT: 1 436 ± 755	1 328 (918–1 876)	11.64 (8.24–15.07)		Phytate intake was assessed by MBIAT/WDR and food composition data based on published articles
				WDR: 1 366 ± 559	1 374 (855–1 707)	11.03 (8.62–14.64)		

Study	Country	Sex (n)	Age (years)	Phytic acid/phytate intake (mg/day)		Phytate–zinc molar ratio median (IQR)	Comments/methods of assessment
				Mean ± SD or (range)	Median (IQR)		
Brune et al. (1989)	Sweden	Male + female (6)	24–70	369 (230–532) ^(b)			Individuals following a “typical unrestricted Swedish diet”
		Male/female (4/9)	35–76	1 146 (500–2 927) ^(b)			Individuals following a vegetarian (omitting meat, fish and eggs) or vegan (omitting meat, fish, eggs and milk) diet Phytate intake was assessed based on food consumption data obtained by four-day dietary record and the phytate content of foods was determined with the method described in Harland and Oberleas (1986)
Plaami and Kumpulainen (1996)	Finland	nr	nr	370			Phytic acid intake was assessed from intake of cereal products only (consumption data plus the content of phytate in cereals)
Carnovale et al. (1987)	Italy	nr	nr	ISTAT diet: 219 High-plant food diets: 796 (112–1 367) ^(c)		1.54 5.92 (0.90–11.83) ^(c)	Phytic acid content of 12 diets collected over seven days in a rural area of southern Italy; diets were characterised by a high content of plant foods. One diet representative of national meal pattern trends (ISTAT) was also included. Phytic acid determined in whole diets according to a modification of the colorimetric method of Harland and Oberleas (1977)
Torelm and Bruce (1982)	Sweden	nr	nr	181			Calculated phytic acid intake assessed on the basis of selected foods and their content of phytic acid

Study	Country	Sex (n)	Age (years)	Phytic acid/phytate intake (mg/day)		Phytate–zinc molar ratio median (IQR)	Comments/methods of assessment
				Mean \pm SD or (range)	Median (IQR)		
Children							
Amirabdollahian and Ash (2010)	UK	Male (298)	1.5–2.5	601	465 (353–733)	11.50 (8.08–17.37)	Phytate intake was assessed based on food consumption data obtained in the National Diet and Nutrition Survey and the content of phytate in food according to published and unpublished data (the phytate content of food in the UK is unavailable)
		Male (300)	2.5–3.5	636	515 (408–718)	12.41 (9.47–16.92)	
		Male (250)	3.5–4.5	605	526 (406–725)	11.84 (9.03–15.83)	
		Male (184)	4–6	640	576 (435–770)	10.90 (8.99–13.08)	
		Male (256)	7–10	733	627 (519–831)	10.61 (9.08–13.50)	
		Male (237)	11–14	792	714 (540–929)	10.39 (8.15–13.10)	
		Male (179)	15–18	855	780 (616–1 010)	9.34 (7.26–11.79)	
		Female (278)	1.5–2.5	615	463 (332–695)	11.90 (8.10–17.18)	
		Female (306)	2.5–3.5	577	483 (337–688)	11.90 (8.94–15.78)	
		Female (243)	3.5–4.5	566	497 (379–680)	11.58 (9.10–16.27)	
		Female (172)	4–6	564	494 (369–657)	10.54 (8.39–13.50)	
		Female (225)	7–10	644	566 (461–740)	10.02 (8.42–12.90)	
		Female (238)	11–14	657	594 (480–789)	10.60 (8.25–12.76)	
		Female (210)	15–18	674	574 (459–829)	10.19 (7.90–13.91)	

IQR, interquartile range; MBIAT, meal-based intake assessment tool; WDR, weighed diet record; nr, not reported; ISTAT, National Institute for Statistics.

(a): Mean and 95 % confidence interval.

(b): As reported in Schlemmer et al. (2009), values in the paper by Brune et al. (1989) are for “phytate-phosphorus”.

(c): Mean (range); mean calculated from individual values given in the paper.

Appendix H. Evaluating data when endogenous faecal zinc was estimated using the zinc absorption–intestinal balance method

In the data used to estimate physiological requirements, the most critical measurements are those of EFZ and TAZ. The techniques used to determine EFZ fall into two categories: those that measure EFZ directly with the use of isotope tracers administered intravenously and sampled in the faeces (Kirchgessner and Weigand isotope dilution, compartmental modelling), and techniques that rely on tracer measurements of zinc absorption along with measurements of elemental zinc intestinal balance to determine EFZ. The latter techniques have shortcomings and are less reliable than the direct measurement methods.

During compilation and inspection of the individual data, it was observed that several of the studies using the absorption–intestinal balance technique had one or more negative EFZ values. As this is physiologically impossible, these anomalies were attributed to limitations of the intestinal balance technique and the data were removed prior to further analysis. The presence of the negative values prompted concern that the accuracy of the remaining data from these studies were also compromised. To address this concern, the EFZ data were evaluated in comparison to those acquired with the more reliable, and most likely more accurate, direct measurement methods.

The EFZ and TAZ data from studies using the zinc absorption–intestinal balance method but containing no negative EFZ values (Turnlund et al., 1984; Taylor et al., 1991; Knudsen et al., 1996) are referred to as the “balance A” data, and the data from studies having negative EFZ values (Wada et al., 1985; Hunt J et al., 1992; Hunt et al., 1995; Hunt et al., 1998) are called the “balance B” data in the following discussion. They are compared to the data from the “direct” EFZ measurement method (Jackson et al., 1984; Sian et al., 1996; Lowe et al., 1997; Miller et al., 2000; King et al., 2001; Pinna et al., 2001; Sheng et al., 2009).

The balance A EFZ and TAZ mean values were not different from the direct means (two-sided t-test p-values of 0.60 and 0.95, respectively). While the balance B TAZ means were not different from the direct means ($p = 0.13$), the EFZ means were ($p = 0.019$). Furthermore, the distribution of the combined direct and balance A data (Figure 3) was found to be different from the distribution of the balance B data, as assessed with the non-parametric Anderson–Darling test ($p = 0.009$).




Figure 3: Frequency distributions of endogenous faecal zinc (EFZ) data. The red bars show the negative EFZ values, which were removed prior to the analyses described here

More importantly, the relationships between EFZ and TAZ and EFZ and body weight were different for the balance B data. Figure 4 shows that the direct and balance A data exhibited the expected positive relationship between EFZ and TAZ and had similar slopes and intercepts. In contrast, there was not a corresponding relationship in the balance B data.




Figure 4: Data and regression lines showing the relationships between endogenous faecal zinc (EFZ) and total absorbed zinc (TAZ)

Figure 5 confirms the positive relationship between EFZ and body weight in the direct data. The balance A data suggested a positive relationship, although it was not significant. Again, the balance B data showed no evidence of a relationship.




Figure 5: Data and regression lines showing the relationships between endogenous faecal zinc (EFZ) and body weight

Finally, as would be expected from the preceding information, the fitting to the balance B data of the model used to estimate the physiological zinc requirement as a function of body weight (Section 5.1.1) produces significantly different results. An analysis comparing the model's fit with both datasets demonstrated that the weight and the (TAZ – total endogenous zinc losses) slope parameters were significantly different, with p-values of 0.044 and 0.011, respectively.

Based on the findings that the EFZ data and the balance B studies differed in important ways from the direct measurement data, the Panel decided to not include the balance B studies in the estimation of physiological zinc requirements.

Data from Sandstrom et al. (2000)

The EFZ data from the study of Sandstrom et al. (2000) were generally found to be implausibly high, with most values exceeding the range of values observed in the accepted studies (Figure 6). As with the studies described above, this is most likely attributable to the use of the zinc absorption–intestinal balance method.




Figure 6: Frequency distribution of endogenous faecal zinc (EFZ) data from the study of Sandstrom et al. (2000) (red) compared with the data from the included studies (black)

Appendix I. Data extracted from the selected studies for estimating physiological zinc requirement of adults

Study	Sex (n)	Age (years)	Body weight mean, (range), SD (kg)	Body height mean, (range), SD (m)	Body mass index mean, (range), SD (kg/m ²)	Body surface area ^(a) mean, (range), SD (m ²)	EFZ method	EFZ mean, (range), SD (mg/day)	Urine zinc mean, (range), SD (mg/day)	Total dietary zinc mean, (range), SD (mg/day)	Fractional absorption of zinc mean, (range), SD	Total absorbed zinc mean, (range), SD (mg/day)	Total dietary phytate mean, (range) (mg/day)
Taylor et al. (1991)	Male (2 × 4)	29–40	66 (60–70) 3.9	1.78 (1.7–1.9) 0.08	21 (19–22) 1.1	1.81 (1.7–1.9) 0.09	Balance ^(b)	1.6 (0.6–3.4) 1.0	0.56 (0.2–1.4) 0.40	3.2 (0.8–5.6) 2.6	0.69 (0.3–1) 0.34	1.5 (0.8–2.8) 0.74	NA
Turnlund et al. (1984)	Male (4)	25–32	68 (60–81) 9.0	1.74 (1.7–1.8) 0.06	22 (21–24) 1.6	1.82 (1.7–2.0) 0.15	Balance	2.8 (1.8–4.3) 1.1	0.53 (0.3–1.0) 0.29	15	0.34 (0.2–0.5) 0.12	5.1 (3.3–7.7) 1.9	0
Knudsen et al. (1996)	Female (3)/ male (5)	23–27	72 (60–87) 9.1	1.81 (1.7–1.9) 0.09	22 (19–26) 2.1	1.90 (1.7–2.1) 0.15	Balance	3.0 (0.5–4.7) 1.4	0.43 ^(c) (0.3–0.5) 0.10	10.2 (9.4–11) 0.88	0.29 (0.02–0.5) 0.12	3.0 (0.2–5.1) 1.4	660 (NA)
Jackson et al. (1984)	Male (1)	29	80	NA	NA	NA	K&W ^(d)	3.0	0.63	7.1	0.48	3.4	NA
Sian et al. (1996)	Female (20)	17–27	53 (42–65) 6.2	1.58 (1.5–1.7) 0.06	21 (18–24) 1.6	1.53 (1.3–1.8) 0.11	K&W	1.8 (0.9–3.3) 0.70	0.3 ^(c)	6.6 (4.0–8.9) 1.6	0.32 (0.2–0.5) 0.10	2.2 (0.8–3.5) 0.92	673 (NA)
Lowe et al. (1997)	Female (6)	21–52	57 (40–64) 8.7	1.63 (1.5–1.8) 0.08	21 (17–24) 2.4	1.61 (1.3–1.8) 0.16	Comp model ^(e)	1.9 (1.2–2.6) 0.50	0.21 (0.03–0.4) 0.12	7.1 (5.7–8.8) 1.1	0.31 (0.1–0.6) 0.15	2.1 (1.3–3.2) 0.73	585 (NA)
Sheng et al. (2009)	Female (21)	21–49	64 (51–97) 13	1.63 (1.4–1.8) 0.09	24 (18–35) 4.2	1.71 (1.4–2.2) 0.19	K&W	2.7 (1.4–5.1) 0.80	0.39 (0.08–0.7) 0.18	11.7 (5.6–29) 7.2	0.30 (0.1–0.5) 0.10	3.0 (1.0–4.7) 1.1	835 (250–2080)
Miller et al. (2000)	Female (4)/ male (1)	24–48	67 (47–84) 14	1.70 (1.6–1.8) 0.10	23 (19–27) 3.3	1.79 (1.4–2.0) 0.24	Comp model	2.8 (1.5–4.5) 1.2	0.31 (0.06–0.5) 0.17	11.5 (8–20) 5.3	0.29 (0.2–0.4) 0.05	3.1 (2.2–4.4) 0.86	NA
King et al. (2001)	Male (5)	21–35	74 (67–93) 11	1.77 (1.7–1.8) 0.04	23 (21–28) 3.1	1.91 (1.8–2.2) 0.15	Comp model	2.7 (2.4–3.0) 0.20	0.46 (0.3–0.8) 0.22	12.2	0.26 (0.2–0.3) 0.02	3.2 (2.9–3.4) 0.22	NA

Study	Sex (n)	Age (years)	Body weight mean, (range), SD (kg)	Body height mean, (range), SD (m)	Body mass index mean, (range), SD (kg/m ²)	Body surface area ^(a) mean, (range), SD (m ²)	EFZ method	EFZ mean, (range), SD (mg/day)	Urine zinc mean, (range), SD (mg/day)	Total dietary zinc mean, (range), SD (mg/day)	Fractional absorption of zinc mean, (range), SD	Total absorbed zinc mean, (range), SD (mg/day)	Total dietary phytate mean, (range) (mg/day)
Pinna et al. (2001)	Male (7)	27–47	78 (71–91) 8	1.78 (1.7–1.9) 0.08	25 (21–32) 3.7	1.98 (1.8–2.1) 0.10	Comp model	2.8 (2.1–4.2) 0.81	0.42 (0.07–0.7) 0.20	13.7	0.20 (0.1–0.3) 0.06	2.7 (1.4–3.6) 0.82	NA
Mean (range) of males	Male (31)	30.9 (21–47)	72.7 (60–93)	1.79 (1.7–1.9)	23 (19–32)	1.90 (1.7–2.2)		2.4 (0.6–4.7)	0.54 (0.07–1.4)	10.4 (0.8–20)	0.38 ^(f) (0.02–1)	2.8 (0.2–7.7)	NA
Mean (range) of females	Female (54)	27.5 (17–52)	59.1 (40–97)	1.62 (1.4–1.8)	22 (17–35)	1.64 (1.3–2.2)		2.3 (0.9–4.5)	0.32 (0.03–0.71)	9.0 (4.0–29)	0.31 (0.1–0.6)	2.6 (0.8–4.7)	NA

Where no range or standard deviation is shown, all data had the same value.

EFZ, endogenous faecal zinc; NA, not available.

(a): Calculated with Gehan–George equation (Gehan and George, 1970).

(b): Balance: combination of intestinal balance and “true” absorption measured by zinc stable isotopic labelling of diet.

(c): Some or all of the data are estimated (see text).

(d): K&W, measurements using the isotope dilution method of Kirchgessner and Weigand (Kirchgessner et al., 1980; Weigand and Kirchgessner, 1982, 1992).

(e): Comp model, compartmental modelling.

(f): For the calculation of an overall mean fractional absorption of zinc (FAZ) for men and women, the FAZ of the zinc-depleted subjects in the study of Taylor et al. (1991) were omitted. The overall mean FAZ is 0.30.

Appendix J. Data regression analysis diagnostic results

The physiological requirement model




Figure 7: Residuals plotted against predictor variables and predicted values of the response variable




Figure 8: Normality plot of residuals

The plots of Figures 7 and 8 show no problems with the regression assumptions, although there are two points with large standardised and studentised residuals. The externally studentised residuals for these data are 3.2 and 3.6. The point with the largest residual is also moderately influential, having a Cook's D-value of 0.51. Nonetheless, all data were retained in the model.

The normality of the residuals was tested with the D'Agostino–Pearson and Shapiro–Wilk tests. p-values were 0.020 and 0.051, respectively, but the low p-values were the result of one or two outlying points. When the most extreme outlier was removed, the resulting p-values were 0.33 and 0.45, respectively, indicating that the remaining data have a normal distribution.

The homoscedasticity of the residuals was tested with the Breusch–Pagan and Goldfeld–Quandt tests giving p-values of 0.74 and 0.99, respectively. Thus, the residuals exhibit constant variance.

The variance inflation factors were 1.00, indicating no problem with collinearity of variables.

In addition, there is no evidence that the model is inappropriate.

The saturation response model




Figure 9: Residuals plotted against predictor variables and predicted values of the response variable




Figure 10: Normality plot of residuals

The plots of Figures 9 and 10 show no problems with the regression assumptions, although there is a hint of decreasing variance with increasing value of TDP.

The normality of the residuals was tested with the D’Agostino–Pearson and Shapiro–Wilk tests, giving p-values of 0.098 and 0.28, respectively, indicating normal distributions.

As there are no readily available tests for homoscedasticity of residuals in non-linear regression, the variance of the residuals was examined by doing linear regression of the absolute values of the residuals against the predictor and response variables. p-values from these analyses were ≤ 0.50 , indicating no problems with non-constant variance. The appearance of larger variance at low TDP values is probably the result of the larger number of data at low TDP.

Again, there is no evidence that the model is inappropriate.


Appendix K. Data extracted from the selected studies for the trivariate saturation response model

Study	Total dietary zinc mean (mg/day)	Total dietary phytate mean (mg/day)	Total absorbed zinc mean (mg/day)	n	Sex
Hambidge et al. (2004)	8.3	1 370	2.37	6	6M, 4F
	10	3 460	1.51	4	6M, 4F
	10.1	2 700	1.44	4	6M, 4F
Knudsen et al. (1996)	10.2	660	3.0	8	5M, 3F
Hunt J et al. (1992)	14	670	3.1	14	14M
	7.8	420	2.3	14	14F
Hunt et al. (1995)	13	1 045	3.6	14	14F
	6.7	1 045	2	14	14F
Hunt et al. (1998)	9.1	1 656	2.4	21	21F
	11.1	542	3.7	21	21F
Wada et al. (1985)	16.4	688	4.1	6	6M
	5.5	688	2.7	6	6M
Lowe et al. (1997)	7.1	585	2.1	6	6F
Adams et al. (2002)	4.3	738	1.3	5	2M, 3F
	5	1 820	0.85	5	2M, 3F
Sian et al. (1996)	5.2	552	1.6	10	10F
	8.1	794	2.8	10	10F
Pinna (1999)	4.6	254	2.2	7	7M
Turnlund et al. (1984)	15	0	5.1	4	4M
	15	2 343	2.62	4	4M
Kristensen et al. (2006)	9.4	845	2.6	16	16F
	9.9	845	2.7	16	16F
	7.5	766	1.8	16	16F
Kim et al. (2007)	6.87	1 623	1.7	7	7F
	6.87	690	3.2	7	7F
	6.47	1 713	1.5	10	10F
	6.47	760	2.4	10	10F
Rosado et al. (2009)	3.91	645	1.48	12	12F
	6.56	771	2.03	12	12F
	7.89	2 218	1.56	14	14F
	13.6	2 376	2.08	14	14F
Sheng et al. (2009)	11.7	835	3.0	21	21F
Hunt et al. (2008)	4.8	326	2.4	8	19M, 20F
	7.4	297	3	8	19M, 20F
	10.9	305	3.9	8	19M, 20F
	15.6	311	4.9	7	19M, 20F
	18.1	285	4.2	8	19M, 20F
	6.2	911	2.4	9	23M, 21F
	9.3	1 726	2.5	9	23M, 21F
	11.9	2 748	2.5	8	23M, 21F
	17.8	3 584	2.8	9	23M, 21F
	21	3 728	3.1	9	23M, 21F
	4.3	292	1.8	8	8F
	6.8	273	2.7	6	6F
	9.4	263	3.5	4	4F
13.3	265	3.9	4	4F	

Study	Total dietary zinc mean (mg/day)	Total dietary phytate mean (mg/day)	Total absorbed zinc mean (mg/day)	n	Sex
Hunt et al. (2008) (continued)	15.6	246	4.6	4	4F
	4.8	326	3.3	8	19M, 20F
	7.4	297	3.4	8	19M, 20F
	10.9	305	4.1	8	19M, 20F
	15.6	311	5	7	19M, 20F
	18.1	285	4.2	8	19M, 20F
	6.2	911	2.6	9	23M, 21F
	9.3	1 726	2.7	9	23M, 21F
	11.9	2 748	2.7	8	23M, 21F
	17.8	3 584	3	9	23M, 21F
	21	3 728	3.1	9	23M, 21F
	4.3	292	2.6	8	8F
	6.8	273	3.3	6	6F
	9.4	263	4	4	4F
	13.3	265	3.6	4	4F
15.6	246	5.1	4	4F	
Chung et al. (2008)	11	941	3.91	9	9M
	4	361	2.41	9	9M
	11	941	3.9	9	9M
	4	361	2.73	9	9M
	11	941	3.24	9	9M
	4	361	2.31	9	9M
Hunt and Beiseigel (2009)	11.5	483	3.8	10	10F
	11.3	1 781	3.0	10	10F
	11.4	391	4.5	10	10F
	12.2	1 789	3.2	10	10F

M, males; F, females.

Appendix L. Three-dimensional representation of Figure 1



ABBREVIATIONS

Afssa	Agence française de sécurité sanitaire des aliments
AI	Adequate Intake
AR	Average Requirement
BMI	body mass index
COMA	Committee on Medical Aspects of Food Policy
CV	coefficient of variation
D–A–CH	Deutschland–Austria–Confoederatio Helvetica
DH	UK Department of Health
DRV	Dietary Reference Value
EAR	Estimated Average Requirement
EFZ	Endogenous faecal zinc
EU	European Union
EURRECA	EUROpean micronutrient RECommendations Aligned
F	female
FAO	Food and Agriculture Organization
FAZ	fractional absorption of zinc
IAEA	International Atomic Energy Agency
IOM	US Institute of Medicine of the National Academy of Sciences
IZiNCG	International Zinc Nutrition Consultative Group
M	male
MRE	metal-response element
MTF	MRE-binding transcription factor
NNR	Nordic Nutrition Recommendations
PRI	Population Reference Intake
RDA	Recommended Dietary Allowance
REE	resting energy expenditure
RNI	Reference Nutrient Intake

SCF	Scientific Committee for Food
TAZ	total absorbed zinc
TDP	total dietary phytate
TDZ	total dietary zinc
UNICEF	United Nations Children's Fund
UL	Tolerable Upper Intake Level
WHO	World Health Organization