

Technical University of Denmark



Exploring the planetary boundary for chemical pollution

Diamond, Miriam L.; de Wit, Cynthia A.; Molander, Sverker; Scheringer, Martin; Backhaus, Thomas; Lohmann, Rainer; Arvidsson, Rickard; Bergman, Åke; Hauschild, Michael Zwicky; Holoubek, Ivan; Persson, Linn; Suzuki, Noriyuki; Vighi, Marco; Zetzsch, Cornelius

Published in:
Environment International

Link to article, DOI:
[10.1016/j.envint.2015.02.001](https://doi.org/10.1016/j.envint.2015.02.001)

Publication date:
2015

Document Version
Peer reviewed version

[Link back to DTU Orbit](#)

Citation (APA):
Diamond, M. L., de Wit, C. A., Molander, S., Scheringer, M., Backhaus, T., Lohmann, R., ... Zetzsch, C. (2015). Exploring the planetary boundary for chemical pollution. *Environment International*, 78, 8-15. DOI: 10.1016/j.envint.2015.02.001

DTU Library

Technical Information Center of Denmark

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

1 Exploring the planetary boundary for chemical pollution

2

3

4 Miriam L. Diamond^{†*}, Cynthia A. de Wit[‡], Sverker Molander,[§] Martin Scheringer,[%] Thomas
5 Backhaus,^{||} Rainer Lohmann,[√] Rickard Arvidsson,[§] Åke Bergman,[⊥] Michael Hauschild,[#] Ivan
6 Holoubek,[¶] Linn Persson,[&] Noriyuki Suzuki,[@] Marco Vighi,[□] Cornelius Zetzsch^Δ

7

8 [†] Department of Earth Sciences, University of Toronto, 22 Russell Street, Toronto, M5S 3B1
9 Ontario, Canada

10 [‡] Department of Environmental Science and Analytical Chemistry (ACES), Stockholm
11 University, SE-106 91 Stockholm, Sweden

12 [§]Environmental Systems Analysis, Department of Energy and Environment, Chalmers
13 University of Technology, SE-412 96 Gothenburg, Sweden

14 [%]Institute for Chemical and Bioengineering, ETH Zürich, Wolfgang-Pauli-Str. 10, 8093 Zürich,
15 CH-8093, Switzerland, and Leuphana University Lüneburg, D-21335 Lüneburg, Germany

16 ^{||}Department of Biological and Environmental Sciences, University of Gothenburg, Box 100,
17 SE-405 30 Gothenburg, Sweden

18 [√]Graduate School of Oceanography, University of Rhode Island, South Ferry Road,
19 Narragansett, Rhode Island, 02882, United States

20 [⊥]Department of Materials and Environmental Chemistry, Stockholm University, SE-106 91
21 Stockholm, Sweden

22 [#]Department of Management Engineering, Technical University of Denmark (DTU), Nils
23 Koppels Allé, Building 426 D, DK-2800 Kgs. Lyngby, Denmark

24 [¶]Research Centre for Toxic Compounds in the Environment (RECETOX), Faculty of Science,
25 Masaryk University, Kamenice 753/5, 625 00 Brno, Czech Republic

26 [&]Stockholm Environment Institute, Linnégatan 87D, Box 24218, Stockholm, Sweden

27 [@]Strategic Risk Management Research Section, Center for Environmental Risk Research,
28 National Institute for Environmental Studies, 16-2 Onogawa, Tsukuba, Ibaraki 305-8506, Japan

29 [‡]Department of Earth and Environmental Sciences, University of Milano Bicocca, Piazza della
30 Scienza 1, Milan, 20126 Italy

31 [^]Forschungsstelle für Atmosphärische Chemie, Dr. Hans-Frisch-Str. 1-3, Universität Bayreuth,
32 D-954 48 Bayreuth, Germany

33

34

35

36 ABSTRACT (323 words)

37 Rockström et al. (2009a, 2009b) have warned that humanity must reduce anthropogenic impacts
38 defined by nine planetary boundaries if “unacceptable global change” is to be avoided.
39 Chemical pollution was identified as one of those boundaries for which continued impacts could
40 erode the resilience of ecosystems and humanity. The central concept of the planetary boundary
41 (or boundaries) for chemical pollution (PBCP or PBCPs) is that the Earth has a finite
42 assimilative capacity for chemical pollution, which includes persistent, as well as readily
43 degradable chemicals released at local to regional scales, which in aggregate threaten ecosystem
44 and human viability. The PBCP allows humanity to explicitly address the increasingly global
45 aspects of chemical pollution throughout a chemical’s life cycle and the need for a global
46 response of internationally coordinated control measures. We submit that sufficient evidence
47 shows stresses on ecosystem and human health at local to global scales, suggesting that
48 conditions are transgressing the safe operating space delimited by a PBCP. As such current local
49 to global pollution control measures are insufficient. However, while the PBCP is an important
50 conceptual step forward, at this point single or multiple PBCPs are challenging to operationalize
51 due to the extremely large number of commercial chemicals or mixtures of chemicals that cause
52 myriad adverse effects to innumerable species and ecosystems, and the complex linkages
53 between emissions, environmental concentrations, exposures and adverse effects. As well, the
54 normative nature of a PBCP presents challenges of negotiating pollution limits amongst societal
55 groups with differing viewpoints. Thus, a combination of approaches is recommended as
56 follows: develop indicators of chemical pollution, for both control and response variables, that
57 will aid in quantifying a PBCP(s) and gauging progress towards reducing chemical pollution,
58 develop new technologies and technical and social approaches to mitigate global chemical

59 pollution that emphasize a preventative approach, coordinate pollution control and sustainability
60 efforts, and facilitate implementation of multiple (and potentially decentralized) control efforts
61 involving scientists, civil society, government, non-governmental organizations and international
62 bodies.

63 **KEYWORDS:** planetary boundary, chemical pollution, chemical emissions, Stockholm
64 Convention, tipping point, global threshold, pollution controls, ecosystem health protection,
65 human health protection, chemical management

66 **1. INTRODUCTION**

67 Rockström et al. (2009a, 2009b) presented nine anthropogenic impacts of global relevance,
68 including climate change, biodiversity loss, anthropogenic changes of the nitrogen and
69 phosphorus cycles, stratospheric ozone depletion, ocean acidification, global freshwater use,
70 changes in land use, atmospheric aerosol loading, and chemical pollution. The authors proposed
71 that humanity may be moving beyond a “safe operating space” as the magnitude of these impacts
72 approach or exceed certain thresholds that represent tipping points of the global system or a
73 natural limit for processes without clear thresholds (so-called “dangerous levels” in the
74 Rockström et al. articles) (Fig. 1). As discussed in detail below, the authors defined a “safe
75 operating space” as those global conditions that allow for continued human development.
76 Rockström et al. (2009a, 2009b) challenged the global scientific community to determine these
77 “non-negotiable” thresholds or natural limits, which are science-based limits of the Earth’s
78 systems, reflecting conditions that are favorable for human life and cultural development, and
79 then to define human-determined boundaries at an appropriate distance from these limits that
80 allow humanity to “avoid unacceptable global change” (Carpenter and Bennett, 2011). A critical

81 goal of defining the boundaries is to move governance and management away from a piecemeal
82 and sectorial approach, towards an integrated global approach that is necessary to address global
83 phenomena.

84

85 For chemical pollution, Rockström et al. (2009a, 2009b) did not define the scope of chemicals
86 considered, natural limits or a planetary boundary, but stated that these remain to be determined.
87 However, they suggested that possible measurable control variables for natural limits could be
88 emissions, concentrations or effects of Persistent Organic Pollutants (POPs), plastics, endocrine
89 disruptors, heavy metals and nuclear wastes. Persson et al. (2013) added to the discussion by
90 suggesting three conditions that must be met simultaneously for chemical pollution to present a
91 global threat. Here we consider a broad range of chemicals including synthetic organic
92 substances and metals, and those intentionally and unintentionally released. We do not consider
93 the nutrients nitrogen and phosphorus that are considered under a separate planetary boundary, or
94 sulfates that can also fall under another planetary boundary (atmospheric aerosol loading).

95

96 A large primary literature and numerous reviews document the extent and diversity of chemical
97 pollution and attendant adverse health effects to humans and ecosystems (e.g., UNEP, 2012;
98 AMAP, 2004, 2009; Letcher et al., 2010; WHO and UNEP, 2013; *inter alia*). Indeed, the
99 number of scientific studies providing such evidence fills environmental journals and conference
100 halls. Examples of widespread effects are diminishing populations of wildlife (e.g., Oaks et al.,
101 2004; Tapparo et al., 2012; EFSA, 2013) and increasing burdens of human clinical and

102 subclinical illness related to environmental toxicants (WHO and UNEP, 2013; Grandjean and
103 Landrigan, 2006; Stillerman et al., 2008). Mounting evidence also indicates that the assessment
104 of individual chemicals is insufficient, as complex mixtures might cause significant toxic effects,
105 even if all individual chemicals are present only at individually non-toxic concentrations, as
106 discussed below. This pattern has been observed repeatedly in a broad range of bioassays at
107 different levels of complexity and for different types of chemicals (see reviews by Kortenkamp
108 et al., 2007, 2009; Kortenkamp, 2008; Backhaus et al., 2010; SCHENIHR et al., 2012).
109 Together, this evidence implies that if emissions of increasing numbers and amounts of
110 chemicals continue at current and anticipated increasing rates (UNEP, 2012), concentrations of
111 such chemicals in many parts of the world, alone or as mixtures, will push the global system
112 beyond the safe operating space. In turn, reaching this point will lead to erosion of vital
113 ecosystems and ecosystem services, and threaten human well-being. Some argue that this point
114 has already been reached (WHO and UNEP, 2013; *inter alia*). Furthermore, the boundary of
115 global chemical pollution cannot be ignored because it is inextricably connected to the other
116 planetary boundaries by the manifold impacts across the life-cycle of chemicals at a global scale,
117 e.g., energy and water use for extraction and manufacturing, land use change that accompanies
118 waste disposal with a potential loss of biodiversity.

119

120 This paper explores the definitions and meaning of, and arguments for, a planetary boundary or
121 boundaries for chemical pollution (PBCP). We discuss the many challenges that indicate that
122 defining a boundary or boundaries for chemical pollution is not easily within reach. Our intent
123 here is not to reproduce or re-summarize evidence of widespread adverse effects due to chemical

124 pollution. Rather, we submit that this evidence points to the need for considering a planetary
125 boundary or more likely *boundaries* for chemical pollution to help humanity remain within the
126 Earth's safe operating space. Thus, the paper closes with recommendations for steps that
127 hopefully will move humanity towards a safe operating space with respect to chemical pollution.

128

129 We start the discussion by acknowledging that defining natural limits and a PBCP(s) is
130 challenging for many reasons. In the framework presented by Rockström et al. (2009a, 2009b),
131 defining a PBCP is more difficult than for other planetary boundaries (e.g. for global warming),
132 due to the difficulty of identifying a single or a few measurable control variables. A control
133 variable is defined, according to Rockström et al. (2009a, 2009b), as a measurable parameter
134 that can be related to a specific planetary boundary, e.g., atmospheric CO₂ or temperature for
135 global warming. However, agreeing on one or more control variables for chemical pollution is
136 challenging because chemical pollution is caused by an enormous number of chemicals emitted
137 from innumerable sources and in extremely different amounts in different regions of the world.
138 In the same way, the response variable is difficult to define and measure in a clear-cut way, since
139 chemicals cause a wide variety of adverse effects in a similarly wide variety of species, including
140 humans. The links to the related boundary of biodiversity are evident (Steffen et al. 2015). The
141 critical point is that the Earth's assimilative capacity, or the number and capacities of the sinks
142 capable of degrading or immobilizing anthropogenically-released chemicals, is limited at the
143 global level, even for readily biodegradable chemicals.

2. WHY A PLANETARY BOUNDARY FOR CHEMICAL POLLUTION?


144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165

Several policy instruments aimed at controlling chemical pollution have been developed and are in varying degrees of implementation (Table S1). How does a PBCP differ from existing instruments for chemical management and how or why might it be useful rather than redundant? In order to answer these questions we first expand on the concept of planetary boundaries and a “safe operating space” introduced by Rockström et al. (2009a, 2009b) and then move to put a PBCP into the context of existing instruments for chemicals management.

Rockström et al. (2009a, 2009b) identified that several Earth processes and subsystems behave non-linearly, with thresholds that, once crossed, could tip them into new, undesirable states. For these processes, a sharp “tipping point” may exist beyond which the system may transition into a qualitatively different stage, such as much more rapid global warming at CO₂ concentrations above a certain value (Fig. 1a). Examples of Earth systems with such global thresholds or tipping points include the global climate and ocean acidification (e.g., Lenton et al., 2008; Doney et al., 2009; 2014). The planetary boundary can then be set at a level somewhere below the tipping point.

Other processes and subsystems may not have sharp thresholds (Fig. 1b), but their continued erosion or depletion at continental to global scales may cause functional collapse in an increasing number of globally interconnected systems. Here, examples are freshwater use, land use change and loss of biodiversity (May, 1977; Gerten et al., 2013; Baronsky et al., 2012; Brook et al.,

166 2013). For these, the planetary boundary can be set at a level where the risk of functional
 167 collapse is deemed acceptably low. In aggregate, planetary boundaries may thus be defined as a
 168 set of critical values for one or several control variables defined by humans to be at a safe
 169 distance from such thresholds or dangerous levels (if no threshold is evident) that, if crossed,
 170 could lead to abrupt global environmental change. The domain below the boundary can be
 171 considered a “safe operating space”.



172

173 Figure 1. Illustration of the concept of the planetary boundary (a) for phenomena with a clear
 174 tipping point or threshold, where the system moves into a new state, such as CO₂-driven climate
 175 change, and (b) without a tipping point, where the system is constantly eroded (modified figure
 176 from Rockström et al. (2009a), reprinted with permission of the Stockholm Resilience Center,
 177 Stockholm University, Sweden). We suggest that aggregated chemical pollution is illustrated by
 178 (b) where there is no clear tipping point.

179

180

181 Although the intention was to define planetary boundaries for systems or processes affecting the


182 Earth at the global scale, Rockström et al. (2009a, 2009b) recognized that many of the identified

183 boundaries have thresholds that are more evident at local and/or regional scales where
184 disturbance is concentrated or the affected ecosystem is more sensitive. These were identified as
185 “slow processes without known global scale thresholds”. As such, they become a global
186 problem when they occur at many sites at the same time, aggregating to a level that undermines
187 the resilience of ecosystems or that adversely affects human health. In turn, these effects would
188 make it more likely that a threshold with global consequences will be crossed. Examples include
189 biodiversity loss, land use change, global nitrogen and phosphorus biogeochemical cycles, and
190 chemical pollution (Erisman et al., 2013; Hooper et al., 2012; Diaz and Rosenberg, 2008). Slow
191 processes without global thresholds may also exert their effects by affecting other planetary
192 boundaries, for example, chemical pollution of ecosystems linked to biodiversity loss
193 (Voeroesmartly et al., 2010; Lenzen et al., 2012; Steffen et al. 2015). For example, chemical
194 pollution can increase the vulnerability of ecosystems to species loss and land-use change,
195 notably deforestation, can increase terrestrial-based chemical loadings to surface waters.

196

197 The distance between the planetary boundary and the threshold or natural limit ideally depends
198 on the uncertainty that surrounds the scientific knowledge about the threshold or natural limit
199 (Fig. 2). If the uncertainty is high, a larger distance between the threshold and the boundary is
200 advisable.

201



202

203 Figure 2. Illustration of where global impacts are located with respect to the safe operating space.

204

205 For the planetary boundaries where critical limits were estimated, most of these could be based
 206 on one or two specific control variables, such as atmospheric CO₂ concentrations and radiative
 207 forcing for climate change. Most of the planetary boundaries that were quantified are
 208 preliminary, rough estimates with large uncertainties and for which knowledge gaps were
 209 acknowledged.

210

211 Although some preliminary boundaries have been proposed, Rockström et al. (2009a, 2009b)
 212 pointed out the normative quality of a “safe” distance, as it is based on how societies deal with
 213 risk and uncertainty. By normative we mean that decisions on what constitutes a “safe operating
 214 space” are societal decisions, supported by scientific evidence. This implies that the diversity of
 215 viewpoints held by different societal groups have to be heard in order to come to a decision on
 216 what constitutes a safe operating space.

217

218 What does the PBCP offer that existing pollution control instruments lack? The planetary
219 boundary concept allows us to explicitly address the *global aspects of chemical pollution*. By
220 recognizing the global nature of chemical pollution, including aggregated local effects or where
221 distance separates emissions from effects, we highlight the need for an integrated global response
222 and acknowledge that pollution control activities of local to national entities alone, are
223 insufficient.

224
225 Chemical pollution is a global issue. Several groups of chemicals are distributed around the
226 globe by virtue of their persistence and ability to undergo long-range transport, for example
227 chlorofluorocarbons (CFCs) and persistent organic pollutants (POPs). Others, such as high-
228 production-volume metals that are inherently persistent, are used and emitted globally because of
229 their high production volumes, global trade and widespread use in a broad range of applications.
230 Additionally, the global economy is undergoing chemical “intensification”, as described by the
231 UNEP “Global Chemicals Outlook” analysis (UNEP, 2013). Chemical intensification is due to
232 rapidly increasing global production of chemicals (Wilson and Schwarzman, 2009), to the
233 increasing use of synthetic substances to replace natural materials, and to the use of increasingly
234 complex chemicals in more and more applications. Chemical intensification is predicted to lead
235 to increasing per-capita chemical usage amongst a growing global population (UNEP, 2013).

236
237 In addition, chemical product chains, which span the life cycle stages from resource extraction to
238 product manufacturing, use and disposal, are increasing in complexity, often covering several
239 continents and decades of time, and offer new challenges to pollution control. For example,
240 chemical production today can result in future emissions, particularly for chemicals in

241 infrastructure and goods with long lifetimes. Brunner and Rechberger (2001) have estimated that
242 whereas ~10% of all chemical stocks is contained in waste deposits from primary production and
243 ~10% is contained in land filled waste, ~80% is contained in in-use and “hibernating” stocks.
244 Most documentation of uncontrolled releases concern the two former sources (i.e., 20%) but not
245 the 80% (e.g., Brunner and Rechberger, 2001; Weber et al., 2013; *inter alia*). Examples of the
246 “20%” include long-term emissions from tailings, waste rock piles, nuclear waste repositories,
247 abandoned industrial sites, and numerous landfills in developing countries (Turk et al., 2007;
248 Torres et al., 2013; Weber et al., 2011). One example of long-term emissions from an in-use
249 chemical stock is that of polychlorinated biphenyls (PCBs, listed as a POP under the Stockholm
250 Convention) from equipment that was still in use in Canada in 2006 despite the ban on PCB
251 production nearly 40 years ago (Diamond et al., 2010; Csiszar et al., 2013). Another example is
252 that of CFCs contained in blown building insulation that is subject to uncontrolled releases as the
253 generation of buildings using that foam undergoes renovation or destruction over the next 30
254 years (Brunner and Rechberger, 2001)

255
256 Similar application patterns of chemical technologies and similar uses of chemical products in
257 almost all regions of the world result in widespread chemical releases. Chemical manufacturing
258 and industrial usage are rapidly shifting from Western industrialized countries to developing
259 countries and countries with economies in transition, including BRICS countries (Brazil, Russia,
260 and especially India and China, and most recently South Africa) (UNEP, 2013). New and
261 increasing resource extraction and chemical manufacturing, usage and waste disposal are leading
262 to increased chemical pollution, particularly in jurisdictions with insufficient control mechanisms
263 (Schmidt, 2006; Gottesfeld and Cherry, 2011). Short-lived chemicals are also being released in

264 many regions at rates that exceed degradation rates and hence environmental assimilative
265 capacities. Examples of such chemicals include pharmaceuticals, high production volume
266 plastics and plasticizers such as bisphenol A and di-ester phthalates, and “D4” and “D5”
267 siloxanes (e.g., WHO and UNEP, 2013; Kolpin et al., 2002; Rosi-Marshall et al., 2013; Peck and
268 Hornbuckle, 2004; Fromme et al., 2002; Fries and Mihajlovic, 2011; Wang et al., 2013).

269
270 As pointed out above, the global nature of chemical pollution demands a global response of
271 internationally coordinated control measures, in addition to multiple local, regional and national
272 efforts covering different groups of substances, which are disconnected in time and space. One
273 example of a global governance instrument is the Stockholm Convention on Persistent Organic
274 Pollutants (POPs), which seeks elimination at best, or more broadly, the sound management, of a
275 set of POPs agreed upon through international negotiations (Stockholm Convention, 2008).
276 While achieving many successes (Stockholm Convention, 2012), the Convention is limited to a
277 small number of chemicals or chemical classes (currently 22 are listed, with four more under
278 review), includes numerous exemptions, and has no instrument for sanctions to ensure national
279 implementation. This is not a shortcoming of the Convention because the intention of the
280 Convention is not to address the totality of chemical pollution. As such, the Stockholm
281 Convention is not adequate for challenge presented by developing a PBCP. Similarly, the
282 Montreal Protocol is limited to substances that deplete the stratospheric ozone layer (UNEP
283 2010-2011) and the Minamata Convention is limited to mercury (UNEP 2015). The Convention
284 on Long-range Transboundary Air Pollution, under the aegis of the United Nations Economic
285 Commission for Europe and to which there are 51 parties, addresses a range of chemical
286 pollutants including metals and POPs (UNECE 2004).

287

288 Another example of a global governance tool is the United Nations Framework Convention on
289 Climate Change where global negotiations and agreements have led to reduction goals for
290 greenhouse gases that are intended to be implemented at national levels (UNFCCC, 2013).
291 International climate negotiations have seen the emergence of control instruments of largely two
292 types. The first is an absolute limit for total CO₂-equivalent emissions (a “cap”) to assure that
293 total global emissions are on target to prevent the global atmospheric CO₂ concentration
294 exceeding an agreed-upon boundary. The second type of control scheme links emissions to
295 activity or intensity such as CO₂-equivalent emissions per unit of electricity generated or per
296 kilometre driven, or to an economic cost resulting in reductions of CO₂-equivalent
297 emissions/capita (Azar and Rodhe, 1997; Ellerman and Sue Wing, 2003). These intensity or
298 efficiency-based emission controls acknowledge the need to reduce greenhouse gas emissions
299 but cannot ensure that global emissions are within the global safe operating space because of
300 population and economic growth that increase the demand for energy services, most of which are
301 based on fossil fuels (IEA, 2014).

302

303 Implicit in the concept of a safe operating space for CO₂ and other greenhouse gases, ocean
304 acidification, nitrogen and phosphorus cycles, and “chemical pollution”, is that there is a finite
305 global assimilative capacity. Here we define assimilative capacity as the ability of an ecosystem
306 to render substances harmless, i.e. avoiding adverse effects. By seeing the problem in this light,
307 it leads us towards exploring the need for a globally coordinated cap for emissions, rather than
308 jurisdiction-specific, intensity-based controls, which may be sufficient in some circumstances but
309 fail to account for cumulative, global effects.

310

311 **3. CHALLENGES OF DEFINING A PLANETARY BOUNDARY FOR** 312 **CHEMICAL POLLUTION**

313 Moving the idea of a PB beyond a conceptual model requires that the impact of anthropogenic
314 stressor(s) on all ecosystems can be described and quantified as a function of a measurable
315 control variable(s) that is (are) related to a measurable response variable(s). For a PBCP, the
316 ultimate effect or response variable (Fig. 1) subject to control is widespread adverse impact(s) to
317 ecological and/or human health caused by exposure to (a) substance(s). Exposure can be
318 identified as the critical control variable since it is the necessary prerequisite for any kind of
319 chemically induced effect or response we want to safeguard against. Ideally, chemical exposure
320 can be used to define a threshold(s) or natural limit(s) that, in turn, can be translated into a global
321 boundary (boundaries) and a safe operating space. As noted above, the boundary (boundaries) is
322 (are) established by humans and is (are) a product of societal demands, needs, value judgments
323 and negotiations. The control variable(s) must also be amenable to translation into possible
324 mitigation or control activities, which in this case would reduce exposure and thus, would
325 maintain human and ecosystem health within the safe operating space, the latter reflected in
326 maintained biodiversity, ecosystem functionality and human health.

327

328 Challenges arise at all stages in the definition process that starts with a control variable(s) and
329 ends with “actionable” activities. First, operationalizing “exposure” as the control variable is
330 difficult because of the high and poorly defined number of chemicals that fall under the umbrella
331 of “chemical pollution”. More than 100 000 substances are in commerce (Egeghy et al., 2012),

332 including pesticides, biocides and pharmaceuticals, industrial chemicals, building materials and
333 substances in personal care products and cosmetics (e.g., Howard and Muir, 2010, 2011; ECHA,
334 2013) and very few of them have undergone adequate risk assessment for adverse effects. A
335 recent screening of 95 000 chemicals for persistence (P), bioaccumulation (B) and toxicity (T)
336 properties (REACH criteria) identified 3% or approximately 3000 chemicals as potential PBT
337 chemicals (uncertainty range of 153-12 500 chemicals) (Stempel et al., 2012). Similarly, 93 000
338 chemicals were screened for P, B and long range transport potential according to the Stockholm
339 Convention criteria, plus T (REACH criteria) resulting in the identification of 510 potential
340 POPs (uncertainty range of 190-1 200 chemicals) (Scheringer et al., 2012). Unintentionally
341 produced substances, such as the combustion by-products polycyclic aromatic hydrocarbons
342 (PAH) and polychlorinated and polybrominated dibenzo-*p*-dioxins and furans (PCDD/F and
343 PBDDs/Fs), are emitted as a consequence of human activity and many emitted chemicals are
344 transformed to a multitude of other chemicals by biological and physical-chemical processes.
345 Whereas some limits have been placed on a few selected chemicals that are highly persistent,
346 bioaccumulative and toxic such as PCDD/F, those with intermediate PBT properties have
347 received insufficient attention (Muir and Howard, 2006; Howard and Muir, 2010; Scheringer et
348 al., 2012). In addition, an enormous number of organisms in a diversity of ecosystems are
349 exposed to chemical pollution (which is invariably a complex chemical mixture) and they will
350 respond in myriad ways. Moreover, chemicals have specific modes of actions and can show
351 very different toxicological potencies. Humans take a specific place among affected organisms.
352 Any approach to establishing a PBCP(s) must include impacts on human health, even if this is in
353 contrast to the framework of Rockström et al. (2009a, 2009b) or which the objects of protection

354 are biogeochemical systems and ecosystems, e.g., the climate system, the ozone layer, and
355 freshwater.

356

357 Second, we acknowledge that boundaries for chemical pollution have been developed at a global
358 scale for selected POPs and mercury, and at local and regional scales for chemicals in foods,
359 water and air (Table S1). However, only a few of these boundaries account for exposure to
360 multiple chemicals simultaneously that can act in an additive fashion. Moving beyond a
361 chemical-by-chemical approach to acknowledge mixture effects is of growing importance if
362 limits are to be protective (e.g., Kortenkamp, 2007; Kortenkamp et al., 2007; Backhaus et al.,
363 2010; Meek et al., 2011; SCHENIHR et al., 2012). An increasing body of evidence suggests
364 that, *de facto*, the existing boundaries are not sufficiently protective for endocrine disrupting
365 chemicals that can cause transgenerational effects (e.g., Baccarelli and Bollati, 2009; Bollati and
366 Baccarelli, 2010; Bouwman et al., 2012; Mani et al., 2012; WHO and UNEP, 2013; *inter alia*).
367 This is not surprising since accepted and validated methods for identifying and testing endocrine
368 disrupting chemicals, particularly after exposure during critical early life stages, are generally
369 lacking or have not yet been implemented in chemicals risk assessment (WHO and UNEP, 2013;
370 *inter alia*).

371

372 Third, connecting exposure as the control variable to an “actionable” activity (such as controlling
373 emissions) is difficult because of the diversity of fate and transformation processes at play
374 between an initial emission of a chemical or a chemical mixture and the concentration(s)

375 resulting in exposure and then an adverse effect. Establishing the release-fate-concentration-
376 effect linkage is necessary for other planetary boundaries such as CO₂, stratospheric ozone,
377 phosphorus and nitrogen cycles. Establishing this linkage for chemical pollution is also
378 necessary but it is more challenging because of the large number of chemicals of varying
379 persistence and toxicity that are captured by this boundary.

380

381 Finally, in addition to the scientific challenges of defining a boundary(s), it must be remembered
382 that most of the world's countries do not have the capacity or resources to measure a control
383 variable such as exposure and to implement effective controls such as those listed in Table S1
384 (e.g., Klanova et al., 2009; Adu-Kumi et al., 2012). Furthermore, as noted above, a boundary(s)
385 is normative and as such, a diversity of viewpoints will be held on what constitutes an
386 "acceptable" level of pollution.

387

388 The combination of numerous substances with different use and emission patterns, affecting a
389 multitude of different endpoints in a plethora of exposed species in the vastly different
390 ecosystems of the world, plus consideration of human health, makes the derivation of a single
391 quantitative PBCP or multiple PBCPs a daunting, if not impossible task. However, the situation
392 of increasing chemical production, emissions and adverse effects cannot be allowed to continue
393 unabated. Thus, we believe that the concept of a planetary boundary or boundaries for chemical
394 pollution is a useful framework for global action, but that it needs to be modified to account for
395 these complexities and challenges.

396

397

4. STEPS TOWARD GLOBAL CHEMICALS MANAGEMENT

398

399 Although it may not be possible to establish a single or even multiple PBCP(s) at this time, an
400 increasing body of evidence strongly suggests that we need more effective global chemicals
401 management. What has been accomplished in global chemicals management? Global
402 cooperation amongst nations has, amongst others, resulted in the Stockholm Convention on
403 POPs, the Montreal Protocol on CFCs, the Basel Convention on Control of Transboundary
404 Movements of Hazardous Wastes, and the Rotterdam Convention on Prior Informed Consent
405 Procedure for Certain Hazardous Chemicals and Pesticides in International Trade. These
406 Multilateral Environmental Agreements have come together under the aegis of UNEP. The
407 Stockholm and Montreal agreements strive towards zero-emissions of the listed chemicals. In
408 January 2013, UNEP brokered the Minamata Convention on mercury, the language of which has
409 gained support from 94 signatory countries (UNEP, 2015). The Minamata Convention specifies
410 the banning of production, export and import of a range of mercury-containing products, calls for
411 the drafting of strategies to limit the use of mercury in artisanal and small-scale gold mining, and
412 aims to work towards minimizing mercury emissions from combustion sources such as
413 conventional fossil fuel power plants and cement factories. Like the Stockholm Convention, the
414 Minamata Convention includes the provision to develop a compliance mechanism that will be
415 established through negotiation after the official signing of the Convention.

416

417 These five agreements address priority chemical pollutants at the global scale, reflect the insight
418 that global dilution is not the solution to local or global pollution, and that environmental
419 safeguards are the right of all countries. Well over 100 countries have adopted them (except for
420 the most recent Minamata Convention), which in itself is a great accomplishment. However,
421 these agreements have limitations due to numerous official exemptions and unofficial
422 “loopholes”, they cover only a limited number of chemicals, implementation costs are largely
423 left to individual countries of which many lack such capacity, and sanctions cannot be levied for
424 a lack of compliance. As such, these agreements are not adequate to address the totality of
425 chemical pollution (which was never their intent). Importantly, the fact that these agreements
426 have been enacted is a reflection that humanity has come close to or crossed boundaries for these
427 chemicals. A PBCP provides an overarching conceptual basis to characterize the achievements
428 of these agreements and to accommodate additional necessary controls.

429
430 For chemicals listed by the Stockholm and Minamata Conventions and the Montreal Protocol,
431 the planetary boundary is set at a *de minimus* level (ideally zero emissions but exemptions
432 preclude this). In addition to the zero emissions boundary, several other types of boundaries
433 have been defined during the past decades under many jurisdiction-specific regulations and
434 initiatives spanning local to national scales. As summarized in Table S1, the initiatives, which
435 come from international agencies, Europe, Japan, North America, China, India and Nigeria,
436 include limits to levels of pesticides in groundwater and surface water, levels of priority
437 pollutants in surface waters, and acceptable daily intakes (ADIs) for a wide range of food
438 contaminants. However, as noted above, not all of these agencies are able to monitor for, and
439 enforce compliance.

440

441 Another major global initiative is the Strategic Approach to International Chemicals
442 Management (SAICM), which is also under the aegis of UNEP. The ultimate goal of SAICM is
443 to facilitate activities to ensure that "...chemicals will be produced and used in ways that
444 minimize significant adverse impacts on the environment and human health" (SAICM, 2006).
445 The role of SAICM is advisory by acting as a source of information to governmental and extra-
446 governmental bodies regarding safe chemical management and funding projects to fulfill the aim
447 of the initiative. SAICM is a non-binding agreement with broad participation of countries and
448 other stakeholders such as the chemical industry. In comparison to the five chemical
449 agreements, SAICM is much broader in scope by addressing all agricultural and industrial
450 chemicals from cradle to grave, aiming at overall sound chemicals management. However,
451 SAICM does not have a compliance mechanism.

452

453 To move towards a truly global approach encompassing the aggregated impacts from all
454 anthropogenic chemical pollution, we need to learn from experience and build on successes (and
455 failures). What are the key lessons learned? One lesson learned is that implementation of
456 stringent controls by specific jurisdictions has led to improved local conditions in those
457 jurisdictions. However, increased global trade and the fluidity of global finance have moved
458 more chemical and goods production and waste disposal to locations without stringent controls
459 (e.g., Skelton et al., 2011; Breivik et al., 2011; Sindiku et al., 2014). Thus, one intention of a
460 global boundary is avoiding "pollution free" jurisdictions at the expense of creating "pollution
461 havens" in developing nations (e.g. Gottesfeld, 2013). Examples of developed nations achieving
462 their pollution control goals by shipping waste and waste products to developing nations have

463 been described elsewhere (Schmidt, 2006; Breivik et al., 2011, 2014; Gioia et al., 2011;
464 Abdullah et al., 2013).

465
466 A second lesson learned is that despite the challenges, as scientists we need to avoid calling for
467 more scientific certainty before action is taken as this delays adoption of control measures, which
468 in this case translates to measures that will help stem widespread chemical pollution. Gee and
469 others (Gee, 2006; Gee et al., 2013; Harremoës et al., 2001) have documented examples of where
470 the call for more research to improve risk assessments of chemicals often led to delays in action
471 of up to several decades although early warnings of adverse effects were already apparent (e.g.
472 tobacco smoking and asbestos). Persson et al. (2013) provide a persuasive argument in this
473 regard.

474
475 As a result of these considerations, we submit that the PBCP is a useful aspirational framework
476 that allows natural and social scientists, policy makers, industry and civil society to visualize the
477 idea of a safe operating space, see the limited assimilative capacity of the Earth, recognize
478 chemical pollution at a global scale, and see the inadequacy of current control measures to deal
479 with the totality of global chemical pollution. Having said that, we recognize that defining a
480 single or multiple quantitative PBCP(s), or even a single approach for its definition, is not now
481 within reach. Rather, we recommend advancing in multiple directions that involve globally
482 coordinated action in scientific, technical and political domains (e.g., Conklin, 2005; Horn and
483 Weber, 2007). For the scientific domain we propose the following:

- 484 1. Explore advancing the concept of, and methods for quantifying a PBCP(s). We advocate
485 making stepwise progress using a few well-known chemicals such as POPs, intermediate
486 PBT chemicals (demonstrated toxicity but not highly persistent), and a few high production
487 volume chemicals with demonstrated toxicity.
- 488 2. Continue to identify and develop indicators of global chemical pollution, initially based on
489 proxies for chemical exposure and potency. Information on indicator status should then be
490 used to gauge progress towards staying within the safe operating space for chemical
491 pollution. Useful information to guide this task can be taken from the Drivers, Pressures,
492 States, Impacts, Responses (DPSIR) approach (OECD, 1991; Harremoës, 1998), and
493 suggestions of how this could be accomplished are given in the Supporting information. This
494 proposal builds on the global monitoring networks that have achieved considerable success
495 such as those under the Stockholm Convention (e.g., the Global Atmospheric Passive
496 Sampling network or GAPS (Gawor et al., 2014) and Human milk survey (UNEP et al.,
497 2013)).
- 498 3. Conduct research into new technologies and methods that will aid in implementing the goals
499 of the six global chemical agreements (Montreal Protocol; Stockholm, Minamata, Rotterdam,
500 Basel and UNECE LRTAP Conventions) and in lowering production and emissions of non-
501 POP priority chemicals. This research includes methods for identifying and characterizing
502 stocks of chemicals scheduled for elimination, developing technologies for efficient and
503 effective destruction of stockpiles, research into societal and cultural considerations that will
504 maximize the likelihood of policy implementation, etc.
- 505 4. Connect activities aimed at chemical pollution control in the context of PBCP to efforts
506 aimed at moving towards sustainable resource use. This should include investigating ways to

507 chemically “de-intensify” economies, to use “green chemistry” substitutes and non-chemical
508 solutions, and to implement social solutions aimed at reducing resource consumption.

509 Efforts are underway in this regard, such as the U.S. EPA’s Design for the Environment
510 Program (U.S.EPA, 2014) and the GreenScreen© for Safer Chemicals (Clean Production
511 Action, 2015). These two issues, PBCP and sustainable resource use, are intertwined such
512 that chemical pollution is a manifestation of unsustainable and inefficient resource use.

513 Thus, efforts directed towards achieving both goals would benefit from coordinated action.

514

515 Progressing towards a PBCP(s) will require scientific, political, social and economic strategies.

516 In the political domain, it will be important to raise more awareness for chemical pollution
517 problems in all parts of the world, and to aid individual countries in implementing existing local
518 and regional boundaries and international agreements. The shift of chemical production from
519 OECD countries primarily to the BRICS countries needs to be complemented by a process that
520 helps to develop chemical regulation and enforcement in these regions to a level comparable or
521 better than that of OECD countries.

522

523 To address these needs, organizations at the global level such as WHO and UNEP can be drivers
524 for effective exchange and collaboration amongst the public, environmental NGOs, industry and
525 national government institutions to enable significant pollution control. Civil society and local
526 jurisdictions also have and continue to implement effective pollution controls using a variety of
527 tools. Examples here include the activities of the International POPs Elimination Network

528 (IPEN), the Pesticides Action Network (PAN), and C40 Cities for “Global Leadership on
529 Climate Change” (C40 Cities, 2013).

530

531 In closing, 50 years ago Rachel Carson pointed out for the first time that the extensive use of
532 pesticides is dangerous not only to wildlife, but also to humans. This is still an ongoing concern,
533 emphasized by the recent finding that neonicotinoid pesticides are contributing to the massive
534 collapse of bee populations (Tapparo et al., 2012; Henry et al., 2012; Whitehorn et al., 2012).

535 Now we need to go beyond Rachel Carson’s clarion call about pesticides. Today’s phenomenon
536 of locally to globally distributed chemicals that are causing adverse effects, demands that a wide
537 range of chemical products and uses be restrained and many chemicals in commerce need to be
538 used with much more prudence and precaution. It is time to harness the knowledge, capacity and
539 commitment held by many to see Rachel Carson’s vision moved to a truly global scale.

540

541 **ACKNOWLEDGEMENTS**

542

543

544 The authors gratefully acknowledge financial support by the Swedish Research Council

545 FORMAS and the International Panel on Chemical Pollution, which funded a workshop on this

546 topic.

547

548

549

550 **REFERENCES**

551

- 552 Abdullah, H.M., Mahboob, M.G., Banu, M.R., Seker, D.Z., 2013. Monitoring the drastic growth
553 of ship breaking yards in Sitakunda: a threat to the coastal environment of Bangladesh.
554 Environmental Monitoring & Assessment 185, 3839-3851.
- 555 Adu-Kumi, S., Kares, R., Literak, J., Boruvkova, J., Yeboah, P.O., Carboo, D., Akoto, O.,
556 Darko, G., Osae, S., Klanova, J., 2012. Levels and seasonal variations of organochlorine
557 pesticides in urban and rural background air of southern Ghana. Environmental Science
558 & Pollution Research 19, 1963-1970.
- 559 AMAP (Arctic Monitoring and Assessment Programme), 2004. AMAP Assessment 2002:
560 Persistent Organic Pollutants in the Arctic. Arctic Monitoring and Assessment
561 Programme (AMAP), Oslo, Norway, p. 310.
- 562 AMAP (Arctic Monitoring and Assessment Programme), 2009. AMAP Assessment 2009:
563 Human Health in the Arctic. Arctic Monitoring and Assessment Programme, Oslo,
564 Norway, p. 254.
- 565 Azar, C., Rodhe, H., 1997. Targets for Stabilization of Atmospheric CO₂. Science 276, 1818-
566 1819.
- 567 Baccarelli, A., Bollati, V., 2009. Epigenetics and environmental chemicals. Current Opinion in
568 Pediatrics 21, 243-251.
- 569 Backhaus, T., Blanck, H., Faust, M., 2010. Hazard and Risk Assessment of Chemical Mixtures
570 under REACH—State of the Art, Gaps and Options for Improvement. Swedish Chemicals
571 Agency, Sundbyberg, Sweden.
- 572 Barnosky, A.D., Hadly, E.A., Bascompte, J., Berlow, E.L., Brown, J.H., Fortelius, M., Getz,
573 W.M., Harte, J., Hastings, A., Marquet, P.A., Martinez, N.D., Mooers, A., Roopnarine,
574 P., Vermeij, G., Williams, J.W., Gillespie, R., Kitzes, J., Marshall, C., Matzke, N.,

- 575 Mindell, D.P., Revilla, E., Smith, A.B., 2012. Approaching a state shift in Earth's
576 biosphere. *Nature* 486, 52-58.
- 577 Bollati, V., Baccarelli, A., 2010. Environmental epigenetics. *Heredity* 105, 105-112.
- 578 Bouwman, H., Kylin, H., Yive, N., Tatayah, V., Loken, K., Skaare, J.U., Polder, A., 2012. First
579 report of chlorinated and brominated hydrocarbon pollutants in marine bird eggs from an
580 oceanic Indian Ocean island. *Environmental Research* 118, 53-64.
- 581 Breivik, K., Gioia, R., Chakraborty, P., Zhang, G., Jones, K.C., 2011. Are reductions in
582 industrial organic contaminants emissions in rich countries achieved partly by export of
583 toxic wastes? *Environmental Science & Technology* 45, 9154-9160.
- 584 Breivik, K., Armitage, J.M., Wania, F., Jones, K.C., 2014. Tracking the Global Generation and
585 Exports of e-Waste. Do Existing Estimates Add up? *Environmental Science &*
586 *Technology* 48, 8735-8743.
- 587 Brook, B.W., Ellis, E.C., Perring, M.P., Mackay, A.W., Blomqvist, L., 2013. Does the terrestrial
588 biosphere have planetary tipping points? *Trends in Ecology & Evolution* 28, 396-401.
- 589 Brunner, P., Rechberger, H.H., 2001. Anthropogenic metabolism and environmental legacies.
590 Ed: Munn, T. *Encyclopedia of Global Environmental Change*. Wiley, Chichester, UK.
591 pp. 54-72
- 592 C40 Cities, 2013. C40 Cities: Climate Leadership Group. www.c40.org.
- 593 Carpenter, S.R., Bennett, E.M., 2011. Reconsideration of the planetary boundary for phosphorus.
594 *Environmental Research Letters* 6, DOI: 10.1088/1748-9326/6/1/014009
- 595 Clean Production Action. 2015. GreenScreen© for Safer Chemicals.
596 <http://www.greenscreenchemicals.org/>

- 597 Conklin, J., 2005. Dialogue Mapping: Building Shared Understanding of Wicked Problems.
598 Wiley, West Sussex, UK. 266 pp.
- 599 Csiszar, S.A., Daggupaty, S.M., Diamond, M.L., 2013. SO-MUM: a coupled atmospheric
600 transport and multimedia model used to predict intraurban-scale PCB and PBDE
601 emissions and fate. *Environmental Science & Technology* 47, 436-445.
- 602 Diamond, M.L., Melymuk, L., Csiszar, S.A., Robson, M., 2010. Estimation of PCB stocks,
603 emissions, and urban fate: will our policies reduce concentrations and exposure?
604 *Environmental Science & Technology* 44, 2777-2783.
- 605 Diaz, R.J., Rosenberg, R., 2008. Spreading dead zones and consequences for marine ecosystems.
606 *Science* 321, 926-929.
- 607 Doney, S.C., Fabry, V.J., Feely, R.A., Kleypas, J.A., 2009. Ocean acidification: the other CO₂
608 problem. *Annual Review of Marine Science* 1, 169-192.
- 609 Doney, S.C., Bopp, L., Long, M.C., 2014. Historical and future trends in ocean climate and
610 biogeochemistry. *Oceanography* 27, 108-119.
- 611 ECA (European Chemicals Agency), 2013. Pre-registered substances.
612 <http://echa.europa.eu/en/information-on-chemicals/pre-registered-substances>. Accessed
613 May 6, 2013.
- 614 EFSA (European Food Safety Authority), 2013. EFSA identifies risks to bees from
615 neonicotinoids. European Food Safety Authority.
616 <http://www.efsa.europa.eu/en/press/news/130116.htm>
- 617 Egeghy, P.P., Judson, R., Gangwal, S., Mosher, S., Smith, D., Vail, J., Hubal, E.A.C., 2012. The
618 exposure data landscape for manufactured chemicals. *Science of the Total Environment*
619 414, 159-166.

- 620 Ellerman, A.D., Sue Wing, I., 2003. Absolute vs. intensity-based emission caps. *Global Change*
621 *Science and Policy*, MIT. MIT, Cambridge MA, 11 pp.
- 622 Erisman, J.W., Galloway, J.N., Seitzinger, S., Bleeker, A., Dise, N.B., Petrescu, A.M.R., Leach,
623 A.M., de Vries, W., 2013. Consequences of human modification of the global nitrogen
624 cycle. *Philosophical Transactions of the Royal Society B: Biological Sciences* 368.
- 625 Fries, E., Mihajlovic, I., 2011. Pollution of soils with organophosphorus flame retardants and
626 plasticizers. *Journal of Environmental Monitoring* 13, 2692-2694.
- 627 Fromme, H., Kuchler, T., Otto, T., Pilz, K., Muller, J., Wenzel, A., 2002. Occurrence of
628 phthalates and bisphenol A and F in the environment. *Water Research* 36, 1429-1438.
- 629 Gawor, A., Shunthirasingham, C., Hayward, S.J., Lei, Y.D., Guin, T., Mmereki, B.T.,
630 Masamba, W., Ruepert, C., Castillo, L.E., Shoeib, M., Lee, S.C., Harner, T., Wania, F.,
631 2014. Neutral polyfluoroalkyl substances in the global atmosphere. *Environmental*
632 *Science & Process Impacts* 16, 404-413.
- 633 Gee, D., 2006. Late lessons from early warnings: Toward realism and precaution with endocrine-
634 disrupting substances. *Environmental Health Perspectives* 114, 152-160.
- 635 Gee, D., Grandjean, P., Hansen, S.F., Hove, S.V.D., MacGarvin, M., Martin, J., Nielsen, G.,
636 Quist, D., Stanners, D., 2013. *Late Lessons from Early Warnings: Science, Precaution,*
637 *Innovation*. EEA (European Environment Agency), Luxembourg. 760 pp.
- 638 Gerten, D., Hoff, H., Rockström, J., Jaegermeyr, J., Kummu, M., Pastor, A.V., 2013. Towards a
639 revised planetary boundary for consumptive freshwater use: role of environmental flow
640 requirements. *Current Opinion in Environmental Sustainability* 5, 551-558.

- 641 Gioia, R., Eckhardt, S., Breivik, K., Jaward, F.M., Prieto, A., Nizzetto, L., Jones, K.C., 2011.
642 Evidence for major emissions of PCBs in the West African region. *Environmental*
643 *Science & Technology* 45, 1349-1355.
- 644 Gottesfeld, P., 2013. The West's toxic hypocrisy over lead paint. *New Scientist* 218, 26-27.
- 645 Gottesfeld, P., Cherry, C.R., 2011. Lead emissions from solar photovoltaic energy systems in
646 China and India. *Energy Policy* 39, 4939-4946.
- 647 Grandjean, P., Landrigan, P.J., 2006. Developmental neurotoxicity of industrial chemicals.
648 *Lancet* 368, 2167-2178.
- 649 Harremoës, P., 1998. The challenge of managing water and material balances in relation to
650 eutrophication. *Water Science & Technology* 37, 9-17.
- 651 Harremoës, P., Gee, D., MacGarvin, M., Stirling, A., Keys, J., Wynne, B., Vaz, S.G., 2001. Late
652 Lessons from Early Warnings: The Precautionary Principle 1896-2000. EEA (European
653 Environment Agency), Copenhagen. 211 pp.
- 654 Henry, M., Beguin, M., Requier, F., Rollin, O., Odoux, J.F., Aupinel, P., Aptel, J., Tchamitchian,
655 S., Decourtye, A., 2012. A common pesticide decreases foraging success and survival in
656 honey bees. *Science* 336, 348-350.
- 657 Hooper, D.U., Adair, E.C., Cardinale, B.J., Byrnes, J.E.K., Hungate, B.A., Matulich, K.L.,
658 Gonzalez, A., Duffy, J.E., Gamfeldt, L., O'Connor, M.I., 2012. A global synthesis reveals
659 biodiversity loss as a major driver of ecosystem change. *Nature* 486, 105-U129.
- 660 Horn, R.E., Weber, R.P., 2007. New tools for resolving wicked problems: mess mapping and
661 resolution mapping processes. MacroVU, Inc and Stanford University; Strategy Kinetics,
662 LLC, Stanford, CA; Watertown, MA. 31 pp.

- 663 Howard, P.H., Muir, D.C.G., 2010. Identifying new persistent and bioaccumulative organics
664 among chemicals in commerce. *Environmental Science & Technology* 44, 2277-2285.
- 665 Howard, P.H., Muir, D.C.G., 2011. Identifying new persistent and bioaccumulative organics
666 among chemicals in commerce II: pharmaceuticals. *Environmental Science &*
667 *Technology* 45, 6938-6946.
- 668 IEA (International Energy Agency), 2014. *World Energy Outlook 2014*. OECD (Organization
669 for Economic Cooperation and Development), Paris. 748 pp.
- 670 Klanova, J., Cupr, P., Holoubek, I., Boruvkova, J., Pribylova, P., Kares, R., Tomsej, T., Ocelka,
671 T., 2009. Monitoring of persistent organic pollutants in Africa. Part 1: passive air
672 sampling across the continent in 2008. *Journal of Environmental Monitoring* 11, 1952-
673 1963.
- 674 Kolpin, D.W., Furlong, E.T., Meyer, M.T., Thurman, E.M., Zaugg, S.D., Barber, L.B., Buxton,
675 H.T., 2002. Pharmaceuticals, hormones, and other organic wastewater contaminants in
676 US streams, 1999-2000: A national reconnaissance. *Environmental Science &*
677 *Technology* 36, 1202-1211.
- 678 Kortenkamp, A., 2007. Ten years of mixing cocktails: a review of combination effects of
679 endocrine-disrupting chemicals. *Environmental Health Perspectives* 115, 98-105.
- 680 Kortenkamp, A., 2008. Low dose mixture effects of endocrine disruptors: implications for risk
681 assessment and epidemiology. *International Journal of Andrology* 31, 233-237.
- 682 Kortenkamp, A., Faust, M., Scholze, M., Backhaus, T., 2007. Low-level exposure to multiple
683 chemicals: reason for human health concerns? *Environmental Health Perspectives* 115,
684 106-114.

- 685 Kortenkamp, A., Backhaus, T., Faust, M., 2009. State of the Art Report on Mixture Toxicity.
686 Commission of the European Union (Directorate General for the Environment), London.
687 391 pp.
- 688 Lenton, T.M., Held, H., Kriegler, E., Hall, J.W., Lucht, W., Rahmstorf, S., Schellnhuber, H.J.,
689 2008. Tipping elements in the Earth's climate system. Proceedings of the National
690 Academy of Sciences 105, 1786-1793.
- 691 Lenzen, M., Moran, D., Kanemoto, K., Foran, B., Lobefaro, L., Geschke, A., 2012. International
692 trade drives biodiversity threats in developing nations. Nature 486, 109-112.
- 693 Letcher, R.J., Bustnes, J.O., Dietz, R., Jenssen, B.M., Jorgensen, E.H., Sonne, C., Verreault, J.,
694 Vijayan, M.M., Gabrielsen, G.W., 2010. Exposure and effects assessment of persistent
695 organohalogen contaminants in arctic wildlife and fish. Science of the Total Environment
696 408, 2995-3043.
- 697 Mani, M., Markandya, A., Sagar, A., Strukova, E., 2012. An Analysis of Physical and Monetary
698 Losses of Environmental Health and Natural Resources in India. The World Bank, South
699 Asia Region, Disaster Risk Management and Climate Change, Washington DC. 41 pp.
- 700 May, R.M., 1977. Thresholds and breakpoints in ecosystems with a multiplicity of stable states.
701 Nature 269, 471-477.
- 702 Meek, M.E., Boobis, A.R., Crofton, K.M., Heinemeyer, G., Van Raaij, M., Vickers, C., 2011.
703 Risk Assessment of Combined Exposure to Multiple Chemicals: A WHO/IPCS
704 framework. Regulatory Toxicology & Pharmacology 60, S1-S14.
- 705 Muir, D.C.G., Howard, P.H., 2006. Are there other persistent organic pollutants? A challenge for
706 environmental chemists. Environmental Science & Technology 40, 7157-7166.

- 707 OECD (Organization for Economic Cooperation and Development), 1991. Core Set of Indicators
708 for Environmental Performance Reviews, a Synthesis Report by the Group on the State of
709 the Environment. OECD, Paris.
- 710 Oaks, J.L., Gilbert, M., Virani, M.Z., Watson, R.T., Meteyer, C.U., Rideout, B.A., Shivaprasad,
711 H.L., Ahmed, S., Chaudhry, M.J.I., Arshad, M., Mahmood, S., Ali, A., Khan, A.A., 2004.
712 Diclofenac residues as the cause of vulture population decline in Pakistan. *Nature* 427,
713 630-633.
- 714 Peck, A.M., Hornbuckle, K.C., 2004. Synthetic musk fragrances in Lake Michigan.
715 *Environmental Science & Technology* 38, 367-372.
- 716 Persson, L.M., Breitholtz, M., Cousins, I.T., de Wit, C.A., MacLeod, M., McLachlan, M.S.,
717 2013. Confronting unknown planetary boundary threats from chemical pollution.
718 *Environmental Science & Technology* 47, 12619-12622.
- 719 Rockström, J., Steffen, W., Noone, K., Persson, A., Chapin, F.S., Lambin, E., Lenton, T.M.,
720 Scheffer, M., Folke, C., Schellnhuber, H.J., Nykvist, B., de Wit, C.A., Hughes, T., van
721 der Leeuw, S., Rodhe, H., Sorlin, S., Snyder, P.K., Costanza, R., Svedin, U., Falkenmark,
722 M., Karlberg, L., Corell, R.W., Fabry, V.J., Hansen, J., Walker, B., Liverman, D.,
723 Richardson, K., Crutzen, P., Foley, J., 2009. Planetary boundaries: exploring the safe
724 operating space for humanity. *Ecology and Society* 14. [online] URL:
725 <http://www.ecologyandsociety.org/vol14/iss2/art32/>
- 726 Rockström, J., Steffen, W., Noone, K., Persson, A., Chapin, F.S., Lambin, E.F., Lenton, T.M.,
727 Scheffer, M., Folke, C., Schellnhuber, H.J., Nykvist, B., de Wit, C.A., Hughes, T., van
728 der Leeuw, S., Rodhe, H., Sorlin, S., Snyder, P.K., Costanza, R., Svedin, U., Falkenmark,
729 M., Karlberg, L., Corell, R.W., Fabry, V.J., Hansen, J., Walker, B., Liverman, D.,

- 730 Richardson, K., Crutzen, P., Foley, J.A., 2009. A safe operating space for humanity.
731 Nature 461, 472-475.
- 732 Rosi-Marshall, E.J., Kincaid, D.W., Bechtold, H.A., Royer, T.V., Rojas, M., Kelly, J.J., 2013.
733 Pharmaceuticals suppress algal growth and microbial respiration and alter bacterial
734 communities in stream biofilms. Ecological Applications 23, 583-593.
- 735 SAICM (The Strategic Approach to International Chemicals Management), 2006. Texts and
736 Resolutions of the International Conference on Chemicals Management in 2006. United
737 Nations Environment Programme and the World Health Organization, Geneva,
738 Switzerland. 125 pp.
- 739 Scheringer, M., Stempel, S., Hukari, S., Ng, C.A., Blepp, M., Hungerbühler, K., 2012. How
740 many persistent organic pollutants should we expect? Atmospheric Pollution Research 3,
741 383-391.
- 742 Schmidt, C.W., 2006. Unfair trade - E-waste in Africa. Environmental Health Perspectives 114,
743 A232-A235.
- 744 SCHENIHR (Scientific Committee on Emerging and Newly Identified Health Risks), SCCS
745 (Scientific Committee on Consumer Safety), SCHER (Scientific Committee on Health
746 and Environmental Risks), 2012. Addressing the New Challenges for Risk Assessment.
747 European Commission, DG Health & Consumers, Directorate D: Public Health Systems
748 and Products, Unit D3 – Risk Assessment, Brussels. 157 pp.
- 749 Sindiku, O., Babayemi, J., Osibanjo, O., Schlummer, M., Schleup, M., Watson, A., Weber, R.
750 2014. Polybrominated diphenyl ethers listed as Stockholm Convention POPs, other
751 brominated flame retardants and heavy metals in e-waste polymers in Nigeria.
752 Environmental Science & Pollution Research DOI: 10.1007/s11356-014-3266-0.

- 753 Skelton, A., Guan, D., Peters, G.P., Crawford-Brown, D., 2011. Mapping Flows of Embodied
754 Emissions in the Global Production System. *Environmental Science & Technology* 45,
755 10516-10523.
- 756 Steffen, W., Richardson, K., Rockström, J., Cornell, S.E., Fetzer, I., Bennett, E.M., Biggs, R.,
757 Carpenter, S.R., de Vries, W., de Wit, C., Folke, C., Gerten, D., Heinke, J., Mace, G.M.,
758 Persson, L.M., Ramanathan, V., Reyers, B., Sörlin, S. 2015. Planetary boundaries:
759 guiding human development on a changing planet. *Science* DOI:
760 10.1126/science.1259855.
- 761 Stillerman, K.P., Mattison, D.R., Giudice, L.C., Woodruff, T.J., 2008. Environmental exposures
762 and adverse pregnancy outcomes: a review of the science. *Reproductive Sciences* 15,
763 631-650.
- 764 Stockholm Convention, 2008. Convention text.
765 http://ozone.unep.org/new_site/en/Treaties/treaties_decisions-hb.php?sec_id=5
- 766 Stockholm Convention, 2012. Success Stories Stockholm Convention 2001-2011. UNEP (United
767 Nations Environment Programme), Geneva. 169 pp.
- 768 Strepel, S., Scheringer, M., Ng, C.A., Hungerbühler, K., 2012. Screening for PBT chemicals
769 among the “existing” and “new” chemicals of the EU. *Environmental Science &*
770 *Technology* 46, 5680–5687.
- 771 Tapparo, A., Marton, D., Giorio, C., Zanella, A., Solda, L., Marzaro, M., Vivan, L., Girolami,
772 V., 2012. Assessment of the environmental exposure of honeybees to particulate matter
773 containing neonicotinoid insecticides coming from corn coated seeds. *Environmental*
774 *Science & Technology* 46, 2592-2599.

- 775 Torres, J.P.M., Froes-Asmus, C.I.R., Weber, R., Vijgen, J.M.H., 2013. HCH contamination from
776 former pesticide production in Brazil- A challenge for the Stockholm Convention
777 implementation. *Environmental Science & Pollution Research* 20, 1951-1957.
- 778 Turk, M., Jaksic, J., Miloradov, M.V., Klanova, J., 2007. Post-war levels of persistent organic
779 pollutants (POPs) in air from Serbia determined by active and passive sampling methods.
780 *Environmental Chemistry Letters* 5, 109-113.
- 781 UNECE (United Nations Economic Commission for Europe), 2004. Handbook for the 1979
782 Convention on Long-Range Transboundary Air Pollution and its Protocols. New York
783 and Geneva. United Nations Publication, ISBN 92-1-116895-3.
- 784 UNEP (United Nations Environment Programme), 2010-2011. The Montreal Protocol on
785 Substances that Deplete the Ozone Layer. Ozone Secretariat.
786 http://ozone.unep.org/new_site/en/Treaties/treaties_decisions-hb.php?sec_id=5
- 787 UNEP (United Nations Environment Programme), 2012. Global Environmental Outlook 5:
788 Environment for the future we want. Ed. UNEP, Nairobi.
- 789 UNEP (United Nations Environment Programme), 2013. GCO Global Chemicals Outlook:
790 Towards sound management of chemicals. Ed. United Nations Environment Programme,
791 Nairobi, Kenya.
- 792 UNEP (United Nations Environment Programme), 2015. Minamata Convention Agreed by
793 Nations. <http://www.mercuryconvention.org/>
- 794 UNEP (United Nations Environment Programme), WHO (World Health Organisation), CVUA
795 (The State Institute for Chemical and Veterinary Analysis of Food), MTM Research
796 Centre, 2013. Human exposure to POPs across the globe: POPs levels and human health

- 797 implications. Results of the WHO/UNEP human milk survey. WHO; UNEP; Stockholm
798 Convention, Geneva; Nairobi; Châtelaine.
- 799 U.S. EPA (United States Environmental Protection Agency), 2014. Design for the Environment.
800 <http://www.epa.gov/dfe/>
- 801 UNFCCC, 2013. United Nations Framework Convention on Climate Change.
802 <http://unfccc.int/2860.php>.
- 803 Voeroesmarty, C.J., McIntyre, P.B., Gessner, M.O., Dudgeon, D., Prusevich, A., Green, P.,
804 Glidden, S., Bunn, S.E., Sullivan, C.A., Liermann, C.R., Davies, P.M., 2010. Global
805 threats to human water security and river biodiversity. *Nature* 467, 555-561.
- 806 Wang, D.G., Norwood, W., Alaei, M., Byer, J.D., Brimble, S., 2013. Review of recent advances
807 in research on the toxicity, detection, occurrence and fate of cyclic volatile methyl
808 siloxanes in the environment. *Chemosphere* 93, 711-725.
- 809 Weber, R., Watson, A., Forter, M., Oliaei, F., 2011. Persistent organic pollutants and landfills - a
810 review of past experiences and future challenges. *Waste Management & Research* 29,
811 107-121.
- 812 Weber, R., Aliyeva, G., Vijgen, J., 2013. The need for an integrated approach to the global
813 challenge of POPs management. *Environmental Science & Pollution Research* 20, 1901-
814 1906.
- 815 Whitehorn, P.R., O'Connor, S., Wackers, F.L., Goulson, D., 2012. Neonicotinoid pesticide
816 reduces bumble bee colony growth and queen production. *Science* 336, 351-352.
- 817 WHO (World Health Organisation), UNEP (United Nations Environment Programme), 2013.
818 State of the science of endocrine disrupting chemicals 2012. United Nations Environment
819 Programme and the World Health Organization, Geneva, Switzerland, 260 pp.

820 Wilson, M.P., Schwarzman, M.R., 2009. Toward a new U.S. chemicals policy: rebuilding the
821 foundation to advance new science, green chemistry, and environmental health.
822 Environmental Health Perspectives 117, 1202-1209.

835

836

837

838

Supplementary information

839

840 Exploring the planetary boundary for chemical pollution

841

842 Miriam L. Diamond^{†*}, Cynthia A. de Wit[‡], Sverker Molander,[§] Martin Scheringer,[%] Thomas843 Backhaus,^{||} Rainer Lohmann,[√] Rickard Arvidsson,[§] Åke Bergman,[⊥] Michael Hauschild,[#] Ivan844 Holoubek,[¶] Linn Persson,[&] Noriyuki Suzuki,[@] Marco Vighi,[□] Cornelius Zetzsch^Δ

845

846 [†] Department of Earth Sciences, University of Toronto, 22 Russell Street, Toronto, M5S 3B1

847 Ontario, Canada

848 [‡] Department of ~~Applied~~-Environmental Science and Analytical Chemistry (ATMACES),

849 Stockholm University, SE-106 91 Stockholm, Sweden

850 [§] Environmental Systems Analysis, Department of Energy and Environment, Chalmers

851 University of Technology, SE-412 96 Gothenburg, Sweden

852 [%] Institute for Chemical and Bioengineering, ETH Zürich, Wolfgang-Pauli-Str. 10, 8093 Zürich,853 CH-8093, Switzerland, and Leuphana University Lüneburg, D-21335 Lüneburg, Germany854 ^{||} Department of Biological and Environmental Sciences, University of Gothenburg, Box 100,

855 SE-405 30 Gothenburg, Sweden

856 [√] Graduate School of Oceanography, University of Rhode Island, South Ferry Road,

857 Narragansett, Rhode Island, 02882, United States

858 [†]Department of Materials and Environmental Chemistry, Stockholm University, SE-106 91

859 Stockholm, Sweden

860 [#]Department of Management Engineering, Technical University of Denmark (DTU), Nils

861 Koppels Allé, Building 426 D, DK-2800 Kgs. Lyngby, Denmark

862 [‡]Research Centre for Toxic Compounds in the Environment (RECETOX), Faculty of Science,

863 Masaryk University, Kamenice 753/5, 625 00 Brno, Czech Republic

864 [&]Stockholm Environment Institute, Linnégatan 87D, Box 24218, Stockholm, Sweden

865 [@]Strategic Risk Management Research Section, Center for Environmental Risk Research,

866 National Institute for Environmental Studies, 16-2 Onogawa, Tsukuba, Ibaraki 305-8506, Japan

867 [‡]Department of Earth and Environmental Sciences, University of Milano Bicocca, Piazza della

868 Scienza 1, Milan, 20126 Italy

869 [^]Forschungsstelle für Atmosphärische Chemie, Dr. Hans-Frisch-Str. 1-3, Universität Bayreuth,

870 D-954 48 Bayreuth, Germany

871

872

873

874

875 13 pages, two tables and text

Formatted: Font color: Auto

Table S1. Examples of regulations addressing the occurrence of chemicals in the environment or the human body that establish boundaries for chemical pollution. Regulations are listed according to the type of boundary used: risk-based, concentration-based, emissions-based, technology-driven.

4	Issuing organization and year of entry into force	Chemicals covered	Boundary type	Spatial scale	Protection goal
Acceptable Daily Intake (ADI),	World Health Organization (WHO), Food and Agriculture Organization of the United Nations (FAO), 1961	food additives, veterinary pharmaceuticals and pesticide residues in food	<i>risk-based</i> a lifelong daily uptake below the ADI is considered safe	global human population	human health
Tolerable Daily Intake (TDI)	WHO and Joint FAO/WHO Expert Committee on Food Additives (JECFA), 1961	non-intentionally used xenobiotics in food	<i>risk-based</i> a lifelong daily uptake below the TDI is considered safe	global human population	human health
Provisional Tolerable Weekly Intake (PTWI)	JECFA	non-intentionally used xenobiotics in food that may accumulate in the human body	<i>risk-based</i> a lifelong weekly uptake below the PTWI is considered safe	global human population	human health
Reference Dose (RfD)	US Environmental Protection Agency	toxic chemicals in general	<i>risk based</i> the RfD provides an estimate of the lifelong daily oral exposure to the human population that is likely to be without an appreciable risk of deleterious effects during a lifetime.	US population	human health

Maximum Residue Levels (MRL)	Regulation (EC) 396/2005, 2008	pesticides in food	<i>risk based, technology based</i> the upper legal level of a concentration for a pesticide residue in or on food or feed set in accordance with this Regulation, based on good agricultural practice and the lowest consumer exposure necessary to protect vulnerable consumers	European population	human health
Critical loads and levels	United Nations Economic Commission for Europe (UN ECE) Convention on Long-range Transboundary Air Pollution (LRTAP), 1981	major air pollutants (e.g. SO _x , NO _x)	<i>risk-based</i> a maximum permissible load of a chemical below which no harmful effects occur in an exposed ecosystem	ecosystem (local, regional)	environment
Toxicity Exposure Ratio (TER)	Council Directive 91/414/EEC, which has just been repealed by Regulation (EC) No 1107/2009 (21.10.2009)	pesticides (active ingredients and formulated products)	<i>risk-based</i> a TER above a pre-define threshold is considered safe	ecosystem (local, regional)	environment
Environmental Quality Standards (EQS) and Maximum allowable concentrations (MAC)	Water Framework Directive (WFD), Directive 2000/60/EC, Directive 2008/105/EC on Environmental Quality Standards, Directive on priority substances (2008/105/EC) Oct. 2000	priority pollutants detected in water bodies	<i>risk-based</i> EQS: a level providing protection against long-term exposure, and MAC: protection against short-term exposure	ecosystem (local, regional)	environment (water bodies only)
Environmental quality standards (EQS)	Ministry of the Environment, Government of Japan, for water pollution under basic Environment Law of Japan (Established in 1968, last amendment in 2014) http://www.env.go.jp/en/water/wq/wp.pdf	Substances relating to human health and living environment	<i>risk-based</i> EQS: a level providing protection against long-term exposure, and MAC: protection against short-term exposure	Japanese population and ecosystem (local, regional)	Human health and environment (water bodies only)
Predicted No Effect Concentrations (PNEC)	Regulation EC 1907/2006 (REACH) 1.6. 2007.	industrial chemicals in water, air, soil, sediment	<i>risk-based:</i> a concentration below the PNEC is considered safe	local, regional	environment
Derived No Effect Level (DNEL)	Regulation EC 1907/2006 (REACH) 1.6. 2007.	industrial chemicals	<i>risk-based:</i> a concentration below the DNEL is considered safe	European human population	human health

Zero discharges, emissions and losses of hazardous substances	The Convention for the Protection of the marine Environment of the North-East Atlantic (OSPAR Convention), 1998	hazardous chemicals	<i>concentration-based</i> : concentration of zero for artificial chemicals and concentration at natural background levels for naturally occurring chemicals	regional (north-east Atlantic)	environment (marine ecosystems only)	
<u>Emission Values (ELVs)</u>	<u>Limit Values (ELVs)</u>	<u>Ministry of the Environment, Government of Japan, Regulatory Measures against Air Pollutants Emitted from Factories and Business Sites and the Outline of Regulation (last amended in 1998).</u> http://www.env.go.jp/en/air/aq/air.html	<u>Air pollutants that may affect human health and environment</u>	<u>risk-based ELVs and other regulatory measures for factories and business sites are adopted to achieve EQSs to protect human health and environment</u>	<u>local</u>	<u>human health and environment</u>
Action limits	Guideline of the European Medicines Agency (EMA) on the environmental risk assessment of medicinal products for human use (EMA/CHMP/SWP/4447/00)	human pharmaceuticals	<i>concentration-based</i> concentration below 0.01 µg/l in surface waters are considered inherently safe, unless specific reasons for concern are given (e.g. endocrine activity).	local, regional	environment	
Threshold of toxicological concern (TTC) Threshold of Regulation	EMA Guideline on the limits of genotoxic impurities (EMA/CHMP/QWP/251344/2006), 2006	genotoxic impurities in pharmaceuticals food contact materials	<i>concentration-based</i> the TTC defines a common exposure level (1.5µg/day) for an unstudied chemical that will not pose a risk of “significant carcinogenicity or other toxic effects”.	European human population	human health	
Threshold of Regulation (TOR)	US Food and Drug Administration (FDA), Code of Federal Regulation (CFR), 21, § 170.39	food contact materials	<i>concentration-based</i> Concentrations of ≤ 0.5 ppb (corresponding to dietary exposure levels ≤ 1.5 µg/(person*day)) are considered safe.	US human population	human health	
Maximum Contaminant Level (MCL), maximum contaminant level goals (MCLGs) and	Safe Drinking Water Act (SDWA), enforced by US EPA	contaminants in drinking water	<i>concentration, risk and technology based</i> MCLG: The level of a contaminant in drinking water below which there is no known or	US human population	human health	

Practical Quantitation Limit (PQL)			expected risk to health. MCLGs allow for a margin of safety and are non-enforceable public health goals. MCL describe the highest level of a contaminant that is allowed in drinking water. MCLs are set as close to MCLGs as feasible using the best available treatment technology and taking cost into consideration. MCLs are enforceable standards. For non-carcinogens, MCLGs levels for drinking water are established based on the RfD, average drinking water consumption, etc. For carcinogens the MCLG is set to zero, which is practically ensured by checking whether a contaminant is present above the PQL.		
<u>Canadian Environmental Quality Guidelines</u>	<u>Canadian Council of Ministers of the Environment</u>	<u>VOCs, SVOCs and metals</u>	<u>Concentration-based: chemical specific goals (non-enforceable) for protection of aquatic life, protection of soil quality, protection of groundwater at contaminated sites, protection of environmental and human health.</u>	<u>national</u>	<u>human health and environment</u>
<u>Canadian “tolerances” and “standards” for various chemical contaminants in food</u>	<u>Health Canada Food Directorate</u>	<u>specified chemicals</u>	<u>Concentration-based: Maximum concentrations expressed as tolerances (through regulation) and standards (not regulated) for listed chemicals.</u>	<u>national</u>	<u>human health</u>
<u>Environmental Standards for ambient air and</u>	<u>Ministry of Environment, Forest and climate change, Government of India</u> <u>Reference:</u>	<u>specified chemicals and parameters</u>	<u>Concentration based: chemical or parameter specific goals for protection of environmental and</u>	<u>national</u>	<u>human health and environment</u>

Formatted: Font: Not Italic

Formatted: Font: Not Italic

<u>water quality criteria</u>	<u>http://envfor.nic.in/environmental_standards</u>		<u>human health, protection of aquatic life and water resources.</u>			
	Montreal Protocol, 1989	CFCs	<u>emission-based</u> : production has to reach zero.	global	ozone layer; human health <u>and</u> environment	Formatted: Font: Italic, No underline, Font color: Auto
	Stockholm Convention, 2004	POPs	<u>emission-based</u> : production and use have to reach zero.	global	human health <u>and</u> environment	Formatted: Font: Italic, No underline, Font color: Auto
<u>Schedule I Compounds</u>	<u>Canadian Environmental Protection Act (CEPA) 1999</u>	<u>chemicals deemed "CEPA toxic"</u>	<u>emission-based: limits on production, use and importation of chemicals listed in Schedule 1.</u>	<u>national</u>	<u>human health and/or environment</u>	Formatted: Font: Not Italic
Emission Limit Values (ELVs)	Directive 2008/1/EC concerning integrated pollution prevention and control (IPPC directive), 2008	chemicals produced at a given site	<u>technology-based</u> : ELVs are part of the permit an installation needs to acquire, based on the best available techniques (BAT), as defined in the Directive, and also taking specific local conditions into account. ELVs "should lay down provisions on minimising long-distance or transfrontier pollution and ensure a high level of protection for the environment as a whole". Emissions are regarded in an integrated manner in order to avoid switching from one compartment to another.	local	human health <u>and</u> environment	
<u>National Environmental Regulations</u>	<u>National Environmental Standards and Regulations Enforcement Agency (NESREA) Nigeria Federal Republic of Nigeria Official Gazette No. 92, Vol. 94 of 31st July, 2007.</u>	<u>specified chemicals and parameters</u>	<u>concentration, emission and risk based: Enforceable maximum concentrations, emission limits and tolerance limits for specified chemicals</u>	<u>national</u>	<u>human health and environment</u>	
Environmental Protection Law of the People's	Ministry of Environmental Protection	general environmental protection issues	<u>Concentrations and emission-based</u> : prevention and control of water and air pollution;	national	<u>h</u> Human health and environment	

Republic of China (Regulations, and laws)			management in solid waste, marine environment, hazardous chemicals; pollution discharge and levying; environmental standards and monitoring.		
Environmental Protection Law of the People's Republic of China (Environmental Standards)	Ministry of Environmental Protection	specified chemicals and parameters	<i>Concentration and/technology based</i> : protection of water, air, soil, and eco-environment.	national	Human health and environment

PEC: predicted environmental concentration

PNEC: predicted no-effect concentration

DNEL: derived no-effect concentration

BAT: best available technology

Indicators

Environmental management schemes employ indicators as metrics that allow evaluation of the status of an environmental system that is influenced by human activities (OECD, 1991a; Gallopín, 1996; Harremoës, 1998). In the context of a planetary boundary (PB), the “control variable” is a type of indicator, linking human activities (that hopefully can change under a governance scheme) to a specific threshold – a tipping point - for some of the categories (like global warming) or, for other categories (like biodiversity), to a derived limit. Considering the challenges of establishing one or more planetary boundary/boundaries for chemical pollution (PBCP), precaution, warranted by uncertainties and/or knowledge gaps, can be integrated into the PB analysis by introducing an uncertainty range on the safe side of its defined limit.

Rather than defining a single indicator that can be directly related to a control variable, defining an “interim” indicator may be necessary. An example within the PB context is biodiversity that is addressed at a continental to global scale, since biodiversity loss depends on many factors rather than a single control variable and a single threshold may not exist (Schellnhuber, 2002; Rockström et al., 2009). Here, the present extinction rate is an “interim indicator” of the ultimate mean of long-term maintained biodiversity. A PB can then be obtained by relating the present extinction rate to the long-term mean extinction rate (Rockström et al., 2009).

The construction of indicators of planetary chemical pollution is a formidable task given the large set of difficulties in this particular case. As discussed in the text, one difficulty is the *very large number of specific chemical structures identified* and of potential concern - about 100 000

are expected to be on the market following the definitions used by European REACH-legislation. A second difficulty is the *widespread production, and inclusion of chemicals in manufacturing* of a very wide set of products, which are used and wasted in many different ways wherever humans are found. Globalized production chains and increasing human consumption underline the importance of this aspect. A third difficulty is that the *release of chemical substances* occurs along complex product chains during the life-cycles of the products. The emissions are influenced by a number of factors, including material composition, fragmentation of the product increasing the effective surface for release, and environmental factors like temperature, making only the determination of *emissions* a daunting task. A further difficulty is the *environmental distribution, transformation and transportation*, that all are complicated processes, continue after emission. These processes are influenced by many environmental factors spanning from temperature and light intensity to pH and the ability of (micro)organisms to transform, transport and degrade the substances.

Furthermore the *very large numbers of organisms, exposed under an overwhelming number of conditions, express a wide number of responses to chemicals*. (Eco)toxicologists have identified a huge number of such responses, on different levels of biological complexity, and are employing a large number of test species and measurement endpoints in order to cover the potential effects of chemicals on human health and the environment. Reconnecting to the huge number of chemicals, as mentioned above, these chemicals differ tremendously in their *potency* to exert a particular effect in a particular species.

It is on the combination of these aspects that indicators of planetary chemical pollution must act, giving a simplified, but still meaningful representation of the actual pollution situation.

Furthermore, the indicators must meet practical requirements: they must be unambiguously defined, their values must be measurable and data must be available or possible to gather, the method for acquisition, processing and presenting of values must be clear, transparent and standardized, and the means to do this must be available. Meeting these requirements would bring into focus the benefits and costs of indicators, and therefore their political acceptability and the process to establish them (Gallopín, 1996).

The perception of a simplified cause-effect-chain, along which environmental indicators can be identified, has dominated the development of such indicators since the first OECD State of the Environment report (OECD, 1991a). The DPSIR framework (*Driving forces-Pressures-States-Impacts-Responses*) was adopted for the European environmental indicators by the European Environmental Agency (Harremoës, 1998; Smeets and Weterings, 1999). A similar approach was also taken within life-cycle impact assessment methods (Udo de Haes et al., 1999), and considerable effort has been expended to developing sustainability indicators more or less along these lines (Meados, 1999; OECD, 1998; Bossel, 1999; Lundin et al., 1999; Parris and Kates, 2003; Palme et al., 2005; OECD, 1991b). Here we have adapted the DPSIR framework for the PBCP, placing currently existing indicators of chemical pollution within the DPSIR framework in order to illustrate possible indicators and further required development.

Table S2 suggests a framework for indicators of chemical pollution at different stages in a simplified cause-effect chain, inspired by the DPSIR-approach and applying proxy indicators

reflecting *exposure* and *potency* as the key aspects. The indicators suggested in Table S2 offer the possibility of moving from distant or indirect drivers of chemical pollution (like production or emissions) to more direct indicators of adverse effects. Another explanation of Table S2 begins with indicators that are proxies of exposure (production and emissions), to indicators of the control variable (exposure), to “interim indicators” where effects, which are connected to chemical potency, are identified. It is also possible to develop spatially dependent indicators (e.g. derived from indicators listed in Table S2) related to, for example, the proportion of land (or sea) area impacted by a certain degree of chemical pollution. Such an approach opens the application of GIS-based emission, fate and exposure modeling that is under development (Pistocchi et al., 2010).

Several existing global monitoring efforts of concentrations form an important step towards developing indicators that can be used to define a PBCP. These include monitoring efforts coordinated under the umbrella of the Stockholm Convention, such as the Global Atmospheric Sampling network or GAPS, the Arctic Monitoring and Assessment Programme or AMAP, the East Asia Air Monitoring Program. (Stockholm Convention and UNEP, 2008; Gawor et al., 2014) and the Human Milk Survey (Stockholm Convention and UNEP, 2008; Gawor et al., 2014; UNEP et al., 2013).

Defining a PBCP related to one or several of the suggested indicators is the next step. Here we suggest some possible indicators for control variables and some starting points for further scientific elaboration.

Table S2. Examples of indicators for chemical pollution at different stages along a generalized cause-effect chain for chemicals that can be further elaborated aiming for one or more indicator of PBCP. ADI is the Acceptable daily intake according to [the IPCS/WHO](#) (1987) and Renwick (1998). Toxic unit is the quotient of an actual concentration or intake of a substance and a determined effect measure (e.g. the EC50 or LC50) (Peterson, 1994). A critical volume is the volume of a medium (often water) needed to dilute an emitted mass of a substance to a concentration lower than the no-effect concentration of a representative species or group of species. Disability adjusted life-years (DALY) is an indicator of disease burden that can be connected to human chemical exposure (Murray and Lopez, 1996).

Indicator target	Drivers			Pressures	States		Impacts	
	Innovation	Production	Societal stock	Emission	Environmental concentration	Exposure	Effects	Damage
Single chemicals OR groups of chemicals OR all chemicals	Number of commercially available chemicals [dimensionless]	Annual production of single, groups of, or all chemicals [ton/year]	Current stock of single, groups of, or all chemicals [ton]	Annual emissions of single, groups of, or all chemicals [ton/year]	Environmental concentrations of single, groups of, or all chemicals [mol/l or kg/m ³]	(Potential) Daily intake (PDI) of single, groups of, or all chemicals [mg/kg bw/day] Concentration in tissue of single, groups of, or all chemicals [mg/kg]	PDI/ADI of single, groups of, or all chemicals [dimensionless] Toxic units [dimensionless] Critical volumes [m ³] Potentially Damaged Fraction of species (PDF) [PDF·m ² ·yr] Loss of population(s) or species [dimensionless]	An explicit and relative valuation of effect indicator(s) e.g. DALY [years]

1 **References**

- 2
- 3 Bossel, H., 1999. Indicators For Sustainable Development: Theory, Method, Applications. A
4 Report to The Balaton Group. IISD (International Institute for Sustainable Development),
5 Winnipeg, Canada.
- 6 Gallopín, G.C., 1996. Environmental and sustainability indicators and the concept of situational
7 indicators. A systems approach. *Environmental-Modeling and -Assessment* 1, 101-117.
- 8 Gawor, A., Shunthirasingham, C., Hayward, S.J., Lei, Y.D., Gouin, T., Mmereki, B.T.,
9 Masamba, W., Ruepert, C., Castillo, L.E., Shoeib, M., Lee, S.C., Harner, T., Wania, F.,
10 2014. Neutral polyfluoroalkyl substances in the global atmosphere. *Environmental
11 Science-Processes & Impacts* 16, 404-413.
- 12 Harremoës, P., 1998. The challenge of managing water and material balances in relation to
13 eutrophication. *Water Science & Technology* 37, 9-17.
- 14 IPCS (International Programme on Chemical Safety), 1987. Principles for the Safety Assessment
15 of Food Additives and Contaminants In Food - Environmental Health Criteria No 70.
16 WHO (World Health Organisation), Geneva.
- 17 Lundin, M., Molander, S., Morrison, G.M., 1999. A set of indicators for the assessment of
18 temporal variations in the sustainability of sanitary systems. *Water Science &
19 Technology* 39, 235-242.
- 20 Meadows, D.H., 1998. Indicators and Information Systems for Sustainable Development: A
21 Report to The Balaton Group. The Sustainability Institute. Hartland Four Corners, VT.
- 22 Murray, C.J.L., Lopez, A.D., 1996. The Global Burden of Disease. Harvard University Press;
23 WHO, Boston, MA; Geneva.

- 24 OECD (Organisation for Economic Co-operation and Development), 1991a. State of the
25 Environment. OECD, Paris. 298 pp.
- 26 OECD (Organisation for Economic Co-operation and Development), 1991b. Core Set of
27 Indicators For Environmental Performance Reviews: A Synthesis Report By The
28 Group On The State Of The Environment. OECD, Paris.
- 29 OECD (Organisation for Economic Co-operation and Development), 1998. Towards Sustainable
30 Development - Environmental Indicators. OECD, Paris.
- 31 Palme, U., Lundin, M., Tillman, A.M., Molander, S., 2005. Sustainable development indicators
32 for wastewater systems - researchers and indicator users in a co-operative case study.
33 Resources Conservation & Recycling 43, 293-311.
- 34 Parris, T.M., Kates, R.W., 2003. Characterizing and measuring sustainable development. Annual
35 Review of Environment & Resources 28, 559-586.
- 36 Peterson, D.R., 1994. Calculating the aquatic toxicity of hydrocarbon mixtures. Chemosphere 29,
37 2493-2506.
- 38 Pistocchi, A., Sarigiannis, D.A., Vizcaino, P., 2010. Spatially explicit multimedia fate models for
39 pollutants in Europe: State of the art and perspectives. Science of the Total Environment
40 408, 3817-3830.
- 41 Renwick, A.G., 1998. Toxicokinetics in infants and children in relation to the ADI and TDI.
42 Food Additives and Contaminants 15, 17-35.
- 43 Rockström, J., Steffen, W., Noone, K., Persson, A., Chapin, F.S., Lambin, E., Lenton, T.M.,
44 Scheffer, M., Folke, C., Schellnhuber, H.J., Nykvist, B., de Wit, C.A., Hughes, T., van
45 der Leeuw, S., Rodhe, H., Sorlin, S., Snyder, P.K., Costanza, R., Svedin, U., Falkenmark,
46 M., Karlberg, L., Corell, R.W., Fabry, V.J., Hansen, J., Walker, B., Liverman, D.,

- 47 Richardson, K., Crutzen, P., Foley, J., 2009. Planetary boundaries: Exploring the safe
48 operating space for humanity. *Ecology & Society* 14.- [online] URL:
49 <http://www.ecologyandsociety.org/vol14/iss2/art32/>
50 ~~35 pp.~~
51
52 Schellnhuber, H.J., 2002. Coping with Earth system complexity and irregularity. Eds: Steffen,
53 W., Jager, J., Carson, D.J., Bradshaw, C. *Challenges of a Changing Earth*, pp. 151-156.
54 Smeets, E., Weterings, R., 1999. *Environmental Indicators: Typology and Overview*. European
55 Environment Agency (EEA), Copenhagen. 19 pp.
56 Stockholm Convention and UNEP (United Nations Environment Programme). 2008. *Air*
57 *Monitoring*. Stockholm Convention.
58 [http://chm.pops.int/Implementation/GlobalMonitoringPlan/MonitoringActivities/Airmoni](http://chm.pops.int/Implementation/GlobalMonitoringPlan/MonitoringActivities/Airmonitoring/tabid/269/Default.aspx)
59 [toring/tabid/269/Default.aspx](http://chm.pops.int/Implementation/GlobalMonitoringPlan/MonitoringActivities/Airmonitoring/tabid/269/Default.aspx) .
60 Udo de Haes, H.A., Jolliet, O., Finnveden, G., Hauschild, M., Krewitt, W., Müller-Wenk, R.,
61 1999. Best available practice regarding impact categories and category indicators in life
62 cycle impact assessment. *International Journal of Life Cycle Assessment* 4, 167-174.
63 UNEP (United Nations Environment Programme), WHO (World Health Organisation), CVUA
64 (The State Institute for Chemical and Veterinary Analysis of Food), MTM Research
65 Centre, 2013. *Human Exposure to POPs Across the Globe: POPs Levels and Human*
66 *Health Implications. Results of the WHO/UNEP Human Milk Survey*. WHO; UNEP;
67 Stockholm Convention, Geneva; Nairobi; Châtelaine.

68

Formatted: Indent: Left: 0 cm,
Hanging: 1.27 cm, Line spacing:
Double